This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

ISTURBANCES
OF THE
HEART

LANE MEDICAL LIBRARY STANFORD
L661 .O61 1913 STOR
Disturbances of the heart : a discussion

LANE
MEDICAL LIBRARY
SEAL OF THE LELAND STANFORD JUNIOR UNIVERSITY
SEMPER VIRENS
1885

GIFT
DR.H.C.McCL_NAHAN

DISTURBANCES OF THE HEART

Discussion of the Treatment of the Heart in Its Various Disorders

EDITED BY

OLIVER T. OSBORNE, A.M., M.D.
Professor of Therapeutics and formerly Professor of Clinical Medicine in Yale Medical School; Member of the Council on Pharmacy and Chemistry; Author of "Introduction to Materia Medica and Pharmacology"; Editor of the "Handbook of Therapy"
NEW HAVEN, CONN.

* * *

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
Five Hundred Thirty-Five North Dearborn Street, Chicago

COPYRIGHT, 1913
BY THE AMERICAN MEDICAL ASSOCIATION

LANE LIBRARY

081
1913

PREFACE

That marvelous organ which, moment by moment and year by year, keeps consistently sending the blood on its path through the arteriovenous system is naturally one whose structure and function need to be carefully studied if one is to guard it when threatened by disease. This series of articles deals with heart therapy, not discussing the heart structurally and anatomically, but taking up in detail the various forms of the disturbances which may affect the heart. The cordial reception given by the readers of *The Journal* to this series of articles has warranted its issue in book form so that it may be slipped into the pocket for review at appropriate times, or kept on the desk for convenient reference.

54037

CONTENTS

PAGE

Preface 3

Heart Disturbances in General 7

Acute Pericarditis 17

Myocardial Disturbances 27

Endocarditis 38

Chronic Disease of the Valves 62

Acute Cardiac Symptoms: Acute Heart Attack 84

Diet and Baths in Heart Disease 114

Heart Disease in Children and During Pregnancy 123

Degenerations 130

Disturbances of the Heart Rate 169

Miscellaneous Disturbances 200

Index 213

conductivity will cause a heart-block, an intermittent disturbance will cause intermittent heart-block (Stokes-Adams disease), and a prolonged disturbance, death. It has also been shown that extrasystoles, meaning irregular heart action, may be caused by impulses originating at the apex, at the base or at some point in the right ventricle.

Anatomists and histologists have more perfectly demonstrated the muscle fibers of the heart and the structure at and around the valves; the physiologic chemists have shown more clearly the action of drugs, metals and organic solutions on the heart; and the physiologists and clinicians with laboratory facilities have demonstrated by various new apparatus the action of the heart and the circulatory power under various conditions. It is not now sufficient to state that the heart is acting irregularly, or that the pulse is irregular; the endeavor should be to determine what causes the irregularity, and what kind of irregularity is present.

CLINICAL INTERPRETATION OF PULSE TRACINGS

A moment may be spent on clinical interpretation of pulse-tracings. It has recently been shown that the permanently irregular pulse is due to fibrillary contraction, or really auricular fibrillation — in other words, irregular stimuli proceeding from the auricle — and that such an irregular pulse is not due to disturbance at the auriculoventricular node, as believed a short time ago. These little irregular stimuli proceeding from the auricle reach the auriculoventricular node and are transmitted to the ventricle as rapidly as the ventricle is able to react. Such rapid stimuli may soon cause death; or, if for any reason, medicinal or otherwise, the ventricle becomes indifferent to these stimuli, it may not take note of more than a certain portion of the stimuli. It then acts slowly enough to allow prolongation of life, and even considerable activity. If such a heart becomes more rapid from such stimuli, 110 or more, for any length of time, the condition becomes very serious. ? Digitalis in such a condition is, of course, of supreme value on account of its ability to slow the heart. Such irregularity perhaps most frequently occurs with valvular disease, especially mitral stenosis, and in the muscular degenerations of senility, as fibrosis.

Lewis[1] believes that 50 per cent. of cardiac arhythmia originates in muscle disturbance or incoordination in the auricle. These stimuli are irregular in intensity and the contractions caused are irregular in degree. If the wave lengths of the pulse-tracing show no regularity — if, in fact, hardly two adjacent wave-lengths are alike — the disturbance is auricular fibrillation. Injury to the auricle, or pressure for any reason on the auricle, may so disturb the transmission of stimuli and contractions that the contractions of the ventricle are very much fewer than the stimuli proceeding from the auricle. In other words, a sort of heart-block may occur. Various stimuli coming through the pneumogastric nerves, either from above or from the peripheral endings in the stomach or intestines, may inhibit or slow the ventricular contractions. It seems to have been again shown, as was earlier understood, that there are inhibitory and accelerator ganglia in the heart itself, each subject to various kinds of stimulation and various kinds of depression.

Of course it is generally understood that children have a higher pulse-rate than adults; that women normally have a higher pulse-rate than men at the same age; that strenuous muscular exercise, frequently repeated, without cardiac tire, while causing the pulse to be rapid at the time, slows the pulse during the interim of such exercise and may gradually cause a more or less permanent slow pulse. It should be clinically remembered that athletes have slow pulse, and the severity of their condition must not be interpreted by the rapidity of the pulse. Even with high fever the pulse of an athlete may be slow. It is well known that high blood-pressure slows the pulse-rate; that low blood-pressure generally increases the pulse-rate; that arteriosclerosis, or the gradual aging of the arteries, slows the pulse, except when the cardiac degeneration of old age makes the heart again more irritable and more rapid. The rapid heart in hyperthyroidism is also well understood. It is not so frequently noted that hypersecretion of the thyroid may cause a rapid heart without any other tangible or discoverable thyroid symptoms or symptoms of hyperthyroidism. Bile in the blood almost always slows the pulse.

---

[1] Lewis: Brit. Med. Jour., 1909, ii, 1528.

BLOOD-PRESSURE

As above suggested, new and better instruments have been invented, and laboratory and clinical studies of tracings of heart and vessels have increased our knowledge of the varying blood-waves and of the pressure under which the blood is distributed. The development of these instruments of precision started with the sphygmograph of Marey, in 1860. While the sphygmomanometer was discovered in the first third of the last century, an instrument was not devised or used clinically until perhaps first in 1887 — the instrument of von Basch. Soon followed Marey's and Potain's instruments, then Ludwig's suggestion of the graphic method of study of blood-pressures and pulse-tracings, then the finger-ring suggestion which is not very accurate, then the Riva Rocci blood-pressure apparatus and the modifications that followed, such as Cook's, Sahli's, von Recklinghausen's, Janeway's, Erlanger's, Satterthwaite's and Faught's, all based on the reading of the mercurial pressure, and the Tycos air instrument gauged to mercurial pressure.

From the careful clinical studies of blood-pressure it has been shown that the pressure varies greatly with the age, sex, different physical and mental conditions, position and with disease. The pressure in a child may be about 100 mm. of mercury, in a woman about 120, and in a man about 130. It is now recognized that accuracy requires the taking of diastolic as well as systolic pressure, and that however carefully taken, errors in reading and interpretation are likely to occur. It is also recognized that whether or not a given pressure, if not excessive, is decidedly abnormal to a given patient must be determined individually; and it may be asserted here that a pressure deemed too high should not always be tampered with.

Wonderful accuracy in determining heart action has been developed by Crehore of New York by his method of light-interference rings. Crehore's registrations of heart action by this method and Meara's clinical interpretation of these cardiograms represents scientific work of a high order.

A short and clear exposition of the various polygraphic methods of studying circulatory ability is presented by Satterthwaite[2] of New York. As the number of circulatory tracings is multiplied, careful readings of the composite picture give greater and greater accuracy to the diagnosis and the selection of the advisable treatment. Polygraphy means two or more tracings taken simultaneously, as that of the heart, carotid, jugular, radial, liver or any two of them. The electrocardiograph, when used by an expert, becomes the most accurate of the clinical circulatory instruments. This has been well shown by Lewellys F. Barker of Baltimore. If the apex, carotid, radial and jugular tracings are superimposed the relationship of each one to the others is graphically demonstrated. As above stated, Satterthwaite's article is brief but illuminating in its explanation of graphic analysis of the circulation.

INTERPRETATION OF ARTERIAL TRACINGS

The interpretation of the arterial tracing shows that the nearly vertical up-stroke is due to the sudden rise of blood-pressure caused by the contraction of the ventricles. The long and irregular down-stroke means a gradual fall of the blood-pressure. The first upward rise in this gradual decline is due to the secondary contraction and expansion of the artery; in other words, a tidal wave. The second upward rise in the decline is called the recoil, or the dicrotic wave, and is due to the sudden closure of the aortic valves and the recoil of the blood-wave. The interpretation of the jugular tracing, or phlebogram as the vein tracing may be termed, shows the apex of the rise to be due to the contraction of the auricle. The short downward curve from this apex means relaxation of the auricle. The second lesser rise, called the carotid wave, is believed to be due to the impact of the sudden expansion of the carotid artery. The drop of the wave-tracing after this carotid rise is due to the auricular diastole. The immediate following second rise not so high as that of the auricular contraction is known as the ventricular wave, and corresponds to the dicrotic wave in the radial. The next lesser decline shows ventricular diastole, or the heart rest. A tracing of the jugular vein shows the activity of the right side of the heart. The tracing of the carotid and radial shows the activity of the left side of the heart. After normal tracings have been carefully taken and studied by the clinician or a laboratory assistant, abnormalities in these readings are readily shown graphically. Especially characteristic are tracings of auricular fibrillation and those of heart-block.

---

[2] Satterthwaite, Thomas E.: Month. Cyclop. and Med. Bull., April, 1912.

SYMPTOMS AND SIGNS OF CARDIAC DISTURBANCE

These preliminary suggestions for a careful study, both theoretically and clinically, of circulatory disturbances are to urge a more careful interpretation of the symptoms and signs in patients who may not show very evident symptoms of cardiac disability or insufficiency. As recently so well expressed by Dr. Charles L. Greene[3] many a patient who has circulatory debility is branded with the "misnomer of neurasthenia." Many a patient is treated for gastric indigestion, or intestinal disturbances, or muscular rheumatism, or myalgias, or "malaria," or general debility, or even for the more scientific interpretation of intestinal fermentation and putrefaction (the latter of which, of course, must often be corrected) when the trouble really is cardiac tire and cardiac muscular weakness.

---

[3] Greene, Charles L.: THE JOURNAL A. M. A., Aug. 31, 1912, p. 685.

The pulse is not necessarily irregular, the heart may show no manifest enlargement and no murmurs, and yet many of the symptoms which occur under the conditions above named will be corrected when the heart is aided or given rest. A slight exercise test for the heart will often demonstrate weaknesses or inability to sustain effort, when the examination of a patient after some rest or without any exertion would seem to show that the circulation was perfectly normal. Many an ache, referred not only to the left chest but to the right, not only toward the left shoulder but to the right, not only toward the left side but around over the right side of the liver, or down into the abdomen, or up into the throat, with the diagnosis of the cause most uncertain, will be corrected when the heart is given a little more physical rest by resting the patient, or by the administration of small doses of digitalis. In fact, with the cause of such disturbances so uncertain, the therapeutic test of the administration of digitalis is satisfactory in any number of instances.

In this age of cardiac and circulatory tire, owing to the recklessness, speed and intensity of the age, some physicians frequently postpone the diagnosis of the cause of a disturbance until after a week of the action of digitalis, and many times the improvement of the condition is so rapid as to be almost phenomenal. This is not to suggest that digitalis be used carelessly or recklessly or without a careful decision as to whether or not the drug is contra-indicated, but is to urge a more careful analysis of disagreeable internal sensations before we term the patient a hysteric, a neurasthenic, a melancholic or a malingerer.

To be specific, if the patient complains of pains in any part of the chest or upper abdomen, or of leg-aches, or of being weary, or exhausted, or of sleeplessness at night, or of pains in the back of his head, we should investigate the cardiac ability, besides ruling out all of the more frequently recognized causes of these disturbances.

If there is more dyspnea than normally should occur in the individual patient after walking rapidly or climbing a hill or going upstairs; if after a period of a little excitement one finds that he cannot breathe quite so deeply, or that something feels tight in his chest, the heart needs resting; if after one has been driving a motor car or even sitting at rest in one which has been going at speed or has come unpleasantly near to hitting something or to being run into, it is noticed that the little period of cardiac disturbance and chest tension is greater than it should be, the heart needs resting.

If the least excitement or exertion increases the cardiac speed abnormally, it means that for many minutes, if not actually hours during the twenty-four, the heart is contracting too rapidly, and this alone means muscle tire and muscle nutrition lost, even if there is no actual defect in the cardiac muscle or in its own blood-supply. If we multiply these extra pulsations or contractions by the number of minutes a day that this extra amount of work is done, it will easily be demonstrable to the physician and the patient what an amount of good a rest, however partial, each twenty-four hours will do to this muscle. Of course anything that tends to increase the activity or the disturbance of the heart should be corrected. Overeating, overdrinking (even water), overuse or perhaps any use of alcohol, tobacco, tea or coffee should all be prevented. In fact, we come right to the discussion of the proper treatment and management of beginning high blood-pressure, of the incipiency of arteriosclerosis, of the prevention of chronic interstitial nephritis, and the prevention of cardiovascular-renal disease.

Besides the instruments of precision above referred to, more careful percussion, more careful auscultation, more careful measurements, *x*-ray pictures and fluoroscopic examinations of the heart, and a study of the circulation with the patient standing, sitting, lying and after exercise, makes the determination of circulatory ability a specialty and the physician who becomes an expert a specialist. It is a specialization needed to-day almost more than in any other line of medical science.

Greene, referring to the work of Henry Head, Sherrington, MacKenzie, and Alexander Lambert's interpretation of their investigations, shows that the heart is a sensitive organ and can cause sensations from disturbances of its various parts much more subtle than the angina pectoris which we recognize. The auricles may refer their disturbances to the axillæ and shoulders; the ventricles may refer their disturbances to the lower parts of the thorax, to the upper abdomen, to the inner surfaces of the upper arm or to the ulnar side of the forearm and wrist; if the aorta is in trouble, pain may be referred up into the neck, or even into the back of the head.

PROPHYLAXIS

As so much of our science, to-day, is aimed toward prevention, we should by every possible care prevent the heart from becoming weakened; we should, by treating every tonsillitis with the greatest of care and by removing or partially eradicating all tonsils that are diseased, take means to prevent the occurrence of rheumatism and chorea; we should never allow a patient to become active too soon after any serious illness, or even immediately after any slight illness, and certainly should keep a patient at rest for hours after every anesthesia, and in bed for at least two weeks after a severe operation. It is to be regretted that the aim of many surgeons to-day is to have their patients up and about, after an operation, in from a week to ten days. This is an unnecessary cardiac test, even if the operative region excused it. Even after so simple a thing as a migraine, a biliary colic, a renal colic or any other acute pain in which there has been a short starvation period and the administration of active narcotic drugs, with the reflex tire to the circulation from the pain, to say noting of actual pneumogastric reflexes on the heart, if the patient rises too soon he will invariably feel cardiac exhaustion, and his heart will not recuperate for hours and perhaps days. If this is repeated several times the patient will develop a cardiac muscle weakness.

CLASSIFICATION OF CARDIAC DISTURBANCES

For the sake of discussing the therapy of cardiac disturbances in a logical sequence, they may be classified as follows:

- Pericarditis
    - Acute
    - Adherent
- Myocarditis
    - Acute
    - Chronic
    - Fatty
- Endocarditis
    - Acute, simple
        - malignant
    - Chronic
- Valvular Lesions
    - Broken compensation
    - Cardiac drugs
    - Diet
    - Resort treatment
- Cardiac disease in children
- Cardiac disease in pregnancy
- Coronary sclerosis
- Angina pectoris
    - Pseudo-angina
- Stokes-Adams disease
- Arterial hypertension
- Cardiovascular-renal disease
- Arhythmia
- Auricular fibrillation
- Bradycardia
- Paroxysmal tachycardia
- Hyperthyroidism
- Toxic disturbances
- Physiologic hypertrophies
- Simple dilatation
- Shock
- Stomach dilatation
- Anesthesia in heart disease

ACUTE PERICARDITIS

As this inflammation is generally secondary to some other condition, its treatment cannot be positively outlined. Also, it is often a terminal condition, and in such instances the results of treatment are of necessity *nil.* The most frequent terminal cause is nephritis; other terminal causes are pulmonary tuberculosis, adjacent abscesses, cancer or other growth.

The most frequent infectious cause is rheumatism; other infectious causes are cerebrospinal fever, typhoid fever, acute miliary tuberculosis, pneumonia and sepsis. Accidental causes are traumatism and an adjacent inflammation of the pleura.

The result of an inflammation of the pericardium may be a fibrous exudate, or an exudate which is both serous and fibrous, or one in which pus is present in considerable amount.

The onset of pericarditis may be more or less acute, or it may commence insidiously. For this reason, during severe illness, and especially in those diseases which are known to have pericarditis often as a sequence, frequent examination of the heart should be made as a routine procedure.

SYMPTOMS AND SIGNS

If there is pain or much aching in the cardiac region, it tends to disappear with the exudate, if such is to occur, in the same way as the pain of pleurisy. If there is much exudate, the pressure on the heart of course increases, the cardiac dulness enlarges, dyspnea occurs and even perhaps later cyanosis. As the exudate accumulates the patient must lie higher and higher in order that the fluid may gravitate to the lowest part of the sac and give the heart the greatest ability to work. Reflex pains may occur from disturbances of the pneumogastric nerve, or from the weight and pressure of the enlarged and heavy pericardium. Reflex vomiting may be a troublesome and distressing symptom.

Acute pericarditis occurring in rheumatism, in acute infections and from simple injuries tends to recovery. In dry pericarditis with serious adhesions, or if adhesions occur as a sequence of acute pericarditis, the future prognosis is bad, as myocarditis may develop and sudden death or acute dilatation may occur. As above stated, if pericarditis develops during the progress of chronic disease, such as interstitial nephritis, or during sepsis, or from abscesses or growths in the region of the pericardium, the prognosis is bad.

TREATMENT OF ACUTE PERICARDITIS

In acute pericarditis absolute mental as well as physical rest is essential. Even if the patient does not appear to be seriously ill and has not much fever, he should not be allowed to have visitors, to discuss business matters or to carry on any conversation, however little exciting. Anything that increases his heart-beat increases the irritation of the inflamed surfaces of the pericardium. He should not be allowed to sit up, either to eat or to attend to the calls of Nature. These rules are imperative, and when they are followed the pain is less, the heart beats less rapidly, is less hampered by pressure from whatever exudate may be present, and the adhesions which are likely to form will be less in amount and less serious for the future work of the heart.

The treatment, of course, depends largely on the cause of the pericarditis, as, if the cause is one of those just enumerated in which the prognosis is dire, any treatment directed toward the pericardial inflammation is almost useless. The pericarditis under these conditions will be more or less benefited, if at all affected, by the treatment directed toward the cause.

The indications for treatment in all other instances are:

1. To attempt to abort the inflammation.
2. To stop the pain.
3. To limit, if possible, the amount of exudate, and to diminish the exudate already present.
4. To diminish the rapidity of the heart and to strengthen it.

1. Abortive Treatment. — For many years bloodletting was considered of the greatest importance in the early treatment of this disease; but owing to the fact that, except from traumatism, pericarditis rarely occurs except as a sequela of acute disease after the patient has been sick a long time, or as a terminal condition in a patient who has long been chronically diseased and therefore has already lost more or less strength, venesection has been nearly abandoned. Leeches may be used over the region of the pericardium, and cups are sometimes used. Dry cupping is more frequently used. These measures sometimes seem to reduce the inflammation, and certainly often relieve pain, but the most valuable local treatment is cold, which may be applied either in the form of an ice-bag or by a small coil through which ice-water is caused to flow by siphonage. Cold may be applied more or less continuously, depending on the sensations of the patient. The bag or ice-cap must not be overfilled and must not be heavy, as the patient often cannot stand pressure over the pericardium. Sometimes the relief from pain and the diminution of the number of the heart-beats is marked, and from this reason alone the cardiac inflammation may be inhibited. If cold applications are not tolerated by the patient (and they often are not in children) warm applications may be used, such as flaxseed poultices, or cloths wrung out of hot water and covered with oil-silk, and the pain will often be relieved thus. While hot applications would not tend to abort the inflammation, they probably do not tend to promote it.

A diminished diet, of small amount at a time, and such purging as the patient's strength will allow are essential in attempting to curtail the seriousness or amount of this inflammation.

Just what can be done locally or generally to combat the inflammation actively must depend on the cause. When the inflammation occurs as a complication of acute rheumatism, it has been suggested that salicylates, which do not inhibit rheumatism and may be depressant to the heart, should be stopped if they are being administered; but if the salicylates are apparently improving the inflammation in the joints, pericarditis would not contra-indicate their continued use. Except in large doses, salicylates probably do not depress the heart. In pericarditis it is perhaps well always to administer an alkali in some form unless otherwise contra-indicated, whether the cause is rheumatism or not. A diminished alkalinity of the blood would always increase the likelihood of an augmented amount of pericardial or endocardial inflammation. The blood must be kept strongly alkaline. It is possible that one of the reasons why pericarditis or endocarditis occurs so frequently in serious prolonged fevers is that the patient has not eaten enough cereals or other carbohydrates, and the system has become more or less endangered by acidosis. In other words, carbohydrate starvation is inexcusable with our present understanding of the danger from acidemia and even from a diminished amount of alkalies in the blood.

The causes of pericarditis being so varied, any antitoxin treatment or any vaccine treatment could be indicated only if the cause of the inflammation rendered the serum or vaccine advisable.

2. Stopping the Pain.—Nowhere else in the body should pain be so speedily combated as when it occurs in the region of the heart. Morphin, with or without atropin, as deemed best, should be administered hypodermatically in the amount and with the frequency necessary to stop the pain and quiet the restlessness. As above stated, the frequent need for morphin may be prevented by use of the ice-bag. Morphin might even be considered an abortive treatment, as nothing tends so much to inhibit this inflammation as the quietude of the heart caused by the absence of pain, the production of sleep and the prevention of restlessness, muscle twitching and muscle movements. In other words, the more quiet the patient is, the more quiet is the heart.

If for any reason morphin is contra-indicated, and if pain is not a symptom, the patient's nerves may be quieted and rest may be given by sodium bromid, or by veronal-sodium, the dose of the former being 2 gm. (30 grains) two or three times in twenty-four hours, according to its action and the necessity for it, and the dose of the latter 0.2 gm. (3 grains) once in six hours, if deemed necessary.

Especially if there are cerebral symptoms, as typically presented in cerebrospinal meningitis, and especially if the arterial tension is low, the subcutaneous administration of an aseptic ergot will quiet the central nervous system, increase the blood-pressure, quiet the heart and prolong the action of a single dose of morphin. It is the best plan to administer ergot deep into the muscles, with the deltoid as the place of choice. If the skin is properly cleansed, the syringe clean and the preparation of the drug aseptic, no inflammation or abscess will ever occur. If there is any painful swelling a wet alcohol dressing to the part will soon relieve it. The frequency with which ergot should be so administered depends on the results and the indications. Once in twelve hours for several doses is generally the best method for its use.

3. *The Exudate.*—When a fluid exudate into the pericardium has occurred from inflammation — that is,

when it is not an exudate from disturbed kidneys or circulation — it will continue to increase to some extent in spite of any treatment. Just how much this exudate may be prevented by the use of small blisters over or around the heart, and just how much watery stools and diuresis may prevent the advance of the exudate is difficult to determine. Small blisters, properly applied, have many times seemed to be the determining factor in stopping the increase in the fluid, or to have been the starting cause of the resorption of the exudate.

The amount of purging that should be caused by saline cathartics such as sodium sulphate (Glauber salt), potassium and sodium tartrate (Rochelle salt), or the official compound jalap powder cannot be declared dogmatically. Saline purging should be governed by the character of the circulation. If the heart is strong, the pulse not weak, and the blood-pressure good, nothing is more valuable in the above condition. Portal depletion is of great advantage, especially if the amount of liquid ingested is kept as low as possible, so that the blood-vessels may become thirsty and thus tend to absorb an exudate wherever they find it. Much harm has been done, however, and death has been caused by saline purgatives in endeavoring to relieve edemas from a failing heart or to prevent a uremia from kidney inflammation. The depression following such purging is often serious. If the circulation is weak, dependence should be placed on purgation by some of the simple vegetable cathartics or a small dose of calomel. While it is advisable to give a saline in concentrated solution, it should not be so strong as to cause vomiting. With our better understanding of magnesium absorption and the depressant effect of magnesium on the nervous system, magnesium salts should not be used in serious conditions.

Diuretics often do not act well when most needed. The simplest diuretic is potassium citrate, given in wintergreen or peppermint water, in doses of 2 gm. (30 grains), three or four times in twenty-four hours. One or more of the vegetable, non-irritant diuretics may be tried if preferred. If the sickness preceding the pericarditis was not a long fever, and the heart muscle is considered in good condition, digitalis in small doses may be the best possible diuretic. Incidentally it will slow the heart, if there is not much elevation of temperature, and will give some cardiac rest.

Although the patient's diet should be limited in bulk, and especially in amount of liquids, good nutrition should soon be given. Systemic weakness certainly tends to increase the exudate; systemic strength aids in absorption of the exudate.

Iron is early indicated, and nothing is better than 5 drops of the tincture of chlorid of iron in a little lemonade or orangeade, administered once in eight hours.

If the exudate tends to decrease, it perhaps may be hastened by the local application of tincture of iodin over the cardiac region. Also the administration of small doses of an iodid, as 0.3 gm. (5 grains) of sodium iodid, given in plenty of water three times a day, is useful. An iodid circulating in the blood seems to aid absorption. It has long been believed that iodin in the blood tends to promote absorption of thickened, left-over material from exudates, and to prevent the formation of strong fibrous adhesions. Until our knowledge is more exact in this matter, it is advisable to use iodid as suggested. If the above-named dose is not tolerated, less should be given.

If in spite of all the therapeutic measures suggested, the fluid increases and the pericardium becomes more distended and the heart's action more labored, paracentesis must be done. The point where the aspirating needle should be inserted into the pericardium depends somewhat on the conditions in each individual case. It is often best to insert an exploratory needle first. This will determine the fluidity and character of the exudate. If pus is found, a more radical surgical procedure than simple paracentesis must be done immediately. The point of puncture for aspiration most frequently chosen is in the fourth or fifth intercostal space, about an inch to the left of the sternal margin. Paracentesis is also often done in the region of the normal apex-beat. The position of the patient is determined by his dyspnea; he should lie in the position most comfortable for him. The fluid should be withdrawn slowly and the pulse carefully watched. The withdrawal of a small amount of fluid may later seem to be the starting cause of resorption of the rest of the fluid. On the other hand, it may often be not of more value than the simple removal of the immediate pressure, the fluid may again accumulate, and more radical surgery must be performed.

4. To Strengthen the Heart.—Most of the methods of meeting this indication have already been stated, namely, absolute rest; absolute quiet; the use of the bed-pan; any movement that must be made should be deliberate; the nurse and other attendants must be quiet; necessary conversation must be brief, and every method must be used to quiet and prevent the heart's action from becoming rapid. The food taken should be small in amount and non-stimulating; that is, no tea or coffee should be given, and nothing too hot or too cold. Movements of the bowels should be caused with the least possible general disturbance. If the patient does not sleep, he must be made to sleep. The whole body and the nervous system must have periods of rest. If the heart is very weak, small doses of morphin may be used. If the heart is not weak, bromids or chloral may be given. If the blood-pressure is high, such hypnotics will lower it, or if the heart is strong and the condition does not contra-indicate it, aconite may be used in small doses, for a day or two, unless the fever is high and it seems advisable to use one of the coal-tar antipyretics, which will of themselves reduce the blood-tension and the heart activity.

As above stated, pain must not be allowed. Sometimes, when there is thought not to be myocardial degeneration, or the heart has not been injured by prolonged fever, digitalis in small doses may slow the heart and act for good.

Convalescence.—The convalescence should be prolonged as in any other cardiac inflammation. The patient should be given more and more nourishing food, and the iron tonic may be changed to a capsule containing 0.05 gm. of quinin and 0.05 gm. of reduced iron, three times a day.

It is a question as to when patients convalescent from pericarditis should be permitted exercise. It has been thought that gentle movements and possibly exercise, sooner than theoretically justified, might cause the heart to beat a little more actively and possibly prevent the formation of tight adhesions between the two layers of the pericardium. Whether such activity of the heart will prevent adhesions is something that has not been determined.

The small doses of sodium iodid, perhaps 0.2 gm. (3 grains) two or three times a day, should be continued for some time. Iodid in this dosage does no harm and may do a great deal of good.

ADHERENT PERICARDITIS

Following dry pericarditis or pericarditis with an exudate, especially when the exudate is fibrinous in character, the fibrous substance which is not absorbed or resorbed may develop into connective tissue, and the two pericardial surfaces become permanently grown together, causing the so-called adherent pericarditis. These adhesions between the two surfaces of the pericardium may be general throughout the entire pericardial sac, or they may be limited to some one or more parts of the pericardium. Perhaps one of the most frequent points of adhesion is the anterior part of the pericardium, while the apex is the part most likely to be free even when other parts of the pericardium have grown together.

This freedom of the apex is probably due to the constant and more extensive motion of the apical portion of the heart, and is the reason that it has been suggested, as referred to under acute pericarditis, that, other conditions not contra-indicating, the patient may be allowed to move about a little during convalescence to cause the heart to beat more actively. Sometimes the surfaces of the pericardium are not closely adherent to each other, but bands of adhesion stretch from one surface to the other.

After adhesions have taken place between the two layers of the pericardium the action of the heart is impaired, serious interference with the cardiac action may develop, and sudden death may occur. If the heart is given all the rest possible during the acute phase of the disease, there will be less likelihood of the surfaces becoming so irritated that adhesions readily form. Anything that permits complete absorption and resorption of the exudate will tend to prevent these hampering adhesions. If the adhesions are such as to cause irregular heart, recurrent pain and the danger of sudden death, surgical help has been suggested. This surgical procedure is to remove a portion of the ribs, perhaps of the third, fourth and fifth, to allow the heart more freedom of action to compensate for the impairment of its activity from the adhesions. Such an operation was first suggested by Brauer of Heidelberg in 1902.

The question of the best method of producing anesthesia in this condition of the heart is a serious one. A patient might die during the anesthesia; but he might also die at any time from cardiac spasm. Probably chloroform, administered by a skilled anesthetist, would be the safest anesthetic for this operation. In certain instances, in adults, perhaps local anesthesia would be sufficient. Pain reflexes, however, would be as serious as would the chloroform anesthesia.

Such an operation would be indicated when the apex is fixed so that there is a constant sensation of hugging of the heart at the fourth and fifth ribs, with paroxysms of pain and cardiac weakness.

MYOCARDIAL DISTURBANCES

While the myocardium is the most important muscle structure of the body, it has but recently been studied carefully or well understood clinically or pathologically. A heart was "hypertrophied" or "dilated" or perhaps "fatty." It suffered from "pain," "angina pectoris," from some "serious weakness" or from "coronary disease," and that ended the pathology and the clinical diagnosis. This is the age of heart defects; no one can understand a patient's condition now no matter what ails him, without studying his heart. No one can treat a patient properly now without considering the management of his circulation. No one should administer a drug now without considering what it will do to the patient's heart.

Although we are scientifically interested in the administration of specific treatments, antitoxins and vaccines; although we have a better understanding of food values, and order diets with more careful consideration of the exact needs of the individual, and although we are using various physical methods to promote elimination of toxins, poisons and products of metabolism, we have until lately forgotten the physical fact that one-thirteenth of the weight of a normal adult is blood. A man who weighs 170 pounds has 13 pounds of blood. The above proportion of course is not true in the obese, and is not true in children. Whether the individual is sick in bed, miserable though up and about, or beginning to feel the first sensations of slight incapacity for his life-work, his ability properly to circulate this one-thirteenth of his weight through the various arterial and venous channels and capillary tracts must now, with the increasing tension and speed of our lives, be taken into consideration.

The more and more frequently repeated statements that the operation was successfuly performed but that the patient died of shock, and that the typhoid fever and the pneumonia were being successfully combated, but that the patient died of heart failure, and together with the increase in arteriosclerosis, cardiac disturbances and renal disease, emphatically present the necessity of more carefully studying the circulation. A better understanding and the constant study of the blood-pressure shows nothing but the necessity of the age. The unwillingness of the patient to suffer pain, even for a few minutes, without some narcotic, generally a cardiac debilitating drug, means that, if he is a sufferer from chronic or recurrent pain, he has taken a great deal of medicine that has done his heart no good. Repeated high tension of life raises the blood-pressure and puts more work on the heart. Therefore the heart is found weary, if not actually degenerated when any serious accident, medical or surgical, happens to the patient.

The requirements of the age have, then, necessitated that the heart be more carefully studied, and therefore the heart strength and its disturbances are better understood. The mere determination as to where the apex-beat is located, as to what murmurs may be present is not sufficient; we must attempt to determine the probable condition of the myocardium. The following conditions are recognized: (1) acute myocarditis, (2) chronic myocarditis (fibrosis, cardiosclerosis), (3) fatty degeneration, and (4) fatty heart.

ACUTE MYOCARDITIS

Probably most acute infections cause more or less myocarditis, depending on their intensity and their prolongation. This disturbance of the heart is often unrecognized, and has been simply referred to as "the heart growing weaker from the fever process." The acute infections most likely to cause a myocarditis are rheumatism, influenza, sepsis, cerebrospinal meningitis, diphtheria, typhoid fever and scarlet fever. It is probably rare when acute endocarditis occurs that more or less myocarditis is not present. The acute myocarditis may develop some fatty degeneration, and with this softening and weakening of the heart muscle acute dilatation readily occurs, which may be a cause of sudden death, or if less serious, may be the cause of prolonged disability, if the heart ever recovers its original size and strength.

The symptoms are often indefinite and the diagnosis of the condition hardly possible. It may be taken for granted, however, that hardly any serious illness can long continue without cardiac muscle disturbance. If endocarditis is present, soft systolic murmurs soon appear. With the acute myocarditis developing the apex-beat is less positive, less accentuated, and later it becomes diffuse and even feeble. The closure of the aortic valve is less typically sharp, showing that the blood-vessels are not so thoroughly filled. The peripheral circulation is not so active, the blood-pressure falls and the heart becomes more rapid, especially on the least exertion. All of these signs indicate myocardial weakness.

The treatment of this condition is largely preventive. It should be well recognized that prolonged high fever, prolonged insufficient or improper nutrition, prolonged acute pain and especially prolonged septic processes will always cause myocardial degeneration. It should be recognized that after ether and chloroform anesthesia, especially after chloroform, the heart muscle may be disturbed and the tonicity be lost. Therefore after anesthesias, after operations, after all illnesses that have lasted more than a few days the convalescence of the patient must be more or less deliberate. Sudden rising, sudden erect posture, the exertion of walking too early, going up stairs too early or taking moderate, and later severe exercise too early, may cause dilatation of the heart muscle that has become weakened by acute myocarditis. If myocarditis is believed or known to be present, cardiac tonics such as digitalis should not be given; large doses of strychnin should not be given; vasocontractors such as ergot should not be given; large amounts of food or large bulks of liquid should not be taken into the stomach at one time; in fact, unless there is some special indication, the twenty-four-hour amount of fluid should be diminished. The surface circulation and the muscle circulation should be improved by such cold or warm water applications as the disease or condition call for. Massage should be early inaugurated to promote the return circulation. The heart should be treated as though it were the frailest of Venetian glass and would crack with the least rough handling, or even with a rapid change of temperature, great cold or too much heat. A prolonged, tedious convalescence, with the return to activity so graded as to give the heart no strain and to keep its work always just below what it is able to do will often mean return to perfect strength and health.

No cardiac debilitating drug should be administered when myocarditis has been surmised or diagnosed. The safest hypnotic, if one is needed, is morphin in small doses. If there are weakening perspirations, atropin should be given, especially as it is also a circulatory stimulant. Calcium in almost any form seems to be of value in the majority of heart conditions. It is a sedative to the nervous system, and is certainly indicated in acute myocarditis. Calcium lactate is perhaps the best salt to administer, in doses of 0.25 gm. (4 grains), three or four times in twenty-four hours. Calcium glycerophosphate may be used, in powder form or in capsule, in doses of 0.30 gm. (5 grains) three or four times in twenty-four hours; or lime-water may be given.

An exact prognosis of this inflammation is impossible. We do not know how far an acute myocarditis may progress and entire recovery take place; we do not know how slight a myocarditis may cause serious symptoms. Clinically we know that many patients after serious illness never again have perfect circulatory strength. Other patients almost die of heart failure, and yet apparently absolutely recover their ability to do hard physical work.

CHRONIC MYOCARDITIS: FIBROSIS

Chronic myocarditis may develop on an acute myocarditis, but is generally a slowly progressive chronic process from the beginning; it occurs mostly in persons past middle life, and as a rule is not primarily associated with rheumatism or valvular disease of the heart. Perhaps generally the term "chronic myocarditis" is incorrect, as a real inflammatory condition is not present and has not been present; it is really a degenerative process with the development of connective tissue, a fibrosis and more or less hardening of the arterioles, a cardiosclerosis. In many instances this fibrosis is associated with fat deposits or fatty degeneration. The disease is often caused by a narrowing or obstruction or calcareous degeneration of the coronary arteries, thus diminishing the blood-supply to the heart muscle. This chronic myocardial degeneration is often a part of the general arteriosclerosis, and is an important factor in what is termed cardiovascular-renal disease. In simple chronic renal diseases the heart first normally hypertrophies to overcome the increased blood tension and increased resistance.

The principal causes of this degeneration are normal old age, or premature age caused by various conditions. In other words, anything that hastens arteriosclerosis will cause myocardial degeneration. The causes recognized as most frequently producing this condition are syphilis; gout; repeated attacks of rheumatism; excess in the use of alcohol (meaning repeated daily too large amounts, as well as actual dipsomania); the overuse of case; to reduce the amount of meat eaten, especially if there is intestinal indigestion; to relieve intestinal indigestion; to cause free daily movements of the bowels; to abstain from any food that tends to cause gastric or intestinal flatulence; to abstain from such foods as contain nucleins, if the patient is gouty; to take frequent warm baths (not too hot) to promote the secretions and the circulation in the skin; and to take such daily exercise as seems advisable. If the patient cannot take exercise, simple calisthenics or massage should be instituted.

Whether nitroglycerin or other nitrite is advisable depends on the peripheral blood-pressure. If the blood-pressure is low, or not higher than is best for the patient, such treatment would be inadvisable. If, from the supposed cause, iodid seems to be indicated, it should be given in small doses and continued for some time. It is often wise, however, to give small doses, as 0.10 or 0.20 gm. (2 or 3 grains) once or twice in twenty-four hours, for a long period, to any patient who has fibrosis or sclerosis in any form. Iodid tends to prevent the progress of connective tissue formation. It is quite possible that some of its value is in activating a sluggish or imperfectly acting thyroid gland. If the patient is old, his thyroid is subinvoluting, and a little more of its activity will be of advantage. Many diseases that cause chronic myocarditis also cause, later, subactivity of the thyroid. Thyroid extract may be indicated if the patient is obese.

If, in spite of this management and treatment, the patient has cardiac asthma attacks, with or without pain, especially if there are pendent edemas, the question arises as to whether or not digitalis should be given. In such cases one cannot tell without trying whether digitalis will be of benefit or will cause more discomfort. A small dose of an active preparation should be given at first twice in twenty-four hours, and after a week once in twenty-four hours, its action being carefully watched and the decision as to whether the dose is too large or too small arrived at.

It may do a great amount of good; it can cause increased cardiac pains. If used carefully and stopped when it appears not to be acting well, it will do no harm.

Chilling of the surface of the body should be avoided; in other words, sudden cold or sustained severe cold, by increasing the contraction of the peripheral blood-vessels and putting more strain on the heart muscle, is to be avoided if possible. More hours in bed at night and lying down after the heavier meals of the day will tend to give the heart the kind of rest it needs. Also complete rest for one day a week, or a rest of several days at a time, and a rest, both mental and physical, with such walking, golfing or riding as seems advisable, for at least one month every year, will prolong the lives of these patients, and may make an imperfect heart act well for months and years. If the patient is anemic he should, of course, receive some non-astringent iron, and an *Eisenzucker* tablet (saccharated oxid of iron), in small doses, 0.20 gm. (3 grains), once or twice in twenty-four hours, is sufficient.

PROGNOSIS

The prognosis of a case diagnosed as chronic myocarditis or chronic degeneration of the heart is doubtful, as one cannot tell until several weeks or months of observation whether this particular heart also has fatty degeneration or not. If there is fatty degeneration, the prognosis is bad. If there is no serious fatty degeneration, the patient, with the modified life outlined, may live for a long time. Acute dilatation from any serious strain on the heart may occur, and if there is fatty degeneration it is likely to occur at any time. Attacks of cardiac asthma are always serious and always damage the heart a little more.

FATTY DEGENERATION

Fatty degeneration of the heart muscle may be caused by acute poisoning (as phosphorus, arsenic, etc.), by serious infections, or it may follow fibrosis of the heart or coronary artery disease. The symptoms are those of serious circulatory weakness, which does not seem to improve under any ordinary management. It is difficult, if the heart is enlarged, to determine whether there is more or less serious acute dilatation or whether the heart muscle has suffered fatty degeneration. Prolonged rest in bed and then a slow convalescence followed by graded exercise will help the dilated heart; it does not do much for real fatty degeneration.

The treatment of such a patient requires the best of judgment as to the amount of food and liquid that should be given and as to the regulation of the administration of laxatives, the sponging of the body, and the means of producing sleep if there is insomnia, and how much reading, conversation or amusements should be allowed, how much stimulation by strychnin or other stimulating drug should be given, and to decide whether or not very small doses of digitalis should be tried. These are all matters for individualizing, and are matters for the best medical judgment that we are called on to give. How much repair can take place in a heart muscle when fatty degeneration has started we do not know. Such treatment will give the heart the only chance it has to recuperate, but the prognosis is bad.

FATTY HEART

The cause of deposits of fat around the heart or in between its chambers is the same as the cause of general obesity. These patients are likely to be obese, or at least to have large abdomens with large deposits of fat around the abdomen. This fat in itself will interfere somewhat with abdominal respiration. This tends to cause dyspnea and the heart tends to be disturbed from these causes, if much fat is not really in the pericardium. The symptoms are those of imperfect heart action; the patient is dyspneic on exertion or on leaning over, the heart acts rapidly on such exertion, the patient puffs, perspires easily, becomes leg-weary and becomes sedentary in his habits and more or less incapacitated for work. He may not be a large eater; if he is, and his eating habit is corrected, the prognosis is better than if he is putting on weight in spite of eating sparingly.

The general treatment is that for obesity, and if the heart muscle is intact various depletion methods may be inaugurated. More and more exercise, sweatings from Turkish baths, electric-light baths, body baking, vigorous massage and more or less purging are all valuable. Anything that reduces the general weight will help the heart. The prognosis is often good.

ENDOCARDITIS

It should be understood that especially in acute conditions a positive separation of endocarditis from myocarditis is incorrect. Acute endocarditis can probably not occur without some myocarditis, and myocarditis probably does not occur without some endocardial disturbance and perhaps some pericardial irritation. This is especially true in endocarditis that occurs during any acute infection, even in rheumatism. The greater the amount of pericarditis, the more serious is the acute condition. The greater the amount of myocarditis, the more doubtful is the heart strength in the near future. The greater the amount of endocarditis, the greater the doubt of freedom from future permanent valvular lesions.

Endocarditis may be divided into: acute mild (simple) endocarditis, acute malignant (ulcerative, infective) endocarditis, chronic endocarditis and valvular disease.

ACUTE MILD ENDOCARDITIS

This inflammation of the endocardium is generally confined to the region of the valves, and the valves most frequently so inflamed are the mitral and aortic. There may be a slight inflammation or actual ulceration and loss of tissue. Vegetations more or less constantly occur on the inflamed surfaces, with more or less danger of particles becoming loosened and movng free in the bloodstream, causing embolic obstruction in different parts of the body. There is also more or less probability of serious adhesions or contractions occurring from the healing of the ulcerated surfaces. In other words, the future health and welfare of the valves depends on the fact that the inflammation has healed without contractions or adhesions.

It is often difficult to decide when acute endocarditis has developed, but with the knowledge that the endocardium often becomes inflamed during almost any of the acute infections, the physician should repeatedly examine the heart for murmurs, for muffled closure of the valves, or for other evidences of endocarditis or myocarditis during the acute infective process.

It has been shown positively that acute endocarditis is due to microorganisms, generally streptococci, staphylococci or pneumococci, and, more frequently than once believed, gonococci. The most frequent causes are acute rheumatic fever, diphtheria, pneumonia, cerebrospinal meningitis, scarlet fever, erysipelas, influenza, chorea, gonorrhea, sepsis and typhoid fever. It may also follow a follicular tonsillitis which is rheumatic in type but has not caused arthritis. Tuberculosis may also occasionally cause an endocarditis. Organisms may be found in a terminal simple endocarditis due to a chronic disease, as tuberculosis or cancer; such inflammations may have been caused by circulating toxins.

It will be noticed by the above classification that the terms "mild" and "malignant" endocarditis are used. It is to convey the fact that there may be no etiologic distinction between the two forms, and it is impossible to decide clinically in the beginning of an endocardial inflammation which form is present. In the malignant form the infection is probably more serious or the infective germs are more active, the ulcerations deeper, and the likelihood of emboli and the seriousness of such embolic infarcts more serious and more dangerous. In other words, the difference in inflammation in the two cases is really one of degree, and the classification is made to coincide with this probable fact. It is of course clinically recognized that endocarditis following certain diseases, especially rheumatism, is of the simple or mild type, while that termed ulcerative endocarditis may occur apparently as a primary or general infection and the causative bacteria are generally readily discovered in the blood.

A SECONDARY AFFECTION

Mild endocarditis is rarely a primary affection, and is almost invariably secondary to one of the diseases named above. Nearly 75 per cent. of secondary endocarditis occurs as a complication of acute articular rheumatism and chorea, or subsequently. On the other hand, about 40 per cent. of all patients with acute articular rheumatism develop endocarditis, sometimes perhaps so mild as to be hardly discoverable. This complication is most likely to occur during the second or third week of rheumatic fever. It is not sufficiently recognized that a subacute arthritis, recurring tonsillitis, and even a condition of the system with acute, changeable and varying joint and muscle pains may all develop a mild endocarditis, even with subsequent valvular lesions. Therefore in all of these conditions the decision can be made only as to how much rest the patient must have or how serious the condition is to be considered by careful examination of the heart in every instance.

Children are more liable than adults to this complication, especially with rheumatism. Therefore acute mild endocarditis with future valvular lesions occurs most frequently during childhood and adolescence, and if one attack has occurred, a subsequent infection, especially of rheumatism, is likely to cause another acute endocarditis.

PATHOLOGY

The part of the heart most affected is the part that has the most work to do — the left side of the heart — and of this side the left ventricle and therefore the mitral and aortic valves; the most frequent valve to be inflamed and to suffer permanent disability is the mitral valve, the valve which in its inflamed condition is subjected to the greatest amount of pressure and therefore irritation. Not infrequently soft systolic murmurs are heard at the pulmonary and tricuspid valves during acute endocarditis. It is rare, however, that these valves are so affected during childhood or adult life as to be permanently disabled.

Whether a diminished alkalinity of the blood in rheumatism has anything to do with the cause of the frequent complication of endocarditis has not been determined. Whether the administration of alkalies to the point of increasing the alkalinity of the blood is any protection against the complication of endocarditis has also not been positively demonstrated, although clinically such treatment is believed by a large number of practitioners to be wise.

A chronic endocarditis with permanent lesions of the valves may become an acute inflammation with an infectious provocation.

It has been shown that even in a few hours after endocarditis has started little vegetations composed of fibrin, white blood-cells, red blood-pigment and platelets may develop. Practically in all instances such vegetations develop, and later become more or less organized into connective tissue. These little vegetations, generally minute, perhaps not exceeding 4 mm. in height, are irregular in contour like a wart. Some of these may have small pedicles, and as such, of course, are more likely to become loosened and fly off into the bloodstream. It is of interest to note that these little vegetations are more likely to be on the left side of the heart than the right; more likely to be on the valves than any other part; and more likely to be on the mitral valve than on the aortic. The consequence is a more frequent permanent disability of the valves of the left side of the heart, and of these more frequently the mitral. Although these little vegetations and excrescences sooner or later become mostly connective tissue, still fibrin and white blood-cells may form thin layers over them, more or less permanent. In this fibrin are frequently found bacteria, even when there has been no recent acute inflammation. The deeper layers of the endocardium during acute inflammation may become infiltrated with young cells, with resultant softening and destruction of the intercellular substance. This softening and some swelling of the lower layers of the endocardium allow the pushing up of these extravasated blood-cells which, being covered with fibrin, makes the little vegetations above described; and as just stated, the fibrin may form a more or less permanent cap. If this cap is disintegrated or lost and the cells under it washed away in the blood-stream, ulceration takes place, which may be more or less serious, even to the perforation of a valve or actual erosion of one of its cusps, and the parts of the valves most seriously affected are the parts that strike against each other on closure; in other words, as previously stated, the parts subjected to the greatest strain and the greatest amount of friction during the inflammation are the parts most seriously affected afterward.

If a perforation has occurred, it may make a permanent leak. If an erosion of the edge of the valve has occurred, it may make permanent insufficient closure. If the valve has become thickened and stiffened during the cicatricial healing it may not only be incompetent, but may not open perfectly, and a narrowed orifice be the consequence. During the healing of these granulating ulcers there may be thickening of the part or shrinking of the tissue, and the valve may become shortened by adhesion to the wall, or the cusps of the valve may adhere together so that the valve becomes permanently unable to open properly or to close properly, or to do either.

Not infrequently, and probably more frequently than we recognize, recovery without any of the pathologic lesions just described follows mild endocarditis. The occurrence of simple endocarditis is undoubtedly frequent during acute disease and is unrecognized because there are no lesions of the heart at the time or subsequently; but valvular lesions only too frequently follow the endocarditis that occurs with rheumatism. Occasionally the ulcerations become serious and ulcerative endocarditis or malignant endocarditis develops on the mild inflammation. In this form the little vegetations are likely to become loosened, fly off into the bloodstream, and cause emboli in different parts of the body.

Recently Fraenkel[4] concluded that the microscopic nodules which occur in endocarditis in the myocardium, and which consist of the several varieties of white blood-corpuscles first referred to by Ashoff in 1904, are characteristic only of acute rheumatism. Fraenkel found these nodules in the myocardium in a case of chorea, showing the close relationship between it and rheumatism.

SYMPTOMS AND SIGNS

While repeated careful examination of the heart during acute infections will generally show signs of endocarditis if it is present, even if there are no subjective symptoms, the disease may be so insidious as not to be noted until a valvular lesion occurs. Often, however, during the course of the disease, especially in rheumatism, there is a slight increase in fever and there is a discomfort complained of in the region of the heart, frequently accompanied by slight dyspnea. Real pain is seldom present unless the pericardium is affected. If the myocardium is much inflamed at the same time, the heart becomes more rapid and the blood-tension lowered, the apex-beat diminished in intensity and perhaps not palpable. If there is pain, with or without pericarditis, it is often referred to the epigastrium, especially in children. The patient is often nervous, restless and sleepless. In simple endocarditis emboli rarely occur. If they do, of course the signs will be in the part where the infarct occurs. Besides the diminished intensity of the apex-beat and its greater diffusion, the valve sounds may be muffled, and sooner or later there may be systolic murmurs over the different orifices. Of course systolic murmurs may be due to a disturbed condition of the blood, but if they occur with the above-mentioned symptoms and signs, endocarditis should be diagnosed. If the heart becomes seriously weak and the patient suffers much dyspnea, myocarditis should be known to be present with the endocarditis. If there is a diastolic murmur, there can be no question of serious endocarditis having occurred. Unexplainable palpitation during acute illness has been thought to be a distinct symptom of endocarditis.

---

[4] Fraenkel: Beitr. z. path. Anat. u. z. allg. Path., 1912, lii, 597.

TREATMENT OF ENDOCARDITIS

As mild endocarditis rarely occurs primarily but is almost always secondary to some acute disease, its immediate treatment is only a slight modification of that of the disease that is causing it. A complication that is so frequent should always be expected, and consequently warded off or prevented, if possible. Knowledge of the diseases that most likely cause endocarditis makes frequent heart examinations a necessity, to note when it arrives. While an extra heart tire, sleeplessness, the circulation of unnecessary toxins from a bad condition of the bowels and from improperly selected food all make this complication more likely, its occurrence is, nevertheless, often unpreventable.

The most efficacious preventive measures are sleep, rest, absence of pain, prevention of exertion, proper food that does not cause flatulence or other indigestion, good, sufficient daily movements of the bowels, the prevention of intestinal distention, and maintenance of a clean, moist surface of the body, produced by such sponging and bathing as the temperature demands.

The disease having developed, the indications for treatment are really few; in fact, the treatment is mostly negative. There is generally but little pain; the temperature from simple endocarditis alone is not high, and the acute symptoms tend to abate.

Local Treatment.—Endocarditis having been diagnosed, especially if there is palpitation or pain, the ice-bag over the heart is often of considerable value, but not so efficient as in pericarditis. It often tends to quiet the heart, and may be of some value reflexly in slowing the inflammation. If it causes restlessness, however, and does not diminish, or perhaps in some instances increases, the pain, it certainly should be stopped. Children, in whom this complication so frequently occurs, generally do not bear the ice-bag well. Sometimes it may be advisable to substitute warm applications, and often a great deal of comfort is derived from them, the patient soon going to sleep. One of the greatest values of either cold or hot applications is diminution of the discomfort from the cardiac disturbance, and the stopping of any pain that may be present. If they do not do this, there is no object in using either cold or heat.

The discomfort from blisters over the heart during the acute stage of endocarditis is greater than any good that they can do. In adults a few small blisters may be used intermittently around the borders of the heart, after the acute symptoms are over, to act reflexly on the heart and possibly aid absorption of inflammatory products. Sometimes improvement seems to follow such treatment; it certainly can do no harm.

During convalescence the skin over the heart may be painted with iodin, repeated often enough to cause stimulation without injuring the skin; it seems at times to be of value. Various iodin ointments or iodid ointments have been used, but they probably have no more value than the administration of small doses of iodid.

Systemic Treatment.—As this complication most frequently occurs during acute rheumatism, the question arises as to the value or harmfulness of salicylates and alkaline drugs. With our recent better understanding of the action of pure salicylates on the heart (either natural or synthetic salicylic acid, which have been shown to act identically, if equally pure, we must believe that in any ordinary dosage they will injure the heart but rarely. While salicylic acid will not prevent endocarditis, it should be continued, if it is benefiting the arthritis. In other words, the indication for its use depends on its effect on the joints. As it acts at times almost as a specific in rheumatism, it would seem that it should be of value in the endocarditis caused by rheumatism. On the other hand, the endocarditis occurs during the second or third week of acute rheumatism, after the blood has been thoroughly saturated with salicylic acid. Therefore it certainly does not tend to prevent rheumatic endocarditis; hence for this complication alone salicylic acid is not indicated.

ALKALIES

Anything that tends to increase the acidity of the tissues and to diminish the alkalinity of the blood, whether from starvation or other causes, seems to produce endocardial and myocardial irritation, if not actual inflammation. Therefore in a disease like rheumatism that seems to be made worse by anything that increases the acidity, alkalies are obviously indicated, and it is probable that an increased alkalinity of the blood tends to prevent endocardial irritation, and may soothe an inflammation already present. Until we have some positive knowledge to the contrary, alkalies should be freely administered during endocarditis, especially during rheumatic endocarditis. Potassium citrate in 2-gm. (30-grain) doses, in wintergreen water, should be given every three to six hours, depending on how readily the urine is made alkaline. This may be given with the salicylic acid treatment, and also when the salicylic acid has been stopped. It may be well, if sodium salicylate is being used, to give also sodium bicarbonate, the sodium bicarbonate often preventing an irritation of the stomach from the sodium salicylate, the dose being equal parts of the sodium salicylate and the sodium bicarbonate administered in plenty of water. If some other form of salicylic acid is preferred, novaspirin, which is methylene-citrylsalicylic acid and contains 62 per cent. of salicylic acid, is perhaps the least irritant to the stomach of the salicylic preparations. This drug is decomposed in the intestine into its component parts, salicylic acid and methylene-citric acid. If this drug is combined with sodium bicarbonate, the disintegration into its component parts would be likely to occur in the stomach.

IRON

It is essential for the welfare of the patient, especially after a long illness before the complication of endocarditis could occur, and in rheumatic fever, in which all meat and meat extractives have been kept from the diet, that small doses of iron should be administered daily. Not only the fever process, but the salicylic acid tend to prevent the healthy normal growth of red corpuscles, and such patients suffering from rheumatism are often seriously anemic after the acute inflammation has ceased. The iron administered may be 5 drops of the tincture of the chlorid, in lemonade or orangeade, twice in twenty-four hours (and it should be remembered that lemon and orange burn to alkalies in the system and do not act as acids); or 0.1 gm. (1½ grains) of reduced iron in capsule twice in twenty-four hours, or a 3-grain tablet of *Eisenzucker* (saccharated oxid of iron) twice in twenty-four hours.

OPIUM

As so many times repeated, real pain must be stopped, and morphin, either by the mouth or hypodermatically, should be used to the point of stopping such pain. If the patient is a young child, the sulphate of codein or the deodorized tincture of opium may be used, in the dose found sufficient, and either one will act satisfactorily. The dose given should be small and repeated sufficiently often to stop the pain. The dose necessary for the given individual will soon be learned, and that dose may be repeated at such intervals as the condition may require. Sometimes the hypnotic selected, if one is needed, will be sufficient to quiet the cardiac aches or pains.

BROMIDS AND CHLORAL

If there is much restlessness and the circulation is good, that is, if myocarditis is probably not present, the bromids may be of great value, especially in children. The dose should be sufficient to quiet the nervous system. The drug may be discontinued after a few days, if the conditions improve. If the bromid, except in large doses, will not cause sleep, a sufficient dose of chloral should be given. Chloral is one of the most satisfactorily acting drugs to produce sleep and to cause cardiac rest that we have. While it should not be given if there is real cardiac weakness, the good that it does is so much greater than the possible bad effect on the heart, that it should not be forgotten for some newer hypnotic. The worst part of this drug is its taste, and the best way to administer it is to have it in solution in water and the dose given on cracked ice with a little lemon juice, to be followed by a good drink of water and a piece of orange pulp for the patient to chew. Ordinarily a bad-tasting drug such as chloral is well administered in effervescing water, but effervescing waters are generally inadvisable when there is any kind of inflammation of the heart, as they are likely to cause distention of the stomach and pressure on the heart. Some physicians prefer chloralamid as a less disagreeable drug and one that acts almost as efficiently as chloral. As the dose of this must be larger than the dose of chloral, it is a question of doubt as to which is the better drug to use. Of the newer hypnotics, veronal-sodium (sodium-diethyl-barbiturate) is among the best. It acts quickly, is less depressant and is a safer salt than most of the other newer hypnotics. It is the readily soluble sodium salt of veronal (diethylbarbituric acid). When combined with any active drug, sodium seems to make it less toxic and less depressant. The dose of this drug is 0.2 to 0.3 gm. (3 to 5 grains).

PREVENTION

If the patient is weak, the circulation depressed, the blood-pressure low, and the heart rapid, the drug advisable to produce rest and sleep is almost always morphin or some other form of opium. Morphin, with few exceptions, is a cardiac tonic and a cardiac stimulant, unless the dose is much too large. As long as the bowels are daily moved and the food is not given at the time of the full action of the morphin, when digestion might be delayed or interfered with, in most patients the action of this drug during serious illness is entirely for good. The greatest mistake in using morphin for the production of sleep, or for physical and mental rest and comfort when there is not severe pain, is in giving too large a dose. If pain is not severe, or is due to inflammatory distention of some undilatable part, to pressure on some nerve, to distention of some tube by a calculus or to some serious injury to the nerves, large doses of morphin are not needed. Small doses will act much more efficiently. It is excessively rare that a hypodermic of one-fourth grain of morphin sulphate is needed, except for the conditions enumerated. It is often a fact that so small a dose as one-eighth grain of morphin or even one-sixth grain will cause sufficient stimulation of a nervous patient, because its primary stimulant effect on the spinal cord is greater than its depressant effect on the brain, to require another dose (one-fourth grain altogether) to give such a patient rest. On the other hand, this patient may many times be quieted by one-tenth grain of morphin sulphate on account of the size of the dose being not sufficient to stimulate the spinal cord. Many a time clinically when one-eighth grain has failed, a dose of one-fourth grain having been apparently necessary, a change to one-tenth grain has proved entirely and perfectly satisfactory, much to the surprise of the attending physicians.

DIET

As intimated above in the paragraph on "prevention," the diet during endocarditis must be carefully regulated. It must be sufficient, it must be appropriate for the disease in which the complication occurs, but it must be in such dosage and must be administered with such frequency as to cause the least possible indigestion. Large amounts of milk are rarely advisable. Too much milk is certainly given, even in rheumatism. While pretty well tolerated by children, it is often badly tolerated, as far as digestive symptoms are concerned, by adults. The amount of liquid given should be governed by the amount of urine passed and by the amount of perspiration. The patient should not be overloaded with liquid if he does not need it.

LAXATIVES

If the bowels are known to be in excellent condition and not loaded with fecal matters, brisk catharsis is not needed simply because endocarditis has developed. If the bowels have been neglected, a small dose of calomel, aided by a compound aloin tablet, is necessary and good treatment. Subsequent movements of the bowels should be daily obtained by vegetable laxatives with occasional enemas, as needed. With an inflammation of the heart and the possibility of myocarditis developing or being actually present, it is not advisable to use salines freely or often.

CARDIAC DRUGS

Whether any drug should be used which acts directly on the heart is often a question for decision. As endocarditis is generally secondary to some acute disease, the patient has become weakened already and the circulation is not sturdy; therefore such a drug as aconite is probably never indicated. The necessary diminished diet, catharsis, hypnotic, salicylic acid and alkalies all tend to quiet the circulation and diminish any strenuosity of the heart that may be present. Unfortunately, during fever processes digitalis in ordinary doses rarely slows the heart; and while it might slow the heart if given in large doses, it would also cause too powerful contractions of the ventricles. Digitalis is inadvisable if there is much endocardial inflammation, and especially if there is supposed or presumed to be myocardial inflammation. If a patient had already valvular disease from a previous endocarditis, and during this attack insufficiency of the heart was evidenced by pendent edemas, digitalis should be administered; but it probably should not be given to other patients during the acute period of inflammation.

BATHS

During rheumatism the peripheral blood-vessels are generally dilated and the skin perspires profusely. This is caused not only by the rheumatism, but also by the salicylates. The surface of the body should be sponged with cold, lukewarm or hot water, depending on the temperature, especially of the skin. The cold water will reduce the temperature and tone the peripheral blood-vessels; the hot water, if the temperature is low and the skin moist and flabby, will cleanse it and also tone the peripheral blood-vessels. If the blood-vessels are dilated and the perspiration profuse, atropin is indicated, both as a cardiac stimulant and contractor of the blood-vessels and as a preventer of too profuse sweating. The dose should be from 1/200 to 1/100 grain for an adult, given two or three times in twenty-four hours, depending on its action and the indications. It should be remembered that atropin is not a sleep-producer; it may stimulate the cerebrum. Therefore at night it might well be combined with a possible necessary hypodermatic injection of morphin.

STRYCHNIN

The question of the advisability of strychnin is a constant subject for discussion. Strychnin is overused in most patients who are seriously ill. In a patient in whom we are trying to cause nervous and muscular rest strychnin is certainly contra-indicated. On the other hand, if the heart is acting sluggishly, the peripheral circulation is imperfect, and atropin is not acting well, it is advisable to give strychnin in a dose not too large and not too frequently repeated. Strychnin should be avoided, if possible, in the evening so that the patient may sleep. Whether it should be given by the mouth or hypodermatically would depend entirely on the seriousness of the condition. Once in six hours is generally often enough for strychnin to be administered unless the dose is very small.

ALCOHOL

It is rarely, if ever, advisable to use alcohol. In certain instances, however, especially in older patients who are accustomed to alcohol, a little whisky administered several times a day may act only for good, both as a food and as a peripheral dilator. But it must be remembered that alcohol is not a cardiac stimulant, and that a large dose will be followed by more cardiac depression. Nitroglycerin may act as well as whisky in the kind of cases mentioned. Caffein stimulation in any form is generally inadvisable during inflammation of the heart.

PROGNOSIS AND CONVALESCENCE

The duration of acute endocarditis varies greatly; it may be two or three weeks, or the inflammation may become subacute and last for several months. Although mild endocarditis rarely causes death of itself, it may develop into an ulcerative endocarditis, and of course then be serious *per se*. On the other hand, it may add its last quota of disability to a patient already seriously ill and death may occur from the combination of disturbances.

As soon as all acute symptoms have ceased, rheumatic or otherwise, and the temperature is normal, the amount of food should be increased; the strongly acting drugs should be stopped; especially the alkalies should not be given too long, and the salicylates should only be given intermittently, if at all; iron should be continued, massage should be started, and iodid should be administered, best in the form of the sodium iodid, 0.1 to 0.2 gm. (1½ to 3 grains), twice in twenty-four hours, with the belief that it does some good toward promoting the resorption of the endocardial inflammatory products and can never do any harm. Prolonged bed rest must be continued, visitors must still be proscribed, long conversations must not be allowed, and the return to active mental and physical life must be most deliberate.

No clinician could state the extent to which the valvular inflammation will improve or how much disability of the valves must be permanent. It is even stated by some clinicians that a rest in bed for three months is advisable. While this is of course excessive, certainly when the future health and ability of the patient are under consideration, and especially when the patient is a child or an adolescent, time is no object compared with the future welfare of the individual's heart. It is one of the greatest pleasures of the clinician to note such a previously inflamed heart gradually diminish in size and the murmurs at the valves affected gradually disappear. Although they may have disappeared while the patient is in bed, he is not safe from the occurrence of a valvular lesion for several months after he is up and about.

While the discussion of hygiene would naturally be confined to the hygiene of the disease of which the endocarditis is a complication, still the hygiene of its most frequent cause, rheumatism, should be referred to. Fresh air and plenty of it, and dry air if possible, is what is needed in rheumatism, and a shut-up, overheated and depend on where the emboli locate. If in the abdomen, there are colicky pains with disturbances, depending on the organ affected; if in the brain, there may be paralysis, more or less complete. If an infarct occurs in one of the organs of the body there must of necessity occur a necrosis of the part and an added focus of infection. If a peripheral artery is plugged, gangrene of the part will generally occur, if the patient lives long enough.

TREATMENT

If pneumonia or gonorrhea is supposed to be the cause of the endocarditis, injections of stock vaccines should perhaps be used. If the form of sepsis was not determinable, streptococcic or staphylococcic vaccines might be administered. It is still a question whether such "gunshot" medication with bacteria is advisable. Patients recover at times from almost anything, and the interpretation of the success of such injection treatment is difficult. Exactly how much harm such injections of unnecessary vaccines can produce in a patient is a question that has not been definitely decided. Theoretically an autogenous vaccine is the only vaccine that should be supposed to be successful with our present knowledge. The vaccine treatment of ulcerative endocarditis was not shown to be very successful by Dr. Frank Billings in his investigation a short time ago.[5]

Other treatment of malignant endocarditis includes treatment of the condition that caused it plus treatment of "mild" endocarditis, as previously described, with the meeting of all other indications as they occur. As in all septic processes, the nutrition must be pushed to the full extent to which it can be tolerated by the patient, namely, small amounts of a nutritious, varied diet given at three-hour intervals.

Whether milk or any other substance containing lime makes fibrin deposits on the ulcerative surfaces more likely or more profuse, and therefore emboli more

---

5. Billings, Frank: Arch. Int. Med., November, 1909, p. 409.

likely to occur, is perhaps an undeterminable question. In instances in which hemorrhages so frequently occur, as they do in this form of endocarditis, calcium is theoretically of benefit. Quinin has not been shown to be of value, nor has salicylic acid, unless the cause is rheumatism. Alcohol has been used in large doses, as it has been so frequently used in all septic processes. If the patient is unable to take nourishment in any amount, small doses of alcohol may be of benefit. It is probably of no other value. It is doubtful whether ammonium carbonate tends to prevent fibrin deposits or clots in the heart, as so long supposed. In fact, whenever the nutrition is low and the patient is likely to have cerebral irritation from acidemia, whenever the kidneys are affected, or whenever a disease may tend to cause irritation of the brain and convulsions, it is doubtful if ammonium carbonate or aromatic spirit of ammonia is ever indicated. Ammonium compounds have been shown to be a cause of cerebral irritation.

Perhaps the most advisable internal antiseptic treatment is that of hexamethylenamin. This drug should be given in doses of 0.3 gm. (5 grains), dissolved in plenty of water, once in six hours. The administration of mercury in any amount or of the sulphocarbolates or any phenol preparation is probably of not so great value as hexamethylenamin. Intestinal antisepsis may be attained more or less successfully by the administration of yeast or of lactic acid ferments together with suitable diet. The nuclein of yeast may be of some value in promoting a leukocytosis. It has not been shown, however, that the polymorphonuclear leukocyte increase caused by nuclein has made phagocytosis more active.

Malignant endocarditis may prove fatal in a few days, or may continue in a slow subacute process for weeks or even months.

CHRONIC ENDOCARDITIS

It is not easy to decide just when an acute endocarditis has entirely subsided and a chronic, slow-going inflammation is present. It would perhaps be better to consider a slow-going inflammatory process subsequent to acute endocarditis as a subacute endocarditis; and an infective process may persist in the endocardium, especially in the region of the valves, for many weeks or perhaps months, with some fever, occasional chills, gradually increasing valvular lesions and more or less general debility and systemic symptoms. Such a subacute endocarditis may develop insidiously on a previously presumably healed endocardial lesion and cause symptoms which would not be associated with the heart, if an examination was not made. Sometimes such a slow-going inflammatory process will be associated with irregular and intangible chest pains, with some cough or with many symptoms referred to the stomach, so that the stomach may be considered the organ that is at fault. There may be dizziness, headache, feelings of faintness, sleeplessness, progressive debility and a persistent cough, with some bronchial irritation and with occasional expectoration of streaks of blood, which may cause the diagnosis of incipient tuberculosis to be made. In other words, the need of a careful general examination must be emphasized again before a decision is made as to what ails the patient, or before cough mixtures are given unnecessarily, quinin is prescribed for supposed malarial chills, or various diets and digestants are recommended for a supposed gastric disturbance.

The term "chronic endocarditis" should be reserved for a slowly developing sclerosis of the valves. This may occur in a previous rheumatic heart and in a heart that has suffered endocarditis and has valvular lesions, or it may occur from valvular strain or heart strain from various causes; it is typically a part of the arteriosclerotic process of age and is mostly manifested at the aortic valve.

ETIOLOGY

Rheumatism is the cause of most instances of cardiac disease which date back to childhood or youth, while arteriosclerosis is the cause of most cardiac diseases in the adult. In the former case it is the mitral valve which is the most frequently affected, while in the latter it is the aortic valve. Any cause which tends to induce arteriosclerosis may be a cause of chronic endocarditis, such as gout, syphilis, chronic nephritis, alcoholism, excessive use of tobacco, excessive muscular labor and hard athletic work. Lead is also another, now rather infrequent, cause. Severe infections may tend to make not only an arteriosclerosis occur early in life, but also a chronic endocarditis. Heart strain may also be a cause of chronic endocarditis, especially at the aortic valve. Forced marches of soldiers, competitive athletic stunts, and occupations that call for repeated hard physical strain may all cause aortic valve disease. Tobacco, besides increasing the blood-tension and thus perhaps injuring the aortic valve, may weaken the heart muscle and cause disturbance and irritation and perhaps inflammation of the mitral valve.

There is no age which is exempt from valvular disease, but the age determines the valve most likely to be affected. If endocarditis occurs in the fetus, it is the right side of the heart that is affected; in children and during adolescence it is most frequently the mitral valve that is involved; while in the adult or in old age it is the aortic valve that is most likely to become diseased. Statistics have shown that the valves of the left side of the heart are diseased nearly twenty times as frequently as those of the right side of the heart. They also show that the mitral valve is diseased more than one and one-half times as frequently as the aortic valve. Early in life probably both sexes are equally affected with valvular disease, with perhaps a slight preponderance among females, because of their greater tendency to chorea. Females also show a greater frequency to mitral stenosis than do males. Aortic disease, on the other hand, from the very fact of their strenuous life and occupations, is nearly three times more frequent in men than in women.

PATHOLOGY

If a chronic endocarditis has followed an acute condition, some slight permanent papillomas or warty growths may be left from the healed granulating or ulcerated surfaces. Sometimes these little elevations on the valves become inflamed and then adhere together, or adhere to the wall of the heart, and thus incapacitate a valve. Sometimes these excrescences undergo partial fatty degeneration, or may take on calcareous changes and thus stiffen a valve.

If the chronic inflammation is not superimposed on an acute endocarditis there may be no cell infiltration and therefore no softening, but there is a tendency to develop a fibrillated structure, and a fibroid thickening of the endocardium occurs, especially around the valves. This induration causes contraction and narrowing of the orifices with shortening and thickening of the chordae tendineae, and the valves imperfectly open, or no longer close. Fatty degeneration may occur in the papillary growths with necrotic changes, and this may lead to the formation of atheromatous ulcers which may later become covered with lime deposits, and then a hard calcareous ring may form. Fibrin readily deposits on this calcareous substance and may form a permanent capping, or may slowly disintegrate and allow fragments to fly off into the blood-stream and cause more or less serious embolic obstruction. If this chronic endocarditis develops with a general arteriosclerosis, the same inflammation soon occurs in the aorta, and, following the endarteritis in the aorta, atheromatous deposits may also occur there. Chronic endocarditis of the walls of the heart, not in immediate continuity with endocarditis of the valves, is perhaps not likely to occur, except with myocarditis.

TREATMENT

A subacute or a chronic infective endocarditis should be treated on the same plan as an acute endocarditis, which means rest in bed and whatever medication seems advisable, depending on the supposed cause of the condition.

A chronic endocarditis which is part of a general arteriosclerosis requires no special treatment except that directed toward preventing the advance of the general disease.

CHRONIC DISEASE OF THE VALVES

PATHOLOGIC PHYSIOLOGY

The development of permanent injury to one or more valves of the heart may have been watched by the physician who cares for a patient with acute endocarditis. Or it may have been noted early during the progress of arteriosclerosis or other conditions of hypertension. On the other hand, many instances of valvular lesions may be happened on in a life-insurance examination, or are discovered by the physician making a general physical examination for an indefinable general disturbance or for local symptoms, without the patient ever having known that he had a damaged heart. The previous history of such a patient will generally disclose the pathologic cause or the physical excuse.

As soon as a valve has become injured, the heart muscle hypertrophies to force the blood through a narrowed orifice or to evacuate the blood coming into a compartment of the heart from two directions instead of one, as occurs in regurgitation or insufficiency of a valve. The heart muscle becomes hypertrophied, like any other muscle which is compelled to do extra work. Which part or parts of the heart will become most enlarged depends on the particular valvular lesion. In some instances this enlargement is enormous, increasing a heart which normally weighs from 10 to 12 ounces to a weight of 20 or even 25 ounces, and extreme weights of from 40 to 50 ounces and even more are recorded.

As long as the heart remains in this hypertrophied condition, which may be called normal hypertrophy since it is needed for the work that has to be done in overcoming the defect in the valve, there are no symptoms, the pulmonary and systemic circulation is sufficient, and the patient does not know that he is incapacitated. Sooner or later, however, the nutrition of the heart, especially in atheromatous conditions, becomes impaired, and the lack of a proper blood-supply to the heart muscle causes myocardial disturbance, either a chronic myocarditis or fatty degeneration. If there is no atheromatous condition of the coronary arteries, and arterial disease is not a cause of the valvular lesion, compensation may be broken by some sudden extra strain put on the heart, either muscular or by some acute sickness or a necessary anesthetic and operation. From any of these causes the muscle again becomes impaired and the heart, especially the part which is the weakest and has the most work to do relatively to its strength, becomes dilated, compensation is broken and all of the various circulatory disturbances resulting from an insufficient heart strength develop.

PRECAUTIONS TO BE OBSERVED

As long as compensation is complete, there is no medication and no physical treatment necessary for the damaged heart. The patient, however, should be told of his disability, and restrictions in his habits and life should be urged on him. The most important are that all strenuous physical exercise should be interdicted; competitive athletic work should be absolutely prohibited; prolonged muscular effort must never be attempted, whether running, rowing, wrestling, bicycle riding, carrying a heavy weight upstairs or overlifting in any form. The patient should be taught that he should never rush upstairs, and that he should never run rapidly for a car or a train or for any other reason; he should not pump up a tire, or repeatedly attempt to crank a refractory engine; even the prolonged tension of steering a car for a long distance is inadvisable. He should be told that after a large meal he is less capacitated for exertion than a man who has not a damaged heart. It is better if he drinks no tea or coffee; it is much better if he absolutely refrains from tobacco and alcohol. Prolonged mental worry, business frets and mental depression are all injurious to his heart. Anything that seriously excites him, whether anger or a stimulating drug, is harmful. Any disease that he may acquire, especially lung disturbances, as pneumonia or even a serious cough, requires that he take better care of himself and be more carefully treated and take more rest in bed than a patient who has not a damaged heart. Anything that raises the blood-pressure is of course more serious for his heart than for a perfect heart; therefore the drinking of large amounts of liquid, even water, is inadvisable. It simply means so much more work for the heart to do. Such patients should rarely be given any drug that causes cardiac debility, and should never take one without advice. This applies to all the coal-tar drugs.

One other fact should be impressed on the individual with a valvular lesion and compensation, and that is that he has but little, if any, reserve circulatory power. In other words, while he is in apparently perfect health, it takes little circulatory strain to push his heart to the point of danger or insufficiency. As nothing keeps this reserve so good or increases it more than rest, he should expect to have a restful day at least once a week, and a good rest of at least two or three weeks once or twice a year.

A patient with these restrictions may live for years with a serious valvular defect and may die of some intercurrent disease that has nothing to do with the circulatory system.

It is easily recognizable that as the majority of acute lesions of the valves occur in children, it is impossible to prevent them from taking more or less strenuous exercise, and this is probably the reason that we have so many serious broken compensations during youth or early adolescence.

As referred to under the subject of myocarditis, many symptoms for which a patient consults his physician are indefinite and intangible, though due to cardiac weakness. If a patient with a damaged heart has a sudden dilatation, of course his symptoms are so serious that the physician is immediately summoned. If, however, he has a slowly developing insufficiency of the heart muscle, his first symptoms are more or less indefinite cardiac pains, slight shortness of breath, slight attacks of palpitation, a dry, tickling, short cough occurring after the least exertion, some digestive disturbances, often sluggishness of the bowels, gastric flatulence, possibly nosebleeds, and sooner or later some edema of the lower extremities at the end of the day.

PHYSICS OF BROKEN COMPENSATION

To understand the physiology, pathology and the best treatment for broken compensation, it is necessary to study the physics of the circulation under the different conditions. With the mitral valve insufficient, a greater or less amount of blood is regurgitated into the left auricle, which soon becomes dilated. Distention of any hollow muscular organ, if the distention is not to the point of paralysis, means a greater inherent or reflex attempt of that organ to evacuate itself; the muscular tissue begins to grow and a hypertrophy of the left auricle with the above-named lesion develops. The muscular tissue of the auricle, however, is not sufficient to allow any great hypertrophy. The blood flowing from the pulmonary veins into the left auricle finds this cavity already partly filled with blood regurgitated from the left ventricle. The pulmonary blood, being impeded, tends to flow more slowly, and therefore dams back into the lungs, causing a passive congestion of the lungs. The pulmonary artery thus finds the pressure ahead unusually great, and the right ventricle reflexly learns that it requires a greater force to empty itself than

The dry, troublesome, tickling cough is generally due to a congestion of the blood-vessels at the base of the tongue, in the lingual tonsil region or possibly in the larynx. Later the passive congestion of the lungs may be sufficient to cause a bronchitis, with expectoration.

Sometimes these patients have, as indicative of primary cardiac distress, sharp pains through the heart. Such pains are the exception rather than the rule, and are more likely to occur in chronic myocarditis or in coronary disease; in other words, in true angina pectoris.

If there is considerable venous congestion there may be more or less frequent recurrent venous hemorrhages. This frequently is an epistaxis, or a bleeding from hemorrhoids, or in women profuse menstruation or a metrorrhagia.

It is perfectly understandable from the physics of the condition of broken compensation that anything that improves the tone of the heart and makes it again compensatory removes all of these many disabilities, congestions and subacute inflammations. If, however, these passive congestions are long continued, some organs soon become chronically degenerated. This is especially true of the liver and kidneys.

PHYSICS OF MITRAL STENOSIS

Mitral stenosis, though less common than mitral regurgitation, is a frequent form of disease of the valves, especially in women. Often this condition is associated with regurgitation, but in a simple mitral stenosis the greatest hypertrophy is of necessity in the right ventricle. The left auricle finds it difficult to empty all of its blood into the left ventricle during the ordinary diastole of the heart. This auricle then somewhat hypertrophies, but is unable to prevent more or less damming back of the blood into the lungs through the pulmonary veins. This causes passive congestion of the lungs, and the right ventricle finds that it must labor to overcome the increased resistance in the pulmonary artery, and hypertrophies to overcome this increased amount of work.

When this condition has become perfected, compensation is established and the circulation is apparently normal. Nature causes these hearts, when they are disturbed or excited, to pulsate slowly, causing the diastole to be longer than in a heart with mitral regurgitation. This allows more blood to enter the left ventricle, and the left ventricle, acting perfectly on the blood that it receives, causes a good systolic pressure in the aorta and the systemic arteries. The left ventricle in this condition does not become hypertrophied. If the heart does act rapidly and the left ventricle contracts on an insufficient amount of blood, the peripheral pulse is necessarily small and the arterial tension is diminished. Very constant in this condition, and of course noticeable whenever there is pulmonary congestion, is the sharp, accentuated closure of the pulmonary valve. The lungs on the least exertion are always a little overfilled with blood. The pulmonary circulation is always working at a little disadvantage.

The first symptoms of lack of compensation with the lesion of mitral stenosis are lung symptoms — dyspnea, cough, bronchitis, slight cyanosis, sometimes blood-streaks in the expectorated mucus and froth, and if the congestion is considerable some edema of the posterior part of the lungs, if the patient is in bed. Sooner or later during this failing compensation the right ventricle becomes dilated, and the symptoms of cardiac insufficiency and venous congestion occur, as above described with mitral insufficiency.

Again, as in mitral insufficiency, if compensation is restored in mitral stenosis, these symptoms are improved. These patients, however, are never quite free from dyspnea on exertion. Any inflammation of the lungs, even a severe bronchitis, is more or less serious for the patients and their hearts. The mucous membrane of their bronchial tubes and air vesicles is always hyperemic, and it takes little more congestion to all but close up some of the passages, and dyspnea or asthma or suffocating, difficult cough is the consequence.

PHYSICS OF AORTIC LESIONS

Next in frequency to mitral insufficiency is *aortic insufficiency,* which occurs most frequently in men. The cavity of the heart that is most affected by this lesion is the left ventricle, which receives blood both from the left auricle, and regurgitantly from the aorta. This part of the heart being the strongest muscular portion is the part most adapted to hypertrophy, and the hypertrophy with this lesion is often enormous. For a long time this large muscular section of the heart can overcome all difficulties of the aortic insufficiency. The pulse, however, will always show the quickly lost arterial pressure of every beat on account of the aorta losing its pressure through the regurgitant flow of blood. Sooner or later, from the impaired aortic tension causing a diminished or imperfect flow of blood through the coronary arteries, impaired nutrition of the heart muscle occurs. In other words, an interstitial or chronic myocarditis or fibrosis develops, with perhaps later a fatty degeneration. When this condition occurs of course the repair of the heart is impossible.

This form of valvular lesion is the one that is most likely to cause sudden death. In aortic regurgitation Nature causes the heart to beat rapidly. Such a heart must never beat slowly, as the longer the diastole prevails the more blood will regurgitate into the left ventricle, and death may occur from sudden anemia of the base of the brain. Such a heart may of course receive a sudden strain, or the left ventricle may dilate, and yet serious myocarditis or fatty degeneration may not have occurred.

The signs of lack of compensation are generally cardiac distress, rapid heart, insufficiency of the systolic force of the left ventricle, and therefore impaired peripheral circulation, a sluggish return circulation, pendent edemas, and soon, with the left auricle finding the left ventricle insufficiency emptied, the damming back of the blood as in broken compensation with the mitral lesions.

AORTIC STENOSIS

Aortic narrowing or *stenosis* is a frequent occurrence in the aged and in arteriosclerosis when the aorta is involved. It is not a frequent single lesion in the young. If it occurs in children or young adults, it is likely to be combined with aortic regurgitation, meaning that the valve has been seriously injured by an endocarditis.

The first effect of this narrowing is to cause hypertrophy of the left ventricle, and as long as this ventricle is able to force the blood through the narrowed opening at the aortic valve, the general circulation is perfect. Nature again steps in to cause such a heart to beat deliberately, allowing time for the contracting ventricle to force the blood through the narrowed orifice. The blood-pressure may be sufficient, or even increased if arteriosclerosis is present, although the rise of the sphygmograph tracing is not so high as normal. If the pressure in the aorta is sufficient from the amount of blood forced into it, the coronary arteries receive enough blood to keep up the nutrition of the heart muscle. Sooner or later, however, the left ventricle will become weakened, especially when there is arteriosclerosis or other hypertension, and chronic endocarditis and fatty degeneration result. If the left ventricle becomes sufficiently weakened or dilated the same damming back of the blood through the lungs and right heart occurs, and more or less serious signs of broken compensation develop. The main danger, however, with a heart with this lesion, occurring coincidently with arteriosclerosis, is a progressive chronic myocarditis.

OTHER LESIONS

Tricuspid insufficiency, except as rarely found in the fetus, is generally due to a relative insufficiency rather than to an actual disease of the tricuspid valve. In other words, if the right ventricle dilates the valve may be insufficient. *Tricuspid stenosis, pulmonary stenosis* and *pulmonary insufficiency* are rare, and are probably nearly always congenital.

The diagnosis as to whether the murmurs heard in the heart are hemic, functional, accidental, or indicative of valvular lesions would be without the scope of this book. It is always presumed that a correct diagnosis has been made, or at least a presumptively correct diagnosis. Frequently more than one murmur and more than one lesion in a heart is present. Often one murmur denotes a permanent lesion and another may be one that will become corrected when compensation is improved.

SYMPTOMATOLOGY AND TREATMENT OF CHRONIC VALVULAR LESIONS

Before discussing the treatment of broken compensation in general, it may be well to describe briefly the differences in the symptoms and treatment of the various valvular lesions.

MITRAL STENOSIS: MITRAL NARROWING

This particular valvular defect occurs more frequently in women than in men, and between the ages of 10 and 30, and is generally the result of rheumatic endocarditis or chorea, perhaps 60 per cent. of mitral stenosis having this origin. Other causes are various infections or chronic disease, such as nephritis. Of course, like any valvular lesion, it may be associated with other lesions, and sooner or later in many instances, when the left ventricle becomes dilated or weakened, mitral insufficiency also occurs.

It has sometimes seemed that high blood-pressure has caused the left ventricle to act with such force as to irritate this mitral valve and later develop from such irritation a sclerosis or narrowing, and stenosis occurs. It has been suggested that, though lime may be of advantage in heart weakness, as will be stated later, if the blood is overfull of calcium ions the valvular irritations may more readily have deposits of calcium, in other words, become calcareous, and therefore cause more obstruction. It is quite likely, however, that this sort of deposit is only a piece of the general calcification of tissue in arteriosclerosis and old age and could not be caused by the administration of calcium to a younger patient, and might then occur in older patients even if substances containing much calcium were kept out of the diet. Calcium metabolism in arteriosclerosis, and in softening of the bones is not well understood.

Patients with this lesion are seriously handicapped when any congestion of the lungs occurs, such as pneumonia, pleurisy or even bronchitis. Asthma is especially serious in these cases, and these patients rarely live to old age.

The pulse is generally slow, unless broken compensation occurs; dyspnea on exertion is a prominent symptom; the increased secretion of mucus in the bronchial tubes and throat is often troublesome, and there is likely to be considerable cough. If these patients have an acute heart attack a feeling of suffocation is their worst symptom and the dyspnea may be great, although there may be no tachycardia, these hearts often acting slowly even when there is serious discomfort. When compensation fails there is an occurrence of all the usual symptoms, as previously described.

The distinctive diagnostic physical sign of this lesion is the diastolic and perhaps presystolic murmur heard over the left ventricle, accentuated at the apex and transmitted some distance to the left of the heart. There is also an accentuated pulmonary closure. To palpation this lesion often gives a characteristic presystolic thrill at and around the apex.

The first symptoms of weakening of the compensation are irregularity in the beat and venous congestion of the head and face, causing bluing of the lips, often nosebleed and sometimes hemoptysis and insomnia. Later the usual series of disturbances from dilatation of the right ventricle occurs. As previously stated, with the absence of good coronary circulation and the consequent impaired nutrition, the left ventricle may also dilate and the mitral valve may become insufficient. Sudden death from heart failure may occur with this lesion more frequently than with mitral insufficiency and less frequently than with aortic insufficiency.

A particularly dangerous period for women with this lesion is when the blood-pressure rises after the menopause and the patients become full-blooded and begin to put on weight. Also, these patients always suffer more or less from cold extremities. In most cases they sleep best and with least disturbance with the head higher than one pillow.

Besides the usual treatment for broken compensation in patients with this lesion, digitalis is of the greatest value, and the slowing of the heart by it, allowing the left ventricle to be more completely filled with the blood coming through the narrowed mitral opening during the diastole, is the object desired. This drug acts similarly on both the right and left ventricles, and though there is no real occasion for stimulation of the left ventricle, and it is the right ventricle that is in trouble, dilated and failing, still a greater force of left ventricle contraction helps the peripheral circulation. The action on the right ventricle contributes greatly to the relief of the patient by sending the blood through the lungs and into the left auricle more forcibly, and the left ventricle receives an increased amount of blood, the congestion in the lungs is relieved and the dyspnea improves. Perhaps there is no class of cardiac diseases in which more frequent striking relief can be obtained than in these cases of mitral stenosis.

If the congestion of the lungs is very great and death seems imminent from cardiac paralysis, if cyanosis is serious and bloody, frothy mucus is being expectorated, venesection and an intramuscular injection of aseptic ergot may be indicated. Digitalis should also be given, hypodermatically perhaps, but its action would be too late if it was not aided by other more quickly acting drugs. The physician may often save life by such radical measures.

MITRAL INSUFFICIENCY: MITRAL REGURGITATION

This is the most frequent form of valvular disease of the heart, and is due to a shortening or thickening of the valves, or to some adhesion which does not permit the valves to close properly, and the blood consequently regurgitates from the left ventricle into the left auricle during the contraction of the ventricle. Such regurgitation may occur without valvular disease if for any reason the left ventricle becomes dilated sufficiently to cause the valve to be insufficient. Such a dilatation can generally be cured by rest and treatment. As with mitral stenosis, the most frequent causes are rheumatism and chorea, with the occasional other causes as previously enumerated.

The characteristic murmur of this lesion is a systolic blow, accentuated at the apex, transmitted to the left of the thorax, generally heard in the back, near the lower end of the scapula, and transmitted upward over the precordia.

Of all cardiac lesions, this is the safest one to have. Sudden death is unusual, the compensation of the heart seems to be most readily maintained, and the patient is not so greatly dangered by overexertion or by inflammations in the lungs. As in mitral stenosis, any increase in blood-pressure—whether the normal increase after the age of 40, any continued earlier high tension or increase from occupation or exercise—is serious as causing the left ventricle to act more strenuously, so that more blood is forced back into the left auricle, the lungs become congested, and the right ventricle, sooner or later, becomes incompetent.

When compensation fails with these patients, the first sign is pendent edema of the feet, ankles and legs; subsequently, if there is progressive failure of compensation, the usual symptoms occur.

The treatment is principally rest and digitalis, and the recovery of compensation is often almost phenomenal. Patients with this lesion are likely to be children and young adults, and the heart muscle readily responds *as a rule to* the treatment inaugurated. Later, in these patients, or if the lesion occurs in older patients, the return to compensation does not occur so readily. If the condition is developed from a myocarditis or from fatty degeneration of the heart, it may be impossible to cause the left ventricle to improve so much as to overcome this relative dilatation or relative insufficiency of the valve. If the dilatation of the left ventricle is due to some poisoning such as nicotin, with proper treatment—stopping the use of tobacco, administration of digitalis, and rest—the heart muscle will generally recover and the valve again properly close.

AORTIC STENOSIS: AORTIC OBSTRUCTION

Valvular disease at the aortic orifice is much less common than at the mitral orifice, and while stenosis or obstruction is less common from rheumatism or acute inflammatory endocarditis than is insufficiency of this valve, a narrowing or at least the clinical sign of narrowing, denoted by a systolic blow at the base of the heart over the aortic opening, is in arteriosclerosis and old age of frequent occurrence. If such narrowing occurs without aortic insufficiency at the age at which it usually occurs, it may not seriously affect the heart. It may follow acute endocarditis, but it most frequently follows chronic endocarditis or atheroma, in which the aortic valves become thickened and more or less rigid; this condition most frequently occurs in men.

Anything that tends to increase arterial tension, as tobacco, lead or hard work, or anything that tends to cause arterial disease, as alcohol or syphilis, is often the cause of this lesion.

At times the edges of the valves may grow together from ulcerative inflammation, and the lumen thus be diminished in size; or projecting vegetations may interfere with the opening of the valve and with the flow of blood. With such narrowing the left ventricle more or less rapidly hypertrophies to overcome its increased work.

The murmur caused by this lesion is a systolic one, either accentuated in the second intercostal space at the right of the sternum, or perhaps heard loudest just to the left of the sternum in this region. The murmur is also transmitted up the arteries into the neck, and may at times be heard in the subclavian arteries. It may also be transmitted downward over the heart. The pulse is slow, the apex of the rise of the sphygmographic arterial tracing is more or less sustained and rounded, and the rise is much less than normal.

If this lesion occurs in old age, there is general arterial disease present and the tension and compressibility of the arteries vary, depending on how much they are hardened. The disturbed circulation is evidenced by imperfect peripheral circulation and capillary sluggishness, with at times pendant edema of the feet and ankles, but, perhaps, little congestion of the lungs. The left ventricle being sufficient, there is no damming back through the left auricle to the lungs. The left ventricle may, however, become weakened, either by some sudden strain or by a chronic myocarditis, and relative insufficiency of the mitral valve may occur. The subsequent symptoms are typically those of loss of compensation.

This lesion may allow a patient to live for years, provided no other serious disturbance of the heart occurs, such as myocarditis or coronary disease; but sooner or later, with the failing force of the blood-flow and the lessened aortic pressure, slight attacks of anemia of the brain occur, causing syncope or fainting. Also, sooner or later these patients have little cardiac pains. They begin to "sense" their hearts. There may not be actual anginas, but a little feeling of discomfort, with perhaps pains shooting up into the neck, or a feeling of pressure under the sternum. Little excitements or overexertions are likely to make the heart attempt to contract more rapidly than it is able to drive the blood through the narrowed orifice, and this alone causes cardiac discomfort and the feeling of cardiac oppression.

It is essential, then, that these patients should not hasten and should not become excited; and any drug or stimulant that would cause cardiac excitement is bad for them. On the other hand, these are the very patients in whom, sometimes, alcohol in small doses may be advisable, especially if the patient is old; and a dose of alcohol used medicinally when an attack of cardiac disturbance is present is good treatment. The quick dilatation is valuable. Nitroglycerin will also do good work in these cases, and with high blood-tension may be the only safe drug for the patient to have on hand. As soon as his attack occurs, with or without real angina pectoris, let him dissolve in his mouth a nitroglycerin tablet. If he feels faint, he will feel better the moment he lies down, and in this instance he may be improved by a cup of coffee, or a dose of caffein, camphor or ammonia.

If the left ventricle becomes still weaker and shows signs of serious weakness, or if there is actual dilatation, the question of whether or not digitalis should be used is a subject for careful decision. The left ventricle should not be forced to act too sturdily against this aortic resistance. Consequently the dose of digitalis must be small. On the other hand, it frequently happens, especially in old age, that myocarditis or fatty degeneration has already occurred before this cardiac weakness develops in the presence of aortic narrowing, and digitalis may not be indicated at all. We cannot tell how far degeneration may have gone, however, and small doses of digitalis used tentatively and carefully, perhaps two or three drops of an active tincture two or three times a day, and then the drug carefully increased to a little larger dose to see whether improvement takes place, is the only way to ascertain whether digitalis can be used with advantage or not. If it increases the cardiac pain and distress, it should not be used. Strychnin is then the drug that should be relied on, with such other general medication as is needed, combined with the coincident administration of nitroglycerin, which may also be given in conjunction with digitalis, if deemed advisable. Generally, however, if a heart with aortic stenosis needs stimulation the blood-pressure is generally none too high, although there may be arteriosclerosis present. Therefore when nitroglycerin is indicated to lower blood-pressure, digitalis is not usually indicated; when digitalis is indicated to aid the heart, nitroglycerin is generally not indicated. These patients must have high blood-pressure to sustain perfect circulation at the base of the brain.

Patients who have this lesion should not use tobacco in large amounts, or sometimes even small amounts, as tobacco raises the blood-pressure and thus puts more work on the left ventricle; in the second place, if the left ventricle is failing, much tobacco may hasten its debility. On the other hand, with a failing left ventricle and a long previous use of tobacco it is no time to prohibit its use absolutely. A failing heart and the sudden stoppage of tobacco may prove a serious combination.

AORTIC INSUFFICIENCY: AORTIC REGURGITATION

This lesion, though not so common as the mitral lesion, is of not infrequent occurrence in children and young adults as a sequence of acute rheumatic endocarditis. If it occurs later in life it generally is associated with aortic narrowing, and is a part of the general endarteritis and perhaps atheroma of the aorta. Sometimes it is caused by strenuous exertion apparently rupturing the valve.

This form of valvular disease frequently ends in sudden death. On the other hand, it is astonishing how active a person may be with this really terrible cardiac defect. This lesion, from the frequent overdistention of the left ventricle, is one that often causes pain. While the left ventricle enlarges enormously to overcome the extra distention due to the blood entering the ventricle from both directions, the muscle sooner or later becomes degenerated from poor coronary circulation. Unless the left ventricle can do its work well enough to maintain an adequate pressure of blood in the aorta, the coronary circulation is insufficient, and chronic myocarditis is the result. If the left ventricle has maintained this pressure for a long time, edemas are not common unless the cardiac weakness is serious and generally permanently serious; that is, slight weakness, in this lesion, does not give edemas as does slight loss of compensation in mitral disease, and unless the weakness of the ventricle is serious, the lungs are not much affected.

The physical sign of this lesion is the diastolic murmur, which is loudest at the base of the heart, is accentuated over the aortic orifice, and is transmitted up into the neck and the subclavians, and down over the heart and down the sternum with marked pulsations of the arteries (Corrigan pulse) and often of some of the peripheral veins, notably of the arms and throat.

If the left ventricle becomes dilated the mitral valve may become insufficient, when the usual lung symptoms occur, with hypertrophy of the right ventricle; and if it fails, the usual venous symptoms of loss of compensation follow. This lesion not infrequently causes epistaxis, hemoptysis and hematemesis.

Digitalis is always of value in these cases, but it should not be pushed. If a heart is slowed too much the regurgitation into the left ventricle is increased. Therefore such hearts should not be slowed to less than eighty beats per minute, or sudden anemia of the brain and sudden death may occur. These patients must not do hard work.

TRICUSPID INSUFFICIENCY

This rarely, if ever, occurs alone; it is generally a sequence of other valvular defects, and represents an overworked, dilated right ventricle. It may, however, occur from lesions of the lungs which impede the blood-flow through them. Such are fibroid changes in the lungs, emphysema, prolonged chronic bronchitis, the last stages of pulmonary tuberculosis, old neglected pleurisies with cirrhosis or fibrosis of the lung, and repeated attacks of asthma — anything, whether valvular defect or pulmonary circulatory disturbance, that increases the pressure ahead and the work of this ventricle.

The symptoms are those of loss of compensation as described under other valvular lesions. There may be jugular pulsation, especially evident in the external jugular on the left side. The liver enlarges and may pulsate. There are edemas, dropsies, ascites and perhaps hemorrhages. The heart is enlarged and there is a soft systolic blow heard at the lower end of the sternum. The dyspnea is sometimes very great, and cyanosis may be present, especially during paroxysms of coughing.

This lesion of the heart is always benefited by digitalis, but the continuance of the improvement and its amount depend, of course, on the cause of the dilatation of the ventricle. Strychnin is often of advantage. These patients should rarely receive vasodilators, and hot baths, overheating, overloading the stomach and vigorous purging should never be allowed. Sometimes improvement will not take place until ascitic or pleuritic fluid, if present, has been removed.

TRICUSPID STENOSIS: TRICUSPID OBSTRUCTION

This is rare and probably always congenital, and is supposed to be due to an inflammation of the endocardium during intra-uterine life. In early childhood it is possible that it may be associated with left-side endocarditis.

A special treatment of the heart, if any is needed, would probably not be indicated unless there was associated tricuspid insufficiency, when digitalis might be used.

PULMONARY INSUFFICIENCY: PULMONARY REGURGITATION

If this rare condition occurs, it is probably congenital. A distinctive murmur of this insufficiency would be diastolic and accentuated in the second intercostal space on the left of the sternum. It should be remembered that aortic murmurs are often more plainly heard at the left of the sternum. Sooner or later, if this lesion is actually present, the right ventricle dilates and cyanosis and dyspnea occur. Digitalis would therefore be indicated.

PULMONARY STENOSIS: PULMONARY OBSTRUCTION

If stenosis is actually present in this location, the lesion is probably congenital. It might occur after a serious acute infectious endocarditis, but then it would be associated with other lesions of the heart. It has been found to be associated with such congenital lesions of the heart as an open foramen ovale or foramen Botalli, or with an imperfect ventricular septum, and perhaps with tricuspid stenosis — in short, a cardiac congenital defect. The right ventricle becomes hypertrophied, if the child lives to overcome the obstruction.

The physical sign is a systolic blow at the second intercostal space on the left; but as just stated, such a murmur must surely be dissociated from an aortic murmur if found to develop after babyhood, and it should also be diagnosed from the frequently occurring hemic, basic and systolic murmurs; that is, if signs of pulmonary lesions are not heard soon after birth or in early babyhood, the diagnosis of pulmonary defects can be made only by exclusion.

Unless the right ventricle is found later to be in trouble, there is no treatment for this condition. If the right ventricle dilates, digitalis may be of benefit.

ACUTE CARDIAC SYMPTOMS: ACUTE HEART ATTACK

It is not proposed here to describe the condition of sudden cardiac failure, or acute dilatation during disease, or after a severe heart strain, but to describe the terrible cardiac agony that occurs, sometimes repeatedly, with many patients who have valvular lesions. These patients may not have the symptoms of loss of compensation. Probably some one or more chambers of the heart become overdistended and act irregularly, or the blood is suddenly dammed up in the lungs, with the oppression and dyspnea caused by such passive congestion, or perhaps it is the right ventricle that is suddenly in serious trouble.

A physician receives an emergency call, and knows, if it is not a patient who has hysteria, that it is his duty to see the patient immediately. The friends of the patient all anxiously await the physician's arrival; front doors are often wide open, and the servants and the whole household are in a great state of excitement and anxiety. The position in which the patient will be found is that which he has learned gives him the greatest comfort. If the physician knows his patient, he will know how he will find him. He may be sitting up in bed; he may be standing, leaning over a chair; he may be sitting in a chair leaning over a table or leaning over the back of another chair; but he is using every auxiliary muscle he possesses to respire. He is generally bathed in cold perspiration; the extremities are often icy cold; he calls for air, and to stop fanning all in one breath; he wishes the perspiration wiped off his brow, and nearly goes frantic while it is being done;

there is agony depicted on his face; his eyes stare; his expirations are often groaning. Sometimes there is even incontinence of urine and feces, often hiccup or short coughs, perhaps vomiting, and possibly sharp pangs of pain in the cardiac region. A patient with these symptoms may die at any moment, and the wonder is that so many times one lives through these paroxysms.

The patient can hardly be questioned, can certainly not be carefully examined; and herein lies the advantage of the family physician who knows the patient and his heart, and in whom the patient has confidence. In fact, this confidence which such a patient has in the physician who has more or less frequently aided him in weathering these terrible attacks is alone the greatest boon the patient can have.

MANAGEMENT

The immediate conditions to meet are the rapid, fluttering heart, the nervous excitation and cardiac anxiety, and perhaps the most important of all, the vasomotor spasm that is often so pronounced. Physically we have, then, a heart with leaking or constricted valves; in either case more blood is entering the chambers of the heart than can be expelled in one contraction, while the peripheral resistance due to the spasm of the blood-vessels, because of fear, becomes greater every minute and tends still more to interfere with the peripheral circulation and the complete emptying of the heart of its surplus blood. Owing to the well-known stimulus to distention of hollow muscular organs, the heart contracts faster and faster. Soon, by some disarrangement of the inhibitory apparatus, the pneumogastric nerves, the heart loses its governor, and the beats increase to even 150 a minute, with irregular contractions, the blood being sent through the arteries with irregular force, as evidenced by the varying volume of the pulse. At this time, with or without cardiac pain, which upsets the rhythm of the heart, the patient becomes frightened at the feeling of impending demise, and the cerebral reflexes begin to add to the cardiac difficulty. The breathing becomes nervously rapid, besides that which is due to the rapid heart. The chill of fear is added to the already contracted peripheral vessels, and the surface of the body becomes cold, the extremities sometimes intensely so. Next it seems as if the strongly contracted arterioles begin actually to prevent some of the peripheral circulation, the blood is piled up in the large arteries, and the venous circulation becomes more and more sluggish, while the lips, finger-nails and forehead become cyanotic. Respiration becomes more rapid and deep, the inspiration being as strong as possible with every auxiliary muscle taking part, thus making the negative pressure in the chest aid in bringing the blood back through the veins. Part of the extra respiratory stimulus comes from the imperfectly aerated blood reaching the respiratory center.

Two factors may normally, without treatment, stop these paroxysms, and the "bad heart turn" may be cured spontaneously. The first of these is self-control. If the patient does not lose his head, by an effort of the will he saves himself from becoming nervous or frightened and therefore escapes the result of mental excitement; the increased peripheral blood-pressure from fear does not occur, and in a shorter or longer time the heart quiets down. The physician recognizes this power, and gives his patient immediate assurance that he will soon be all right; the patient who knows his physician immediately feels this assurance and is quickly improved.

The second factor in spontaneous cure of the heart attack is relaxation. The exhaustion from the respiratory muscular efforts together with the drowsy condition caused by the cerebral hyperemia and from the imperfectly aerated blood causes finally a dulling of the mental acuity, and the nervous excitement abates, which, with the exhaustion, gives a relaxation of peripheral arterioles; the resistance to the flow of the blood is removed, the surface of the body becomes warm, the heart quiets down by the equalization of the circulation, and the paroxysm is over.

DRUGS

The part the nervous system plays in this paroxysm is shown by the good result obtained from injections of morphin, even when there is no pain; hence the action of morphin is directly in line with the natural resolution of the symptoms: it quiets the nervous system, causes drowsiness, relaxes spasm, and thus causes increased peripheral circulation; many times this is the only treatment necessary.

During these heart attacks it is more than useless to administer any drug by the stomach, as in this condition there will be no absorption, even if there is no vomiting.

While morphin is generally indicated, as just suggested, a very large dose should not be given, lest the activity of the respiratory center is impaired (it is already in trouble), and undoubtedly death may easily be caused by an overaction of morphin during these heart attacks. The addition of atropin to the morphin will prevent depression from the morphin. Also, atropin sometimes quiets cardiac pain, but it will not steady the heart, may irritate it, and will increase vasomotor tension, although peripheral nerve irritation may be diminished. Hence a fair dose of morphin hypodermatically with a small dose of atropin, if respiratory depression is feared, is a physiologic method of bettering the condition.

In this kind of heart attack a drug that often acts well is nitroglycerin. It may be given hypodermatically in a dose of 1/200 to 1/100 of a grain, or a tablet may be dissolved on the tongue, and the dose be repeated once or twice at fifteen-minute intervals, until there is throbbing in the forehead, which shows that a sufficient amount of the drug has been administered. This headache will generally not last long. In the meantime the peripheral blood-vessels are relaxed, the surface of the body becomes warm, the heart quiets, and the attack is over. To hasten the action of nitroglycerin (that is, to equalize the circulation) a hot foot-bath is often valuable. Amyl nitrite may be inhaled with the same object in view, but the action is very intense, the prostration often severe, and unless there is angina pectoris, nitroglycerin is much better.

The symptoms of a heart attack may not be quite those above described; they may be those of sudden dilatation or semiparalysis of the heart, in which the prostration is intense and the patient is unable to sit up, although he may be leaning against several pillows. There is dyspnea, but the patient cannot aid respiration with the auxiliary muscles by holding the arms and shoulders tense or obtaining support from the arms; in fact, the arms are almost strengthless. The surface of the body may be warm, and the arms may be warm except the hands; the feet, ankles and legs may be cold. There is generally more or less cyanosis, although the face may be pale. The finger-nails often show venous stasis. In these cases the blood-pressure is subnormal, the pulse may be hardly perceptible, and there is none of the tension of the body from fear. The patient may be fearful, but he is completely collapsed. Such an attack may occur suddenly in a heart that is perfectly compensating, or it may accompany general edemas and dropsies.

If the emergency is excessively urgent, the lungs filling up with blood, moist râles beginning to occur, and frothy and blood-tinged sputum being coughed up, venesection may be indicated; combined with proper hypodermatic medication it may save life, and does at times. In fact, a patient who shows every sign of fatal cardiac collapse may be saved.

One of the best drugs to administer to such a patient is an aseptic ergot, injected intramuscularly. The drug of all drugs for future action (as it will not act immediately) is digitalis, and perhaps best, in this condition, in the form of digitalin. (See N. N. R., p. 88.) The dose of German digitalin may be from one-tenth to one-fourth grain, or even more. This dose, or a smaller dose, may be repeated in three or four hours, if deemed advisable. Whether digitalis shall be given at all, or how large the dose shall be depends on whether the patient has been taking digitalis in large quantities or not. He may already be overpowered with digitalis. In that case it would be contra-indicated.

Strophanthin, especially when given intravenously, has been found to be a quickly acting circulatory stimulant. The dose of strophanthin, Merck, ranges from 1/500 to 1/200 grain. The intravenous dose of strophanthin, Thoms, is about 1/130 grain. It should not be repeated within a day or two, if at all.

Atropin in a dose of 1/150 grain, and strychnin in a dose of 1/40 or 1/30 are valuable aids in stimulating the circulation under these conditions. The atropin should not be repeated. The strychnin may be repeated in three, four or five hours, depending on the size of the previous dose.

Of all quickly acting stimulants none is better than camphor in saturated solution in sterile oil as may be obtained in ampules. Alcohol is absolutely contra-indicated in the latter condition. In the former kind of heart attack, vasodilatation from a large dose of whisky or brandy may be of value. The dose should be large to cause immediate increased peripheral circulation, dilatation, and even a little stupefaction of the central nervous system, and it may be effectual in a way not dissimilar to the action of morphin.

TREATMENT OF BROKEN COMPENSATION

The consideration of this subject will include the following topics:

A. Hygiene.
B. Diet.
C. Elimination.
D. Physical measures.
E. Medication.
    1. Cardiac Tonics: Digitalis, strophanthus, caffein, strychnin.
    2. Cardiac Stimulants: Camphor, alcohol, ammonia.
    3. Vasodilators: Nitrites, iodids, thyroid extract.
    4. Cardiac Nutritives: Iron, calcium.
    5. Cardiac Emergency Drugs: Ergot, suprarenal active principle, pituitary active principle, atropin, morphin, and also some of the drugs already mentioned.

A. HYGIENE

Of all treatment for broken compensation or dilated heart nothing equals rest in bed. Sometimes it is the only treatment that is needed. The rigidness of this rest, the length of time that it should endure, and the period at which relaxation of such rest should be allowed depend entirely on the individual patient; no rule can be established. Most of the symptoms must disappear before exercise is allowed. Perhaps a not infrequent exception to the rule is when cardiac weakness, generally a myocarditis, develops in a patient after 50. It is not always wise to keep such a patient in bed; he may be rested and his exercise greatly restricted, but sometimes it is difficult to get him out of bed if he is kept there any length of time.

Fresh air, sunlight and anything else that makes the bedroom attractive and cheerful are essential and will aid in the recovery. The kind of nurse that is needed, trained, untrained, or a member of the family, and the amount of company or entertainment that the patient shall have must be entirely decided for the individual patient. The patient must be distinctly individualized and the proper measures taken to give mental and physical rest, to prevent excitement, worry, melancholia and depression, and to improve the general nutrition of the body as well as the condition of the heart.

Each occurrence of broken compensation in valvular disease causes another attack of cardiac weakness to occur with less excuse than before, and several serious attacks of broken compensation mean before long the loss of the heart muscle's ability to recover, so that permanent dilatation occurs.

B. DIET

The food given should be just sufficient for the needs of the body; the patient should not be overfed or underfed. Any large bulk of food or liquid should not be given. Pressure on the heart causes discomfort and is therefore inadvisable. Food that causes flatulence should be avoided. Theoretically the patient should receive a little meat, an egg or two, cereal or bread, a small amount of simple vegetables, a little fruit, often milk, a sufficient amount of non-effervescent water, perhaps a cup of chocolate or cocoa, a simple dessert, sometimes ice-cream; in other words, a varied, limited diet containing all the elements that are necessary to good nutrition. The diet should be varied from day to day to encourage the appetite.

It has for several years been recognized that a salt-free diet in dropsies due to disease of the kidneys is a valuable aid in causing absorption of such exudates and of preventing greater exudations. For this reason a salt-free diet is often ordered in dropsies occurring in valvular disease. Its value, however, is not so great as in kidney lesions, and if it causes hardship to the patient it should not be continued rigorously. On the other hand, large amounts of salt should of course be interdicted.

A most valuable aid in dropsies due to heart deficiencies is the so-called dry diet, which means that as little liquid as possible should be taken in order that the patient's blood may resorb the exudate in the tissues and not have the blood-vessels filled or overfilled with liquid from the gastro-intestinal tract. When dropsy is present, or even when serious pendent edemas are present, the patient should drink as little liquid as possible with his meals, and between meals should sip water rather than drink a large quantity of it. This is one of the reasons that a large milk diet, even with kidney disturbance due to cardiac lesions, is generally inadvisable. With cardiac or general circulatory weakness, a large amount of liquid to flush out the kidneys and the whole system, so long ordered for all kidney defects or mistakes in metabolism, is a serious mistake.

Whether it is better to give three or four small meals a day or to give a small amount of nourishment every three hours during the daytime must again depend on the individual and his ability to digest without fermentation and putrefaction or discomfort. As previously urged, not too much fluid, even milk, though it digest perfectly, should be given, as the greater the amount of fluid the greater the amount of work thrown on the heart.

C. ELIMINATION

A patient who has developed cardiac incompetency has always imperfect elimination. The skin, bowels and kidneys do not act sufficiently or well. The circulation in the skin is sluggish. The bowels are generally constipated, or there is diarrhea of the fermentative type. The amount of urine excreted is generally insufficient and likely to be concentrated and show various signs of imperfect kidney elimination. Therefore hot sponge baths, with perhaps warm alcohol rubs, are daily necessary. Gentle massage, generally in the direction to aid the circulation, will benefit the skin. If the skin is dry or in places scaly, oil rubs are of great benefit.

The bowels must be moved daily and sufficiently, but there should be no watery purging allowed or caused. If it seems advisable in the beginning of the treatment to give a calomel purge, it should be done, but such purging should ordinarily not be repeated, although occasionally a grain or two of calomel, combined with the vegetable laxatives needed, may act perfectly and without causing depression. Saline purgatives, or even laxatives, are generally not good treatment when there is cardiac weakness. The bowels should be moved by vegetable laxatives, as aloin, cascara sagrada, or some simple combination of either or both of these drugs.

Diuretics are often not satisfactory in cardiac insufficiency. The cardiac tonics that are given the patient, and the improvement of the heart from the rest in bed generally start the kidneys to secreting properly. A diuretic administered when the kidneys are suffering passive congestion from cardiac insufficiency does not generally act, and is therefore useless. If digitalis is administered, it will act as a diuretic; if caffein is deemed advisable, that will act as a diuretic. Squills may be administered, if it seems best. If for any reason the kidneys secrete less urine and become insufficient, the diet should quickly be reduced to a small amount of milk, cereal and water, and hot baths and local heat to the back should be inaugurated.

D. PHYSICAL MEASURES

Hydrotherapy is often of great value in restoring compensation by improving the surface circulation. Sponging with hot, tepid or cold water, as indicated, will increase the peripheral circulation and the normal secretions of the skin.

When compensation is perfect, in valvular lesions, more or less frequent warm baths are advisable, and often relieve the heart by equalizing the circulation. Cold sponging in the morning may be advisable, but may do harm when there is high tension; warm, not too hot, baths are of value. Anything is of value that improves the peripheral circulation and prevents the extremities from being cold.

The value of the Nauheim or other carbonated baths is perhaps often a question. They have seemed in many instances to aid in improving compensation in such patients as have been able to go abroad for the treatment. On the other hand, so many other regimens are ordered and inaugurated for these patients at these "cures" that it is hard to decide how much benefit the baths have really done. At home the artificial carbonated or carbonic acid baths do not seem to be of great value. Baths and bathing can do harm, and the decision as to which hydrotherapeutic measure shall be used and the extent to which they shall be used can be made only after careful observation of the patient by the physician.

Gentle massage while the patient is in bed is of undoubted value; more vigorous massage is later often of value, provided there is no arteriosclerosis. As the patient grows stronger and the circulation improves, the muscles are kept in good condition during the enforced rest by massage. When properly applied, it promotes not only the venous return circulation, but also the lymphatic circulation; it often removes muscle aches and muscle tire and restlessness.

While the patient is still in bed, various resistant exercises are of value, and should be begun. These tend to prepare the patient for his later greater activities; in other words, the surprise to the heart when the patient begins to sit up and walk is not so great if he has previously taken these exercises. Later, when the patient is ambulatory he should by gradual gradation walk a little more about the house and take a few steps of the stairs at a time, until gradually he is able to mount the whole flight. Later he should take outdoor exercise, and when his heart has become compensated for ordinary work, he should be given gradually graded hill-climbing with the idea of increasing his reserve cardiac power. If it is found that these increased exertions cause him to have pain or a more rapid heart than is excusable, even after persisting for a few days, the attempt to increase this reserve power of the heart should be abandoned. There is probably, at least at that particular time, considerable myocarditis, although the heart may eventually recuperate still more. Pushing it to overexertion, however, will not accomplish improvement.

Graded exercise was first used scientifically by Oertel and Schott, and has been for years designated by their names. Modifications of their rigid rules are generally advisable.

E. MEDICATION

1. *Cardiac Tonics.*—Digitalis: There is no drug that can take the place of digitalis in loss of compensation in chronic valvular disease. It acts specifically for good, and it has its greatest success in the valvular lesions that cause enlargement of the left ventricle, on which it acts the most intensely. It also acts for good on the right ventricle. It has but little action on the auricles. This is simply a question of muscle; the part that has the greatest amount of muscle will receive the greatest benefit from digitalis, and the parts that have the least, the least benefit. The heart muscle is somewhat similar to other muscles; when we attempt athletic improvement in any muscle of the body we "train" by stimulating it moderately at first, and are careful not to overwork it; the object, then, is to train the heart muscle. For this reason large doses of digitalis should ordinarily not be given to overstimulate suddenly an overworked and weak heart. While in some instances it has been declared that digitalis should be rapidly pushed to the full extent and then dropped for a time, careful experience shows that this method is often not tolerated, sometimes does positive harm, and has at times seemed to hasten death.

Another valuable activity of digitalis is in slowing the heart by action on the pneumogastric nerves. A dilated heart has lost more or less of its regulating mechanism; this is the cause of its irregularity and its increased rapidity. The action of digitalis in slowing the heart, giving it a longer rest, and preventing it from acting irregularly is of great value. This prolonged rest or diastole of the heart allows the circulation in the coronary arteries to become normal, and the nutrition and muscle tone of the heart improves. Digitalis also increases the blood-pressure, not only by improving the activity of the heart, but also by causing some contraction of the arterioles. This feature of digitalis action in arteriosclerosis renders its use sometimes a question of careful decision. The dose of digitalis under such a condition should not be large. It may be indicated, however, and may do a great deal of good, and it does not always increase the blood-pressure.

If the patient is sufficiently ill to require the best action of digitalis, an active preparation should be obtained. It was long supposed that the infusion presented activities that could not be furnished by the tincture of digitalis. This seems not to be true. The greater value of the infusion is generally because it is freshly made and active; the tincture that had been used previously in a given case was old and useless; also most physicians give a larger dose of the infusion than they ever do of the tincture. Owing to the uncertainty of the value of the digitalis leaves found in the various drug shops, however, and to variations in the preparation of the infusion, it is generally better to use a tincture of known character. The beginning dose of such a tincture should generally not be more than 5 drops, and it should not be repeated more frequently than once in eight hours. It is generally advisable, in two or three days, to increase this dose to 10 drops once in twelve hours, later perhaps to 15 drops twice a day, and still later to 20 drops once a day. This amount may then be decreased gradually, if the action is satisfactory. In other words, enough should be given to procure results, and then the dose should be brought down to what seems sufficient and best, administered once a day. The frequency advised in the administration of this drug is because it is eliminated slowly. Its greatest action develops a number of hours after it has been taken, and then the action lasts for many hours; the administration of digitalis once in twenty-four hours is perfectly satisfactory for many patients, and more satisfactory than any more frequent administration. On the other hand, some patients do better on a smaller dose once in twelve hours. This frequency is always sufficient.

Digipuratum, digitol, a fat-free tincture, and digalen, claimed to contain "amorphous digitoxin," three proprietary preparations accepted by the Council on Pharmacy and Chemistry for inclusion in N. N. R. may be employed. They are standardized preparations and may thus be more satisfactory than some pharmacopeial preparations of digitalis, although their claims to lessened emetic action are not borne out by recent experiments of Hatcher and Eggleston.

Digipuratum may be obtained in tubes of twelve tablets. The advice has been given for patients with loss of compensation to receive four tablets the first day, three the second, three the third, and two the fourth day. This, however, is generally an overdosage. The most that should generally be given is one of these tablets in twelve hours.

Digitol is a fat-free tincture of digitalis which is physiologically standardized and which bears on each package the date of manufacture. The dose is 0.3 to 1 c.c. (5 to 15 minims).

Digitalinum, one of the active principles of digitalis, is not very satisfactory. It may be given hypodermatically, but often causes irritation, and the proper dose, and its value are likely to be uncertain.

Digitoxin, another active principle of digitalis, has been declared by some investigators to be harmful, and to be likely to cause serious disturbance of a damaged heart. Other investigators have stated that it acts for good. Digalen, said to contain "amorphous digitoxin," whatever that may be, has been used. The dose suggested in the instructions furnished with the sealed package is generally too large. While it sometimes undoubtedly causes cardiac pain, so may every preparation of digitalis. Digitoxin, and also probably this preparation, do not represent the whole value of digitalis, and in broken compensation digitalis itself, or some preparation embodying the majority of its activities, should be given. Digitoxin, however, is often valuable in conditions of cardiac debility or slight weakening in patients who do not have dilated hearts or edemas. The most satisfactory dose of digalen is from 5 to 10 drops once or twice in twenty-four hours.

Digitalis should not be used when there is fatty degeneration of the heart; it should ordinarily not be used when there is arteriosclerosis, and very rarely, if ever, when it is decided that there is coronary disease. Whether digitalis should be used when there is considered to be much myocardial degeneration is a question for individualization. One can never be sure that the heart muscle is so thoroughly degenerated that no part of it would be benefited by digitalis when compensation is lost; therefore, many times, especially if other drugs have failed, small doses of digitalis should be tried, to see if the heart will respond. Large doses or frequent doses would be contra-indicated.

The signs of overaction of digitalis are nausea, vomiting, a diminished amount of urine, a tight, band-like feeling around the head, perhaps occipital headache and coldness of the hands and feet, or frequently of one extremity only, combined with a feeling of numbness. The pulse is generally reduced to sixty or less a minute. Such symptoms require that digitalis be immediately stopped. They are the primary signs of cumulative action.

While many patients with ordinary dosage of digitalis may take the drug for months and years without ever showing cumulative action, other patients show this effect quickly. They are likely to be those in whom the kidneys are not perfect. The signs of such undesired action may develop slowly, as suggested by the symptoms just enumerated, or they may develop suddenly. The pulse becomes rapid and irregular, the heart action weak, there is severe backache in the region of the kidneys, a greatly diminished amount of urine, or even partial suppression, severe headache, vomiting, cold extremities and shiverings.

The treatment of such an undesired behavior of digitalis is, of course, to stop the drug immediately, give saline laxatives, hot sponging or hot baths, nitroglycerin and perhaps alcohol.

Strophanthus: Strophanthus cannot be compared with digitalis, except when the glucosid, strophanthin, is administered subcutaneously or intravenously. Strophanthus is given either in the form of the tincture, or as strophanthin. It has been shown that in neither of these forms, when the drug is administered by the stomach, is the muscle of the heart or the blood-vessels much acted on. In other words, compensation could not be restored by strophanthus. In emergencies of serious cardiac failure strophanthin intravenously has been shown apparently to save life. It acts quickly, and its power of stimulating the heart and contracting the blood-vessels lasts for many hours. It is rarely, however, that the dose should be repeated, and then not for twenty-four hours; but during that twenty-four hours the patient may be saved until other drugs which act more slowly have been absorbed, or perhaps until the emergency has passed.

There are many, however, who believe that they obtain considerable value from the tincture of strophanthus, and there seems to be no doubt that although strophanthus, given in the form of the tincture and by the mouth, may not increase the muscle power of the heart, it many times acts as a satisfactory cardiac sedative. Under its action the patient becomes less nervous, the heart often acts more regularly, and the low blood-pressure may improve. In other words, we should not be quite ready to discard the internal use of the tincture of strophanthus.

The tincture of strophanthus readily deteriorates, and the preparation ordered should be known to be a good one.

Caffein: This should not be given or allowed, even in the form of tea or coffee, to patients who have valvular lesions with perfect compensation, as it is a nervous and cardiac stimulant and may cause a heart to become irritable. It raises the blood-pressure slightly, acts as a diuretic, and hence is often of great value when used medicinally. It should be ranked as a stimulotonic to the heart. It increases its activity, but gives it a little more strength. It will rarely slow a rapid heart; it will often stimulate a sluggish, slow heart; it may increase the irritability of an irritable heart. As it is a cerebral stimulant, it should not be given late in the afternoon or evening, as it may prevent sleep.

The most frequent form of caffein used is the citrated caffein. The dose is 0.1 gm. (1½ grains) two or three times in the early part of the day, or 0.2 gm. (3 grains) once or twice during the morning. A cup of coffee may be given the patient medicinally, as a substitute for the drug, an ordinary cup of strong coffee containing between 2 and 3 grains. Other preparations of caffein may be selected if desired, or a soluble preparation may be given hypodermatically.

Caffein is indicated if digitalis is contra-indicated or does not act satisfactorily, and the patient is not nervously excited, but perhaps is stupid or apathetic, and also when diuresis is desired.

Strychnin: This is a valuable stimulator and heart tonic when properly used. It promotes muscular activity of the heart much as it promotes all muscular activities. It awakens nervous stimuli and nervous transmissions to normal in all sluggish nerve functions. If for these reasons the heart acts more perfectly, and the nutrition of the heart muscle improves it acts as a cardiac tonic. Many times, by improving the action of the heart, and also by the action of the drug on the vasomotor center, the pressure in the peripheral circulation may be increased. On the other hand, strychnin in the low blood-pressure of serious illness, such as pneumonia, by no means always raises the blood-pressure.

It should not be forgotten that strychnin is a general nervous stimulant, especially of the spinal cord. If it makes a nervous patient more nervous, or a quiet patient restless and irritable, it is acting for harm and should be stopped, just as caffein under the same conditions should be stopped. Strychnin may cause diminished secretion of the skin. This is not frequent, but it does occur. It may prevent the patient from sleeping. If such be the fact, strychnin is not acting for good in a patient who has cardiac weakness.

INDICATIONS FOR STRYCHNIN

Strychnin is a much overused drug. It is now given for almost everything and during almost every disease. It is true that the administration of strychnin is largely due to the evolution of the age in which we are now living. We have ceased to purge and bleed and sweat, and to give large doses of aconite or veratrum viride, we have ceased to starve the patient too long; we have ceased to load him with alcohol to the point of circulatory prostration, and we have recognized that he must be braced from start to finish; strychnin is the drug that has been used for this purpose, and, as above stated, overused. Strychnin given too frequently or in too large doses for a laboring heart can prevent its proper rest; the diastole is shortened and the relaxation of the heart is incomplete, its nutrition suffers, or even irregular and fibrillary contractions of a weak heart may apparently be caused. While a large dose of strychnin, even to one-twentieth grain hypodermatically, may be used once in serious emergency when it is deemed the drug to use, a dose larger than one-thirtieth grain hypodermatically is rarely indicated, the frequency of such a dose should seldom be more than once in six hours, and a smaller dose of strychnin may act more satisfactorily.

Strychnin is indicated when the heart is acting sluggishly and the contractions seem incomplete and when digitalis either is not indicated or is not acting perfectly. Small doses of strychnin may aid such a heart during the administration of digitalis. In many instances in which digitalis is contra-indicated, strychnin is of marked value. This is typically true in fatty hearts, and may be true in arteriosclerosis, in which it often does not increase the blood-pressure at all.

2. *Cardiac Stimulants.*—A cardiac stimulant is a drug that makes the heart beat more strongly and the frequency becomes more nearly normal. The drugs named as cardiac stimulants, however, camphor, alcohol and ammonia, do not leave a heart better than they found it—they are not cardiac tonics.

Camphor: This is one of the best cardiac stimulants that we possess. It is a quickly acting nervous and circulatory stimulant, acting principally on the cerebrum and causing a dilatation of the peripheral blood-vessels. No subsequent weakness follows after a dose of camphor. Too much will make a patient wakeful, a little often quiets nervous irritability. It should be used as a cardiac stimulant during serious illness more frequently than it has been, and during the endeavor to make a non-compensating heart again compensatory camphor will often act for good. The dose is 2 teaspoonfuls of the camphor-water every three or four hours, as deemed advisable. Each teaspoonful represents a little more than ¼ grain of camphor. The spirits of camphor of course may be used, if preferred.

For cardiac emergencies ampules of sterile saturated solutions in oil are now obtainable and are valuable. Such hypodermatic stimulation acts quickly, and may be repeated every half hour for several times, if the patient does not respond. The solution should be injected slowly, and as a rule intramuscularly.

Many times while other measures are being used to repair a broken compensation, camphor makes a splendid circulatory and nervous bracer. Camphor has long been used as a so-called antispasmodic in hysteric or other nervously irritable persons. It really acts as a stimulant to the higher centers of the brain, promoting more or less nervous control. Perhaps its ability to increase the peripheral circulation may be one of the reasons that it seems at times to be almost a nervous sedative by relieving internal congestion. As just stated, after the camphor action is over there is no depression. This is not true of alcohol.

Alcohol: It is of course now generally understood that alcohol is not a cardiac stimulant in the sense of its being more than momentarily helpful to a weak heart. If alcohol is pushed when a heart is in trouble, the secondary vasodilatation and more or less nerve prostration and muscle debility will cause greater circulatory weakness than before it was administered. To obtain cardiac stimulation from alcohol it must be given in strong solutions, generally in the form of whisky or brandy, for local irritation of the mouth, esophagus and stomach; reflexly the heart is stimulated and the blood-pressure rises. As soon as complete absorption has taken place the blood-pressure falls. For continuous stimulation another dose of alcohol must be given before this depression occurs. This may be in from one to three hours. To continue such stimulation the dose of alcohol must be increased. The future of such treatment means an alcoholic sleep with depression, alcoholic excitement that is not desired, or profound nausea and vomiting, with peripheral relaxation and cold perspiration. Obviously none of these conditions is desirable; but in arteriosclerosis, or when the blood pressure is high and the heart labors under the disadvantage of contracting against an abnormal circulatory resistance, alcohol may act perfectly to relieve this kind of circulatory disturbance. In this condition the alcohol should not be given concentrated, and as soon as it is thoroughly absorbed vasodilatation occurs, peripheral circulation and therefore warmth is increased, and the heart is relieved of its extra load. In such instances, in proper doses not too frequently repeated, rarely more than 1 or 2 teaspoonfuls every three hours, alcohol is a valuable drug. Such good action of alcohol is often seen when the surface of the body is cold from chilling or the extremities are cold from vasomotor spasm. A good-sized dose of alcohol, best given hot, equalizes the circulation and acts for good. On the contrary, it is obvious that, if the patient is cold from collapse and there is cold perspiration and very low blood-pressure, alcohol is not the drug indicated, although one dose may be of benefit while other more slowly acting cardiac tonics or stimulants are being administered.

During serious prolonged illness and when the patient has not had sufficient food and is not taking sufficient food, alcohol in the form of whisky or brandy, not more than a teaspoonful every three hours, acts as a necessary food, and will more or less prevent acidosis from starvation.

It will be seen that alcohol, except possibly in a single dose occasionally, or for some special reason, is rarely indicated in broken compensation.

When alcohol is administered regularly, whether during a fever process or for any other reason, if it causes a dry tongue, cerebral excitement, flushed face and a bounding pulse, or if there is the odor of alcohol on the breath, the dose is too large, and alcohol is contraindicated.

Ammonia: In the form of ammonium carbonate or the aromatic spirits of ammonia this has long been used with clinical satisfaction as a cardiac stimulant. Probably, however, it is seldom wise to use ammonium carbonate. It is exceedingly irritant, and constantly causes nausea, perhaps vomiting, and often heart-burn or other gastric disturbance. It has no value over the pleasanter aromatic spirits of ammonia, which is essentially a solution of ammonium carbonate. The dose of the aromatic spirits is anywhere from a few drops to half a teaspoonful, given with plenty of water. It is a quickly acting stimulant, with an effect much like alcohol, followed by very little or no depression. It is more of a cerebral irritant than alcohol, and probably has but few, if any, advantages over camphor.

When but little nutriment has been taken for some days, it may be a chemical question, since ammonium compounds so readily form and become cerebral irritants, whether any more ammonium radicals should be given the patient. This is especially true with defective kidneys. In these conditions camphor is better.

To continue the action of ammonia it may be given every hour for several doses and then less frequently.

3. *Vasodilators.*—In various conditions of high blood-pressure, arteriosclerosis and even during the sthenic stage of a fever, vasodilators may be indicated. The most important are nitrites, iodids and thyroid extracts. Alcohol, as above stated, may act as a vasodilator. Aconite and veratrum viride are now rarely indicated, although possibly aconite should be used when there is high tension and the heart is acting irritably and stormily.

Of the nitrites, no preparation seems to act more satisfactorily than nitroglycerin (trinitrin, glyceryl, nitratis, glonoin). Its action may not be so prolonged as other forms of nitrite such as sodium nitrite or erythrol tetranitrate, but it is not irritant, and only a little less rapid than nitrite of amyl, and although the marked dilatation lasts but a short time, often apparently only for minutes, still, when frequently repeated or given a few times (from four to six) in twenty-four hours, it frequently keeps the blood-pressure lower than it would be without the drug. In diseases of the heart the sudden vasodilatation caused by amyl nitrite inhalations is indicated only in angina pectoris. When the surface of the body tends to be cold, however, when the peripheral blood-pressure is increased and the heart is laboring, nitroglycerin in small doses is valuable. The dose may be from 1/400 to 1/100 grain, dissolved on the tongue or given hypodermatically for quick action, or given by the mouth for more prolonged action. In sudden cardiac dyspnea nitroglycerin sometimes acts specifically, especially when there is asthma. When a drop or two of the official spirits, which is a 1 per cent. solution, is given on the tongue, or a soluble tablet of 1/100 grain is dissolved on the tongue, the action is almost as rapid as though the dose had been administered hypodermatically. Many times when such increased peripheral circulation is desired and alcohol seems indicated, nitroglycerin in small doses will act as well. It cannot be termed a cardiac stimulant, although many times a heart acts better and the pulse is fuller and stronger after nitroglycerin than before. It should not be used, except if specially indicated, in broken compensation or in other myocardial weakness.

Iodids: These have no immediate action. The vasorelaxation that often occurs from iodid is quite likely due to the stimulation of the thyroid gland by the iodin, and the thyroid gland secrets a vasodilating substance. Small doses of iodid, however, when indicated in various kinds of sclerosis, have seemed to lower blood-pressure. While large doses may have more of this action, they are not now under consideration, and large doses are rarely indicated. Too much iodid has been given for many conditions. If the indications for an iodid are present, such as sclerosis anywhere, or unabsorbed inflammatory products, exudation in or around the heart, or an apparent insufficiency of the thyroid, from 0.1 to 0.2 gm. (1½ to 3 grains) once or twice in twenty-four hours, after meals, is all that is required to give the action desired, and the circulation is benefited. It is sometimes a question whether small doses of iodid are not actually stimulant to the heart, possibly through the action on the thyroid gland.

Thyroid Extract: In slow hearts and in sluggish circulation, often in old age, quite frequently in arteriosclerosis and in every condition of insufficient thyroid secretion (these instances are frequent), small doses of thyroid extract will benefit the circulation. Its satisfactory action is to increase the cardiac activity, slightly lower the blood-pressure, and increase the peripheral circulation and the health of the skin. If it causes tachycardia, nervous excitement, sleeplessness or loss of weight, it is doing harm and the dose is too large, or it is not indicated. The dose for the cardiac action desired is 2 or 3 grains once a day, continued for a long period.

When an improved peripheral circulation is desired, and especially when a reduction of the pressure in the heart is desired and a diminished amount of blood in overfilled arteries is indicated, the value of the sitz-bath, hot foot-baths, warm liquids (not hot) in the stomach, and warm, moist applications to the abdomen should all be remembered.

4. *Cardiac Nutritives.*—Iron: Nothing is of more value to a weakened heart muscle, when the nutrition is low, the patient anemic, and the iron of the food not properly metabolized, than tonic doses of some iron salt. It has frequently been repeated, but should constantly be reiterated, that there is no physiologic reason or therapeutic excuse for the patient to pay a large amount of money for some organic iron preparation. Small doses of an inorganic salt act perfectly, and nothing will act better. As previously suggested, a drop or two of the tincture of iron, a grain or two of the reduced iron, or 2 or 3 grains of the saccharated oxid of iron, given once or twice in twenty-four hours, is all the iron

the body needs from the points of view of the blood and the heart.

Calcium: It has lately been learned that calcium is an element that a heart needs for perfect activity. Many patients who are ill lose their calcium, and they may not receive a sufficient amount of it unless milk is given them. Even if such patients are taking milk, a heart and the whole general condition sometimes seem to improve when calcium is added to the diet. It may be given either in the form of lime-water, calcium lactate or calcium glycerophosphate. If a medium-sized dose is given three or four times in twenty-four hours, it is sufficient and will often act for good.

Whether calcium can do harm in a chronic endocarditis or an arteriosclerosis to offset the value that it seems to have in quieting the nervous system and in being of value to a weak or nervously irritable heart is a question that has not been decided. Theoretically lime should not be given when there is a tendency to calcification, or when a patient is past middle age. Lime seems to be essential to youth, and to the welfare of nervous patients.

EMERGENCIES

5. *Cardiac Emergency Drugs.*—Besides some of the drugs already mentioned (such as camphor hypodermatically, nitroglycerin when indicated, strophanthin hypodermatically or intravenously, caffein and strychnin), often ergot, suprarenal vasopressor principle, pituitary vasopressor principle, atropin and morphin should be considered.

When there is low blood-pressure, venous stasis, pulmonary congestion, cyanosis and a laboring, failing heart, intramuscular injections of ergot, with or without coincident venesection, may be the most valuable method of combating the condition. Life has been saved in this kind of sudden acute cardiac failure in valvular disease. When venesection is not indicated in certain conditions of low blood-pressure and heart failure, ergot has saved life. It causes contraction of the blood-vessels and seems to tone the heart. Incidentally it quiets the central nervous system. If the blood-pressure is much increased by it, the ergot should not be repeated, as too much work should not be thrown on the heart muscle. Often, howhowever, it may be administered hypodermatically with advantage in aseptic preparation as offered in ampules, at the rate of one ampule every three hours for two or three times, and then once in six hours for a few times, the future frequency depending on the indications.

Epinephrin and Pituitary Extract: The blood-pressure-raising substance of the adrenals or of the pituitary gland (hypophysis cerebri) has been much used in heart failure. These substances certainly would not be indicated in high blood-pressure; they are indicated in low blood-pressure. They have been given intravenously; they are frequently given hypodermatically. They often act rapidly when a solution in proper dose is dropped on the tongue. The blood-pressure rise from epinephrin is quickly over; that from the pituitary extract lasts longer. In large doses or when it is too frequently repeated epinephrin depresses the respiration. Pituitary extract acts as a diuretic. Sterilized solutions of both, put up in ampules ready for hypodermatic medication are obtainable, the strength offered generally being 1 part of the active principle to 10,000 of the solution. Hypodermatic tablets of epinephrin may also be obtained. Stronger solutions of 1 part to 1,000 may be dropped on the tongue, or tablets may be dissolved on the tongue. The blood-pressure is temporarily raised and the heart stimulated by these treatments, but epinephrin is not used so often for cardiac failure as it was a short time ago.

The most satisfactory action, especially from the epinephrin, is from small doses frequently repeated. Sometimes in serious emergencies it has been found to be of value when given intravenously in physiologic saline solution. The dose, of course, should be very small. In circulatory weakness in acute illness epinephrin has been given regularly, a few drops (perhaps the most frequent dose is 5) of a 1:1,000 solution, on the tongue, once in six hours. Such a dosage may be of value, and certainly is better than the administration of too much strychnin. Much larger or more frequent doses are likely, as just stated, to depress the respiration.

Besides the small amount of blood-pressure-raising substance secreted by the hypophysis cerebri, it has not been shown that any other gland of the body furnishes vasopressor substance except the adrenals. Contrary to the generally accepted belief that this vasopressor principle of the adrenals was necessary for the proper tonus of the vasomotor system, recent experiments by Hoskins and McClure[6] show that this pressor element of the adrenal secretion is unimportant; they say, "Adrenal secretion is not, therefore, a direct factor in the maintenance of the tonus of the vasomotor system." There must be some way of amalgamating previous experimentation and beliefs with these newer investigations. It would be absurd to believe that as active a blood-pressure-raising substance as epinephrin is of no value to the economy and plays no part in the function in which it seems to be the most physiologically active. More confirmatory experimentation is needed before we can believe that adrenal secretion does not play an important part in the blood-pressure of the arterioles of the abdomen as well as of the entire body.

Atropin: When there is great cardiac weakness atropin may be used to advantage. The dose is from 1/200 to 1/150 grain hypodermatically, not repeated in many hours. It will whip up a flagging heart, more or less increase the blood-pressure, cause cerebral awakening, and may often be of value. If there is any idiosyncrasy against atropin, if the throat and mouth are made intensely dry, or if there is serious flushing or cerebral excitement, the dose should not be repeated.

---

[6] Hoskins and McClure: Arch. Int. Med., October, 1912, p. 343.

Morphin: This would rarely be considered as an emergency drug in cardiac weakness. A small dose of it, not more than one-eighth grain, especially if combined with atropin, will often quiet and brace a weak heart, when there is cardiac pain in particular.

Just which drug or drugs should be used and just which are not indicated can never be specifically outlined in a text-book, a lecture or a paper. The decision can be made only at the bedside, and then mistakes, many times unavoidable, are often made.

In all conditions of shock with cardiac failure the blood-vessels of the abdomen and splanchnic system are dilated, and more or less of the blood of the body is lost in these large veins, and the peripheral and cerebral blood-pressure fails. The advantage in such a condition of firm abdominal bandages, and of raising the foot of the bed or of raising the feet and legs, need only be mentioned to be understood.

It is a pretty good working rule, in cardiac failure, not to do too much. On the other hand, life is frequently saved by proper treatment, and the physician repeatedly saves life as surely as does the surgeon with his knife.

CONVALESCENCE

When compensation has been restored, the patient may be allowed gradually to resume his usual habits and work, provided that these habits are sensible, and that the work is not one requiring severe muscular exertion. Careful rules and regulations must be laid down for him, depending on his age and the condition of his arteries, kidneys and heart muscle. It should be remembered that a patient over 40, who has had broken compensation, is always in more danger of a recurrence of this weakness than one who is younger, as after 40 the blood-pressure normally increases in all persons, and this normal increase may be just too much for a compensating heart which is overcoming all of the handicap that it can withstand. Such patients, then, should be more carefully restricted in their habits of life, and also should have longer and more frequent periods of rest.

The avoidance of all sudden exertion in any instance in which compensation has just been restored is too important not to be frequently repeated. The child must be prevented from hard playing, even running with other children, to say nothing of bicycle riding, tennis playing, baseball, football, rowing, etc. The older boy and girl may need to be restricted in their athletic pleasures, and dancing should often be prohibited. Young adults may generally, little by little, assume most of their ordinary habits of life; but carrying heavy weights up-stairs, going up more than one flight of stairs rapidly, hastening or running on the street for any purpose, and exertion, especially after eating a large meal, must all be prohibited. Graded physical exercise or athletic work, however, is essential for the patients' future health, and first walking and later more energetic exercise may be advisable.

These patients must not become chilled, as they are likely to catch cold, and a cold with them must not be neglected, as coughing or lung congestions are always more serious in valvular disease. Their feet and hands, which are often cold, should be properly clothed to keep them warm. Chilling of the extremities drives the blood to the interior of the body, increases congestion there, and by peripheral contraction raises the general blood-pressure. A weak heart generally needs the blood-pressure strengthened, but a compensating heart rarely needs an increase in peripheral blood-pressure, and any great increase from any reason is a disadvantage to such a heart. The patient should sleep in a well-ventilated room, but should not suffer the severe exposures that are advocated for pulmonary tuberculosis, as severe chilling of the body must absolutely be avoided.

The peripheral circulation is improved, the skin is kept healthy, the general circulation is equalized, and the heart is relieved by a proper frequency of warm baths.

Cold baths are generally inadvisable, whether the plunge, shower or sponging; very hot baths are inadvisable on account of causing a great deal of faintness; while warm baths are not stimulating and are sedative. The Turkish and Russian bath should be prohibited. They are never advisable in cardiac disease. With kidney insufficiency, body hot-air treatment (body-baking), carefully supervised, may greatly benefit a patient who has no dilatation of the heart and who has no serious broken compensation. Surf-bathing and, generally, sea-bathing and lake-bathing are not advisable. The artificial sea-salt baths and carbon dioxid baths may do some good, but they do not lower the general blood-pressure so surely as has been advocated, and probably no great advantage is apt to be derived from such baths. If a patient cannot properly exercise, massage should be given him intermittently.

Any systemic need should be supplied. If the patient is anemic, he should receive iron. If he has no appetite, he should be encouraged by bitter tonics. If sleep does not come naturally, it must be induced by such means as do not injure the heart.

Perhaps there is no better place in this series on diseases of the heart to discuss the diet in general and the resort treatment than at this point, as the question is one of moment after convalescence from a broken compensation, at which time every means must be inaugurated to establish a reserve heart strength to overcome the daily emergencies of life.

DIET AND BATHS IN HEART DISEASE

The diet in cardiac diseases has already incidentally been referred to. The decision as to what a patient ought to eat or drink must often be modified by just what the patient will do, and, as we all know, it is absolutely necessary to make some concessions in order for him to aid us in hastening his own recovery or in preventing him from having relapses. Consequently, we cannot be dogmatic with most patients with chronic heart disease. Parents should be prohibited from allowing children or adolescents with heart disease to drink tea, coffee or any alcoholic stimulant. The young boy and young man must absolutely be prohibited from indulging in tobacco at all. There is no excuse for allowing these stimulants or foods in such cases. If the patient is older and has been accustomed to tea and coffee, one cup of coffee in the morning may be allowed, provided the "dekofa" preparation is not found satisfactory. Dekofa is coffee with most of the caffein removed.[7] Whether a small cup of coffee or a cup of tea is allowed at noon is again a matter for individualization; they should rarely be allowed after the noon meal. In a patient who has been accustomed to alcohol regularly (generally an older patient), careful judgment should be used in deciding whether or not a small amount of alcohol daily should be allowed. It should never be in large amounts, even of a dilute alcohol like beer; it may be a weak wine; it may be a small amount of diluted whisky, if it seems best. Ordinarily the patient is better without it. If he is used to smoking, and a small amount does not raise the blood-pressure much, it may do him no harm to smoke a small mild cigar once or twice a day. On the other hand, if a hard smoker suddenly has heart failure, whether from exertion, from chronic disease or from acute illness, a small amount of smoking is of advantage as it tends to remove cardiac irritability, to raise the blood-pressure, and actually to quiet and improve the circulation. It is unwise during acute circulatory failure to take tobacco away entirely from a chronic tobacco user.

---

[7] See Dakofa, a Caffein-Poor Coffee, THE JOURNAL A. M. A., July 1, 1911, p. 37.

The character of the food which each patient should receive depends on his blood-pressure and his age. The older person with a tendency to high blood-pressure should have the protein (especially meat) reduced in amount, as any putrefaction in the intestine with absorption of products of such maldigestion irritates the blood-vessels, raises the blood-pressure, and injures the kidneys. On the other hand, a young patient should receive a sufficient meat diet rather than be overloaded with vegetables and starches, to the easy production of fermentation and gas. Flatulence from any cause must be avoided. It dilates the stomach and intestines, causing them to press on the diaphragm, so that the heart and respiration are interfered with. Also, an increased abdominal pressure, especially if there is any edema or dropsy, is bad for the circulation. A distended, tense abdomen is serious in cardiac failure. On the other hand, a flaccid, flabby, lax abdomen should be well bandaged in cardiac failure with low blood-pressure.

Children do well on a milk diet, but it should be remembered that excessive amounts of any liquids, even milk and water, are inadvisable, if the circulation is poor and there is a tendency to dropsy. It has been recommended at times to limit a patient's diet for a week or so to a small amount of milk, not more than a quart in twenty-four hours. If such a patient is in bed and does not require carbohydrates, sugars or stronger proteins or more fat, such a restricted diet may aid in establishing circulatory equilibrium, although he will lose in nutrition. The excretory organs are relieved by the decreased amount of excretory products, the digestive system is rested and the circulation is improved. Such a limited diet should not be tried longer than a week, but it may be the turning-point of circulatory improvement.

The ordinary diet for a convalescing heart patient should be small in bulk, of good nutritive value, and should represent all of the different elements for nutrition. This means a small amount of meat, once a day to older patients, twice a day to those who work hard or for young patients; such vegetables as do not cause indigestion with the particular patient, and these must be individualized; such fruits as are readily digested, especially cooked fruits; generally plenty of butter, cream, olive oil if the nutrition is low, and milk, depending on the age of the patient or the ease with which it is digested. Soups, on account of their bulk and low nutritive value, should be avoided. Anything that causes indigestion, such as fried foods, hot bread, oatmeal or any other gummy, sticky, gelatinous cereal should be avoided; also spices, sauces and strong condiments. Anything that is recognized as especially loaded with nuclein and xanthin bodies, such as liver, sweetbreads and kidneys, should be prohibited, as tending to cause uric acid disturbance; and the more tendency to gout or uric acid malmetabolism the more irritated are the arteries and the more disturbed the blood-pressure. Sugars should be used moderately, unless the patient is thin and feels cold, in which case more may be given, provided there are no signs of gout or disturbed sugar metabolism. Very cold and very hot drinks or foods should be avoided.

Many times these patients have a diminished hydrochloric acid secretion, and such patients thrive on 5 drops of dilute hydrochloric acid in water, three times a day, after meals. When their nutrition has improved and the digestion becomes perfect, hydrochloric acid will generally be sufficiently secreted and the medication may be stopped.

If the patient is overweight, this obesity must be reduced, as nothing more interferes with the welfare of the heart than overweight and overfat. In these cases the diet should be that required for the condition.

If there are edemas, or a tendency to edemas, the decision should be made whether salt (sodium chlorid) should be removed from the diet. Unless there is kidney defect, probably it need not be omitted, and a long salt-free diet is certainly not advisable. This salt-free diet has been recommended not only in nephritis and heart disease, but also in diabetes insipidus and in epilepsy. It is of value if there is edema in nephritis; it is of doubtful value in heart disease; it is rarely of value in diabetes insipidus; and in epilepsy its value is probably in allowing the bromid that may be administered to have better activity in smaller doses, the bromin salt being substituted in the metabolism for the chlorin salt.

THE RESORT TREATMENT OF CHRONIC HEART DISEASE

In line with the continued growing popularity of special resorts and special cures for different types of disease, resort or sanatorium treatment for chronic heart disease has grown to considerable popularity during the last twenty years or more. The most popular of these resorts owe their success to the personality of the physicians, who have made heart disease a life study.

Perhaps the most noted of these resorts for the cure of heart disease is that at Bad Nauheim, Germany, which was inaugurated by Dr. August Schott and Prof. Theodore Schott, and is now conducted by the latter, Dr. August Schott having died about twelve years ago. Hundreds of patients and many physicians have testified to the value and benefit of the treatment carried out at this institution.

The method of treatment largely employed at these heart resorts is to withdraw all, or nearly all, of the active drugs that the patient may be taking, and to substitute physical and physiologic methods of therapy. These include bathing, regulation of the diet, and exercise. This exercise consists of two varieties: exercise of the muscles against the resistance of an attendant, and exercise by walking on inclined planes or up hills. The treatment is aimed at chronic heart disease, to develop a greater cardiac reserve strength; the whole object of the treatment is to strengthen the myocardium, either in conditions of its debility or in conditions of diminished compensation in valvular disease. Any treatment that will develop a reserve heart strength to be called on in emergencies, more or less similar to the reserve strength of a normal heart, tends to prolong the patient's life and health.

Patients with acute heart failure or acute loss of compensation, with more or less serious edemas, should rarely take the risk of traveling any distance to be treated at an institution. As a general rule they are better treated for a few weeks or months at home. After the broken compensation is repaired, a reserve strength of the heart may well be developed by a visit to one of these institutions, if the patient can afford it.

The Oertel treatment consists chiefly in diminishing the fluids taken into the body, and in graduated mountain-climbing. By diminishing the fluids taken, the work of the heart is diminished, as the blood-vessels are not overfilled and may even be underfilled. The diet is carefully regulated with the object of removing all superfluous fat from the body. The third leg of the tripod of the Oertel treatment is the gradually increasing hill and mountain-climbing to educate the heart by graded muscular training to become strong, perfectly compensatory, and later to develop a reserve strength. This particular cure is especially adapted to the obese, who have weakened heart muscles.

NAUHEIM BATHS

At Nauheim, under the direction of Dr. Theodore Schott, baths form an important part of the treatment. These baths are of two kinds, the saline and the carbonic acid. The medicinal constituents of the saline bath are sodium chlorid and calcium chlorid, the strength of each varying from 2 to 3 per cent. The baths at first are given at a temperature of 95 F., and as the patient becomes used to them and can take them without discomfort, the temperature is gradually reduced. The patient remains in the bath from five to ten minutes. After the bath he is dried with towels and rubbed until the cutaneous circulation becomes active. He must then lie down for an hour. These baths are repeated for two or three days, and are omitted on the third and fourth days, to be resumed on the following day. After a few baths have been taken, the carbon dioxid baths are commenced, beginning with a small quantity of the gas, which is later gradually increased. This course of baths should be continued from four to eight weeks. Unless there is some special reason for taking them at some other period of the year, they are taken more advantageously during the warm months.

Besides the baths, an important part of the treatment at Nauheim consists in the exercises against resistance. These are usually given an hour or more after a bath, and are taken with great deliberation; their effect is carefully watched by an intelligent attendant so that no harm may be done by the exercise.

During this treatment the food is, of course, carefully regulated with the aim of giving a mixed, sufficient, easily digestible and easily assimilated diet. All highly seasoned dishes, all effervescent drinks and anything that tends to cause gas in the stomach and intestines are prohibited. Coffee and tea are not allowed, except the "dekofa," which is coffee without its caffein; and it may be noted that it has recently been shown that caffein is one of the surest of drugs to raise the blood-pressure, and is therefore generally not desirable when the heart muscle requires strengthening. Because of its tendency to raise blood-pressure and weaken cardiac muscle, tobacco is entirely forbidden at Nauheim, except in a few individual instances, and then the amount allowed is a minimum one. Large amounts of liquid are not allowed because they distend the stomach, raise the blood-pressure and increase the pumping work of the heart.

One of the greatest advantages of the treatment at an institution like Nauheim is the general hopeful spirit instilled into the patients, who are so many times seriously depressed by the knowledge of a heart weakness and the realization of their physical inability to do what other persons are able to do. Also, it is of great value to send a patient to a resort where the climate is good and the scenery is lovely and soothing. No disease, perhaps, needs cheerfulness and pleasantness and lack of anxiety or frets more than does cardiac weakness. A tuberculous patient may sit on a mountain-top with snow blowing about him, and recover; a heart patient must have sunshine and comfort.

The results of such sanatorium treatment of heart disease are often evident not only to the patient by an increase of general muscle strength, the ability to do ordinary things and perhaps even sustain muscular effort without dyspnea and cardiac discomfort, but also to the physician by the physical signs. The contraction of the heart becomes stronger and the normal sounds more decided; murmurs which were entirely due to dilated ventricles and insufficiency disappear, while the permanent murmurs may become louder from a more forceful, normal action of the heart muscle. The pulse becomes slower, and the blood-pressure, from being too low, becomes normal for the age of the individual. The heart will often also actually decrease in size, and the apex-beat become localized rather than diffuse. The liver becomes reduced in size; the urine is less concentrated, and if there were traces of albumin after exertion, these disappear.

It should perhaps be emphasized that not a little benefit from these resort treatments may be due to the withdrawal of unnecessary drugs. Many heart patients are overdrugged.

This sort of treatment is contra-indicated in some kinds of heart disease, as heart weakness due to arteriosclerosis with high blood-pressure, to aneurysm of the thoracic or abdominal aorta, and to nephritis.

So many heart patients have been improved by the Nauheim treatment that the question arises as to whether the treatment can be conducted at home or in a sanatorium near home, when the patient is unable to go to this resort; that is to say, Can we establish this treatment for the majority of patients who have chronic heart disease? Of course, even at home, the sodium chlorid and calcium chlorid baths may be given, and one may obtain the salts with which to make the carbon dioxid bath; the exercises may be given, and walking on various ascending grades may be inaugurated. All patients will be more or less benefited, provided they will carry out the treatment. Unfortunately, the surroundings at a patient's home are generally adverse to perpetuating these treatments long enough to develop the muscular strength of the heart to the reserve desired. If a patient appears pretty well, especially if he is stimulated by his family to believe that he is well, he thinks the continuation of the treatment entirely unnecessary, and unless he goes to a resort where he sees other patients with similar conditions able to do what he is not able to do, and therefore is stimulated to acquire their ability by the treatment outlined, he will not follow his physician's directions. There are several sanatoriums in this country where the diet, hydrotherapy and exercise necessary for developing heart strength are carried out, and patients are sent to some of them with great advantage.

It has been found that these stimulant baths do not act well in mitral stenosis, if the left ventricle is small. If the left ventricle is unable to receive and therefore send out into the systemic circulation sufficient blood to dilate the peripheral capillaries under the irritation of the baths or the vasodilator effects of the baths, the bath treatment does harm instead of good. A patient who has mitral stenosis and also a small left ventricle will be found to be poorly developed, badly nourished, and to have poor peripheral circulation.

As elsewhere stated, the improvised carbon dioxid bath, to stimulate the skin so as to reduce the blood-pressure, is not satisfactory. Other methods of reducing blood-pressure, when it is too high, are much more effective.

HEART DISEASE IN CHILDREN AND DURING PREGNANCY

A common characteristic in a large proportion of middle-aged or old patients with heart disease is the presence of degenerative changes in the myocardium, the valves, or the arteries of the heart. In children, on the other hand, the most common disturbances of the heart are acute inflammations affecting its different structures, and due in most instances to acute infections. Myocarditis and endocarditis occur frequently, and pericarditis occasionally. As in adults, rheumatism is the most common cause of inflammation of the structures of the heart, but rheumatism causes inflammation of the heart much more frequently in children than in adults. Besides this infection, the most frequent causes of inflammation of the heart in children are diphtheria, scarlet fever, typhoid fever, measles and influenza, with the frequency, perhaps, in the order named. Diphtheria frequently gives rise to myocarditis, which results in dilatation of the heart. This may occur in the second or third week of the course of the disease, and even up to the eighth and tenth week from the beginning of the disease. The myocarditis due to diphtheria is not always the cause of sudden death occurring during the disease, as such a fatal result may be due to paralysis of nervous origin. In scarlet fever inflammation of the heart may be due directly to the poison of the disease, or it may be secondary to a nephritis which is so frequent a complication of scarlet fever. It is probable that the inflammation of the skin in scarlet fever, preventing normal secretion, may be a cause of a sometimes increased blood-pressure and also of the nephritis, both of which conditions may predispose to the cardiac complication. Erysipelas may cause acute inflammation of the heart, perhaps for the same reasons.

A certain proportion of cardiac diseases in children, especially endocarditis, seems to be due to a general septic infection which results in the so-called septic, infectious or malignant endocarditis. There is sometimes a tendency in certain children, and perhaps in certain families, for the heart to become readily infected during an infectious disease, more than in other children who suffer from the same disease. Sometimes the heart becomes inflamed in rheumatic children without any joint affection occurring; the inflammation in the heart may be the only manifestation of the disease.

This etiology of cardiac affections of children indicates the directions in which therapeutic efforts should be aimed. In children who are under the more or less constant care of the family physician, the possibility of the occurrence of some cardiac affection should be borne in mind, especially in children in families who are known to be affected with what may be called a rheumatic diathesis — families in which several members have suffered from rheumatism. It is reasonable to suppose that children who are delicate and feeble, who do not have sufficient fresh air, who do not take sufficient exercise, and who are not properly fed are more likely to be affected with cardiac complications in the presence of infectious diseases than children who have had plenty of fresh air, an abundance of exercise and a sufficient amount of proper food.

At the present day it is hardly necessary to insist on the importance of giving every child an adequate amount of fresh air. It is possible, however, that this gospel has been overworked, and it is not infrequently necessary to caution some parents that there is danger of impairing their children's health by too much exposure. The old ideas of the influence of exposure to cold and dampness in the production of rheumatism have not yet been so far abandoned that we can entirely neglect the possibility of rheumatism being developed, at least, by the exposure to cold winds and dampness of children who are otherwise predisposed to this disease. It is possible that the enormously increasing number of children with adenoids and enlarged tonsils, who need operative measures for their removal, may have these conditions aggravated by too much exposure to the inclemency of variable, harsh weather.

It is not necessary to state that proper exercise develops the heart, as it does all the other muscles; but at the same time it is necessary to caution parents against allowing their children to indulge in too violent and too prolonged exercise. Young children probably stop often enough in their play not to overwork their hearts. Older boys and girls, especially boys, are inclined to take too severe athletics, such as long-distance running, competitive rowing, violent football and rapid cycling. It should be emphasized to schoolmasters, gymnasium teachers and athletic trainers that a boy who is larger than he should be at his age has not the circulatory ability that the older boy of the same size has. The overgrown boy has all he can do to carry his bulk around at the speed of his age and youth. The addition of competitive labor overreaches his reserve heart power, and he readily acquires a strained, injured heart. On the other hand, moderate indulgence in walking, baseball, swimming, rowing and golf should be commended. It is not exactly the exercise that does him the harm, it is the competitive element in it. Until a boy is well developed in his internal reserve strength, he should not compete with other boys who are better developed. His pride makes him do himself injury.

Dietetic fads are so prevalent to-day that there is danger that many children will not receive an adequate amount of nutriment, that they will be fed an excess of such foods as are likely to produce damage to their constitutions, or that they will be given food that does not contain all the different elements of nutrition to satisfy their economy and their growth. While it is now generally acknowledged that an excess of meat is not beneficial to any one, on the other hand a moderate amount is necessary for individuals who are working or are mentally active, especially for growing children. Also a too great limitation of the child's diet to farinaceous foods, and especially the allowance of too much sugar and sugar-producing food, is likely to encourage the development of rheumatism. A mixed diet, not excessive in amount, and prepared so that it will be digested without difficulty is most useful, and it should include in suitable proportions meat, milk, eggs, vegetables, starches and fruit. These should all be taken at regular intervals, thoroughly chewed, and should not be taken in excess.

If a child has had an attack of heart inflammation, a myocarditis or an endocarditis, greater care should be taken of him not only when he is well but especially when he becomes ill of any other disease. If the child has had a rheumatic inflammation of the heart, or has had rheumatism without such a complication, it is considered by some clinicians wise to give a week's treatment with salicylates at intervals of three or four months, for two or three years, perhaps. It is hard to determine how much value this prophylactic treatment has. If the child's surroundings cannot be changed and he is subjected to the same conditions of possible reinfection, it may be a wise precaution, much like the prophylactic administration of quinin in malarial regions. If a child has developed a cardiac inflammation during any disease, the treatment is that previously outlined.

An important part of prophylaxis and treatment of a cardiac affection during the course of any disease is the prevention of serious anemia. During sickness the patient is likely to become more or less anemic, but the administration of iron, in the manner previously suggested, during the course of the disease, and especially during rheumatism, will prevent the anemia becoming rapid or severe.

CARDIAC DISEASE IN PREGNANCY

It is so serious a thing for a woman with a valvular lesion or other cardiac defect to become pregnant that no young woman with heart disease should be allowed to marry. Perhaps every normal heart during pregnancy hypertrophies somewhat to do the extra work thrown on it, but it may easily become weakened and show serious disturbance as its work grows harder and the distention of the abdomen and the upward pressure on the diaphragm increase. This pressure perhaps generally displaces the apex of the heart to the left and causes the heart to lie a little more horizontal. If the patient is normal, there may be a gradually increasing blood-pressure all through the months of pregnancy, and if the kidneys are at all disturbed this pressure is increased, and there is, of course, much increased resistance to the circulation during labor. The better the heart acts, the less likely are edemas of the legs during pregnancy. It is thus readily seen that pregnancy is a serious thing for a damaged heart. The reserve strength of the heart muscle, as has been previously stated, is much less in valvular compensation than that of a normal heart, and this reserve force is easily overcome by the pregnancy, and loss of compensation occurs with all of its usual symptoms.

The most serious lesion a woman may have, as far as pregnancy is concerned, is mitral stenosis. An increased abdominal pressure interferes with her lung capacity, and her lungs are already overcongested. The left ventricle may be small with mitral stenosis, and therefore her general systemic circulation poor. For these two reasons mitral stenosis should absolutely prohibit pregnancy. While many women with well-compensated valvular disease go through a pregnancy without serious trouble, still, as above stated, they should be advised never to marry. If they do marry, or if the lesion develops after marriage, warning should be given of the seriousness of pregnancies.

If a woman becomes pregnant while there are symptoms or signs of broken compensation, there can be no question, medically or morally, of the advisability of evacuating the uterus. The same ruling is true if during pregnancy the heart fails, compensation is broken, and the usual symptoms of such heart weakness develop, provided that a period of rest in bed, with proper treatment, has shown that the heart will not again compensate. Under such a condition delay should not be too long, as the heart may become permanently disabled. If during pregnancy in a patient with a damaged heart albuminuria develops and the blood-pressure is increased, showing kidney insufficiency, there can be no question of delay from every point of view, labor must be precipitated; the uterus must be emptied to save the mother's life.

If a pregnant woman is known to have a degenerative condition of the myocardium, or arteriosclerosis, the danger from the pregnancy is serious, and the pregnancy should rarely be allowed to continue.

Even if no serious symptoms occur during the term of the pregnancy and the heart continues to compensate sufficiently for its defect, labor should never be allowed to be prolonged. The tension thrown on the heart during labor is always severe and has not infrequently caused acute heart failure by causing acute dilatation, and in these damaged hearts tediousness and severe, intense exertion should not be allowed. Proper anesthetics and proper instrumentation should be inaugurated early.

Patients who have successfully passed through the danger of pregnancy with cardiac lesions, possibly relieved by radical treatments, should be warned against ever again becoming pregnant. If this warning does not prevent future pregnancies, the family physician and his consultant must decide just what is proper to do. It is to be understood that no uterus should ever be emptied until one or more consultants have approved of such treatment.

Sometimes serious heart weakness develops during the later weeks of pregnancy, requiring the patient to remain in bed and receive every advantage which rest, proper care and well-judged medicinal treatment will give the circulation.

If the heart is weak and there have been signs of myocardial weakness or loss of compensation, the sudden loss of abdominal pressure after delivery may allow the blood-vessels of the abdomen to become so overfilled as to cause serious cerebral anemia and cardiac paralysis. Therefore in such cases a tight bandage must immediately be applied, and it has even been suggested that a weight, as a bag of sand weighing several pounds, be placed temporarily on the abdomen. The greatest possible care should be given these women during and after labor.

Acute dilatation is not an infrequent cause of death during ordinary labor, and is more likely to occur in these cardiac patients. If signs of acute dilatation of the heart occur, with associated pulmonary edema, venesection (especially if there has not been much uterine hemorrhage), with the coincident intramuscular injection of one or two syringefuls of aseptic ergot, will often be found to be life-saving treatment.

Septic infections after parturition are prone to cause endocarditis and myocarditis, and a malignant endocarditis may develop from uterine infection or uterine putridity.

DEGENERATIONS

CORONARY SCLEROSIS

While disease of the coronary arteries may occur without general arteriosclerosis, it is so frequently associated with it that it is necessary to give a brief description of the general disease. Arteriosclerosis or arteriocapillary fibrosis is really a physiologic process naturally accompanying old age, of which it is a part, or the cause, and it should be considered a pathologic condition only when it occurs prematurely. It may, however, occur at almost any age after 30, and is beginning to be frequent between 40 and 50. In rare instances it may occur between 20 and 30, and even in childhood and youth. It is much more frequent in men than in women. Its most common cause is hypertension; in fact, hypertension generally precedes it. The most frequent cause of hypertension to-day is the strenuousness of life, the next most frequent cause being the toxins circulating in the blood from overeating, overdrinking, overuse of tobacco and the overuse of caffein in the form of coffee, tea or caffein drinks. Another common cause of arteriosclerosis occurring too early is the occurrence of some serious infection in an individual, typhoid fever and sepsis being most frequent. Syphilis is a frequent cause, especially of that form of arteriosclerosis that shows the greatest amount of disease in the aorta. Mercury used in the treatment of syphilis is more likely, however, than syphilis to be the cause of arteriosclerosis. Although this drug, even with the arsenic injections now in vogue, is necessary for the cure of syphilis, it probably tends to raise the blood-pressure by irritating the kidneys and by diminishing the thyroid secretion, both of these occurrences predisposing to arteriosclerosis. From the fact that lead-poisoning causes an increased blood-pressure, lead is a probable cause of arteriosclerosis. With the greater knowledge of the danger of poisoning possessed by those who work in lead, chronic lead-poisoning is becoming rare, as evidenced by the lessening frequency of wrist-drop and lead colic.

Chronic nephritis is often a coincident disease, but the causes of the arteriosclerosis and the nephritis are generally the same. Alcohol, except as a part of overeating and as a disturber of the digestion, is perhaps not a direct cause of arteriosclerosis, as alcohol is a vasodilator. Hard physical labor and severe athletic work may cause arteriosclerosis to develop, and it is likely to develop in the arteries of the parts most used.

Hypertension is generally a prelude to arteriosclerosis, and everything that tends to increase tension promotes the disease; everything that tends to diminish tension more or less inhibits the disease. Therefore a subsecretion of the thyroid predisposes to arteriosclerosis, and increased secretion of the adrenals predisposes to arteriosclerosis, the thyroid furnishing vasodilator substance and the adrenals vasopressor substance to the blood. Also, if these secretions are abnormal, protein metabolism is more or less disturbed.

While arteriosclerosis often occurs coincidently with gout, and gout apparently may be a cause of arteriosclerosis, still the two diseases are widely dissociated, and the causes are not the same.

Although the arterial pressure has been high before arteriosclerosis developed and may remain high for some time in the arteries, unless the heart fails, the distal peripheral pressure, as in the fingers and toes, may be poor in spite of the high blood-pressure. When the left heart begins to fail, pendent edema readily occurs.

PATHOLOGY

The pathology of arteriosclerosis is a thickening and diminishing elasticity of the arteries, beginning with the inner coat and gradually spreading and involving all the coats, the larger arteries often developing calcareous deposits or thickened cartilaginous plates — an atheroma. If the thickening of the walls of the smaller vessels advances their caliber is diminished, and there may even be complete obstruction (*endarteritis obliterans*). On the other hand, some arteries, especially if the calcareous deposits are considerable, may become weakened in spots and dilatation may occur, causing either smaller or larger aneurysms.

Histologically the disease is a connective tissue formation beginning first as a round-cell infiltration in the subendothelial layer of the intima. This process does not advance homogeneously; one side of an artery may be more affected than the other, and the lumen may be narrowed at one side and not at the other, allowing the artery to expand irregularly from the force of the heartbeat. As the disease continues the internal elastic layer is lost, the muscular coat begins to atrophy, and then small calcareous granules may begin to be deposited, which may form into plates. In the large arteries the advance of the process differs somewhat. There may be more actual inflammatory signs, fatty degeneration may occur, and even a necrosis may take place.

However generally distributed arteriosclerosis is, in some regions the disease is more advanced than in others, and in those regions the most serious symptoms will occur. The regions that can stand the disease least well are the brain and coronary arteries, and next perhaps the legs, at the distal parts at least, where the circulation is always at a disadvantage if the patient is up and about.

SYMPTOMS

The symptoms are increased tension, which means, sooner or later, hypertrophy of the left ventricle and an accentuated closure of the aortic valve. This alone means more and more tendency to aortic irritation and aortic valve irritation, with inflammation, and later deposits of calcareous material perhaps with stiffening of the aortic valve and narrowing, aortic stenosis being the result. If such a patient with the disease advanced to this stage must overwork, or sustains any severe muscle strain, an aneurysm of the aorta may occur. In the meantime, with the advancing degeneration of the cerebral arteries, some sudden cerebral congestion, caused by leaning over, lifting, vomiting or hard coughing, may rupture a cerebral vessel, and all the symptoms of apoplexy are present. If small hemorrhages occur in the arterioles of the extremities, of course the prognosis is not serious. Sometimes some of the smaller vessels of the brain may become obstructed and cerebral degeneration occur. If distal vessels become obstructed, as of the toes or feet, gangrene takes place unless the obstruction occurs at a place where the collateral circulation could save the part from such a death. These are some of the ultimate results of serious and final arteriosclerosis. The more frequent result, when the disease has not advanced so far, is a failing heart, either from degenerative myocarditis, coronary sclerosis or dilatation, with all the symptoms of coronary sclerosis and angina pectoris, or with the symptoms of failing circulation.

With high blood-pressure to the point of beginning endarteritis a gradually increasing force of the apex-beat occurs, the aortic closure is accentuated as just described, the pulse is slow, the tensity of the arteries depends on the stage of the disease, and when the disease is actually present the palpable arteries do not collapse on pressure. They soon lose their elasticity, and if this occurs in parts that are soft and flexible, the arteries become more or less tortuous by the force of the blood-current twisting and bending them, owing to the irregularity of their hardening. The extremities readily become numb, or the part "goes to sleep," as it is termed. This occurs frequently at night. Sooner or later some edema of the feet and legs occurs in the latter part of the day. Sometimes abdominal colic attacks occur, caused by disturbed circulation. Various disturbances of metabolism may occur, depending on the circulation in the different organs or on coincident disease, and the liver, pancreas and kidneys may be affected.

The blood-pressure, if taken in the arms especially, may appear excessively high, but really the actual pressure in the blood-vessels may be low. This is on account of the inability to compress the hardened arteries. A heart may be weak and actually need strengthening even while the blood-pressure reading is high.

The treatment of this disease is successful only in its prevention, and the treatment of hypertension before arteriosclerosis is present will be described later. When the disease is actually present, there is nothing to do except for the patient to stop active labor, never to overeat or overdrink, to prevent, if possible, toxemias from the bowels, to keep the colon as clean as possible, and for the physician to give the heart such medicinal aids as seem needed, vasodilators if the heart is acting too strongly, possibly small doses of cardiac tonics if the heart is acting weakly; always with the knowledge that a degenerative myocarditis may be present in considerable amount, or that coronary sclerosis may be present.

As above stated, coronary sclerosis probably seldom occurs without more general arteriosclerosis. Obstruction of the coronary arteries, however, not infrequently occurs at their orifices in conjunction with sclerosis of that region of the aorta and of the aortic valve. The more these arteries are diseased and the more they are obstructed, the more the myocardium of the heart becomes degenerated, softened and weakened, when dilatation of the ventricles, especially the left, is likely to occur. Sooner or later such a condition will cause attacks of angina pectoris and more or less pronounced symptoms of chronic myocarditis and fatty degeneration, as previously described.

TREATMENT

The treatment of a suspected coronary sclerosis is the same as that of general arteriosclerosis — primarily the treatment of anything that tends to cause high tension or to produce chronic endarteritis. When either general or local arteriosclerosis is present, the treatment that should be inaugurated comprises anything that would tend to inhibit the endarteritis and the calcification — necessary periods of rest, the interdiction of all physical effort or physical strain, and the regulation of the diet, digestion and elimination. Perhaps there is no greater preventive of the advance of this disease than a diet considerably less than would be suitable for the same individual when in perfect health and at his regular work. The amount of protein especially should be reduced, and the meal hours should be regular. Ordinarily all tea, coffee and tobacco should be forbidden, and alcohol should be allowed only to the aged, if allowed at all.

It has long been considered that iodin would inhibit abnormal connective-tissue growth. Iodin most readily reaches the blood as a sodium or potassium iodid. Large amounts of iodin are not needed to saturate the requirements of the system for iodin, 0.1 to 0.2 gm. (1½ to 3 grains) preferably of sodium iodid, twice a day, after meals, given with plenty of water, being sufficient; but it should be continued in one or two doses a day not only for weeks, but for months. Whether this iodid or iodin acts *per se*, or acts by stimulating the thyroid gland to increased activity and therefore to more normal activity, so that it is the thyroid secretion that is of benefit, it is difficult to decide. In view of the fact that in advanced years the thyroid is always subsecreting, and after the very diseases that cause arteriosclerosis or during the diseases that cause arteriosclerosis the thyroid is generally subsecreting, it would appear that the value of iodin is in its effect in stimulating the thyroid gland.

If a small amount of thyroid secretion is evidenced by other symptoms, thyroid extract should be given. The dose need not be large, and should be small, but should be given for a considerable length of time. If the patient seems to be improving on small doses of iodid, however, and the thyroid is supposed not to be very deficient, it is better not to administer thyroid extract. unless the patient is obese.

A serum treatment given intravenously, hypodermatically, by the mouth, and by the rectum was inaugurated some years ago (1901 and 1902), and is known as the "Trunecek serum." This first consisted of sodium sulphate, sodium chlorid, sodium phosphate, sodium bicarbonate and potassium sulphate in water in such amounts as to simulate the blood-plasma. Later small amounts of calcium and magnesium phosphate were added to the solution to be injected. These injections seemed to lower the blood-pressure, but it is doubtful whether they had any greater ability than a proper regulation of the diet to inhibit arteriosclerosis. At any rate, these injections are but seldom used.

An important means of inhibiting disturbance from any arteriosclerosis which should be employed when possible is the climate treatment. Warm, equable climates, in which there are no sudden radical changes, are advantageous when coronary sclerosis is suspected, and warm climates are valuable in promoting the peripheral circulation and lowering the blood-pressure in arteriosclerosis. These patients always require more heat than normal persons, always feel the cold severely, and their hearts always have much less disturbance, fewer irregularities and fewer attacks of pain during warm weather than during cold weather.

Simple hydrotherapeutic measures are also necessary for these patients, but baths should not be used to the point of causing debility and prostration. Applications of cold water in any form are generally inadvisable. Very hot baths are also inadvisable; but pleasantly warm baths, taken at such frequency as found to be of benefit to the individual, relax the peripheral circulation, relieve the tension of the internal vessels, lessen the work of the heart, and promote healthy secretion of the skin, the skin of arteriosclerotic patients often being dry. This dry skin is especially frequent if there is any kidney insufficiency, which so soon and so readily becomes a part of the arteriosclerotic process.

If the patient is old, small doses of alcohol may act physiologically for good. In these arteriosclerotic patients the activities of alcohol should be considered from the drug point of view, not from that of an intoxicating beverage. Other drugs will be considered in the discussion of hypertension.

If the heart actually fails, the treatment becomes that of chronic myocarditis and of dilatation.

Not infrequently in sclerosis of the arteries, especially of the coronary arteries, the blood-pressure is not high, but low, and the heart is insufficient. In such patients cardiac tonics may be considered, but they must be used with great care. Digitalis may be needed, but it should be tried in small doses. It often makes a heart with arteriosclerosis have severe anginal attacks. On the other hand, if the heart pangs or heart aches and the sluggish circulation are due to myocardial weakness without much actual degeneration, digitalis may be of marked benefit. The value of digitalis in doubtful instances will be evidenced by an improved circulation in the extremities, a feeling of general warmth instead of chilliness and cold, an increased output of urine, and less breathlessness on slight exertion.

ANGINA PECTORIS

This is a name applied to pain in the region of the heart caused by a disturbance in the heart itself. Heart pains and heart aches from various kinds of insufficiency of the heart, or heart weakness, are not exactly what is understood by angina pectoris. It is largely an occurrence in patients beyond the age of 30, and most frequently occurs after 50, although attacks between the ages of 40 and 50 are becoming more frequent. It is a disturbance of the heart that most frequently attacks men, probably more than three-fourths of all cases of this disease occurring in men; in a large majority of the cases the coronary arteries are diseased.

Various pains that are not true angina pectoris occur in the left side of the chest; these have been called pseudo-anginas. They will be referred to later. True angina pectoris probably does not occur without some serious organic disease of the heart, mostly coronary sclerosis, fatty degeneration of the heart muscle, adherent pericarditis and perhaps some nerve degenerations. Various explanations of the heart pang have been suggested, such as a spasm or cramp of the heart muscle, sudden interference with the heart's action, as adherent pericarditis, a sudden dilatation of the heart, an interference with the usual stimuli from auricle to ventricle and therefore a very irregular contraction, a sudden obstruction to the blood-flow through a coronary artery, or a sudden spasm from irritation associated with some of the intercostal or more external chest muscles causing besides the pang a sense of constriction. Perhaps any one of these conditions may be a cause of the heart pang, and no one be the only cause.

In a true angina, death is frequently instantaneous. In other instances, death occurs in a few minutes or a few hours; or the patient's life may be prolonged for days, with more or less constant chest pains and frequent anginal attacks. Here there is a gradual failing of the heart muscle, with circulatory insufficiency, until the final heart pang occurs.

Anginal attacks before the age of 40, presumed, from a possible narrowing of the aortic valve, to be due to coronary sclerosis, are frequently due to a long previous attack of syphilis. In these cases active treatment of the supposed cause should be inaugurated, including perhaps an injection of the arsenic specific, and certainly a course of mercury and iodid, with all the general measures for managing and treating general arteriosclerosis, as previously described.

SYMPTOMS

The pain of true angina pectoris generally starts in the region of the heart, radiates up around the left chest, into the shoulder, and often down the left arm. This is typical. It may not follow this course, however, but may be referred to the right chest, up into the neck, down toward the stomach, or toward the liver. The attack may be coincident with acute abdominal pain, almost simulating a gastric crisis of locomotor ataxia. There may also be coincident pains down the legs. It has been shown, as mentioned in another part of this series of articles, that disturbances in different parts of the heart or different parts of the aorta may cause pain and the pain be referred to different regions, depending on the part affected.

Instances occasionally occur in which a patient has an anginal attack, as denoted by facial anxiety, paleness, holding of the breath, and a slow weak pulse, without real pain. This has been called angina *sine dolore*. The patient has an appearance of anxious expectation, as though he feared something terrible was about to happen.

The position of the patient with true angina pectoris is characteristic. He stops still wherever he is, stands perfectly erect or bends his body backward, raises his chin, supports himself with one hand, leans against anything that is near him, and places his other hand over his heart, although he exercises very little pressure with this hand. The position assumed is that which will give the left chest the greatest unhampered expansion, as though he would relieve all pressure on the heart.

Besides the feeling of constriction, even to some spasm, perhaps, of the intercostal muscles, respiration is slowed or very shallow, because of the reflex desire of the patient not to add to the pain by breathing. The face is pale, the eyes show fear, and the whole expression is almost typical of cardiac anxiety. The patient feels that he is about to die. The pulse is generally slowed, may be irregular, and may not be felt at the wrist. The blood-pressure has been found at times to be increased. It could of course be taken only in those cases in which there were more or less continued anginal pains; the true typical acute angina pectoris attack is over, or the patient is dead, before any blood-pressure determination could be made. When there is more or less constant ache or frequent slight attacks of pain the blood-pressure may be raised by the causative disease, arteriosclerosis. During the acute attack with inefficient cardiac action and a diminished force and frequency of the beat the peripheral blood-pressure can only be lowered.

The duration of an acute attack, that is, the acute pain, is generally but a few seconds, sometimes a few minutes, and rarely has lasted for several hours. In the latter cases some obstruction to an artery has been found on autopsy, but not sufficient to stop the circulation at a vital point. Repeated slight attacks, more or less severe, may occur frequently throughout one or more days, or even perhaps a series of days, caused by the least exertion, even that of turning in bed.

While most cases of sudden death with cardiac pain are due to a local disease in or around the heart, it is quite probable that some disturbance in the medulla oblongata may cause acute inhibitory stoppage of the heart through the pneumogastric (vagi) nerves. The power of the pneumogastric reflex to inhibit the action of the heart is, of course, easily demonstrated physiologically. Clinically reflexes down these nerves interfering with the heart's action cause faintness and serious prostration, if not actual shock, and perhaps, at times, death. The most frequent cause of such a reflex is abdominal pain, perhaps due to some serious condition in the stomach, to gastralgia, to an intestinal twist, to intussusception or other obstruction, or to hepatic or renal colic. A severe nerve injury anywhere may cause such a heart reflex. Hence serious nerve pain must always be stopped almost immediately, else cardiac and vasomotor shock will occur. In serious pain morphin becomes a life-saver.

MANAGEMENT

While a number of causes of true cardiac pain may be eliminated by improvement in any loss of compensation, by improvement of the heart tone, by more or less recovery from myocardial or endocardial inflammation, and by the withdrawal of nicotin, which may cause cardiac pains, still, a true angina pectoris once occurring is likely to be caused by a progressive, incurable condition and the attacks will become more frequent until the final one. It is possible that a true angina may be due to a coronary artery disease or obstruction and that a collateral circulation may become established and repair the deficiency. While this probably can take place, occurrences must be rare.

Occasionally when the intense pain has ceased the patient may be nauseated and actually vomit, or he may soon pass a large amount of urine of low specific gravity, or have a copious movement of the bowels.

The first attack, and subsequent ones more and more readily, are precipitated by any exertion that increases the work of the heart, as walking up hill, walking against the wind, going up stairs, physical strains as suddenly getting out of bed, leaning over to put on the shoes, straining at stool, or even mental excitement. Exertion directly after eating a large meal is especially likely to precipitate an attack. Food that does not readily digest or food that causes gastric flatulence may precipitate attacks. Any indiscretion in the use of coffee, tea, alcohol or tobacco may be the cause of an attack.

For treatment of the immediate pain, if the physician arrives soon enough, anything may be given that quickly relieves local or general arterial spasm and spasm of the muscles. The moment that the heart and its arterioles relax, the attack is often over. The most quickly acting drug for this purpose is amyl nitrite, inhaled. If amyl nitrite is not at hand, or has been found previously to cause considerable disturbance of the head or a feeling of prolonged faintness, nitroglycerin is the next most rapidly acting drug. It may be given hypodermatically, or a tablet may be dissolved on the tongue. The amyl nitrite should be in the emergency case of the physician in the form of ampules, or may be carried by the patient after he has had one or more attacks. The ampules now come made of very thin glass with an absorbent and silk covering ready for crushing with the fingers, and are thus immediately ready for inhalation. One of these is generally all that it is necessary to use at any one time. Nitroglycerin, if given hypodermatically, should be in dose of 1/100 grain. If given by mouth the dose should be the same, repeated in ten minutes if the pain has not stopped.

Almost coincidently with the administration of nitroglycerin or the amyl nitrite a hypodermatic injection of ⅛ or 1/6 grain of morphin sulphate should be given without atropin, as full relaxation is desired without any stimulation of atropin.

Alcohol is also a valuable treatment of this pain, when the drugs mentioned are not at hand. The dose should be large; whisky or brandy are best, and should be administered in hot or at least warm water. The physiologic action of alcohol, which dulls or benumbs the nervous system and dilates the peripheral bloodvessels, is exactly in line with the clinical indications.

If a patient is home and at rest at the time of an attack, a hot-water bag but slightly filled, or a pad electrically heated may be placed over the heart sometimes with marked advantage and relief from pain. Occasionally even such gentle applications are not tolerated.

After the attack is over absolute rest for some hours, at least, is positively necessary. If the attack was severe, the patient should rest several days, as there seems to be a great tendency for such attacks to come in groups, the cause being acutely present for at least some time. But little food should be given; nothing very hot or very cold, and no large amount of liquids; gentle catharsis may be induced on the following day, if deemed advisable; no stimulating drugs should be administered, and nothing that would raise the blood-pressure.

The question often arises as to whether the patient shall be told of the seriousness of his condition. It is hardly wise to withhold this knowledge from him, and generally is not necessary. The ordinary alert patient knows how serious the condition is by his own feelings, and will even reprove or joke with his physician for minimizing the danger. It is best that the whole subject be discussed carefully with him and his life regulated and ordered, and emergency drugs prepared and given him with proper instructions to the family, so that he may possibly prevent other attacks, and, if they occur, may have the best immediate treatment.

The acute symptoms being over, a careful analysis of the probable cause of the anginal attack should be made. If it is a general sclerosis, the treatment should be directed to that condition. If it is a myocarditis, a fatty degeneration of the heart or a fatty heart, this should be properly treated as previously described. If it is due to a toxemia from intestinal disturbance, that may readily be remedied. If due to nicotin, it need not again occur from that reason, and perhaps the damage caused by the nicotin may be removed. Any organic kidney trouble must, of course, be managed according to its seriousness, and if there is hypertension without an-serious lesion, the treatment should be directed toward its relief. The treatment for hypertension will be described later.

PSEUDO-ANGINA

While this name is more or less unfortunate, it has long been in vogue as a designation for pains and disturbances referred by a patient to his heart. Therefore with the distinct understanding that if the diagnosis is correct the name is a misnomer, it may be allowable to discuss under this heading some of the attacks that may simulate an angina and must be separated from a true angina.

To decide whether pain in the region of the heart or irregularity of its action is due to organic disease, to functional disturbance, or to referred causes is often extremely difficult. Some of the most disturbing sensations in the region of the heart are not due to any organic trouble, and yet the patient is fearful that such sensations mean some kind of heart disease, and therefore becomes exceedingly anxious and watches and mentally records every sensation in the left chest. This is unfortunate, as the patient may learn to note, if he does not actually count, his heart-beats, while normally he should sense nothing of his heart's activity. On the other hand, as just stated, it may be almost impossible to decide that this disturbance of the heart is not due to an organic cause, but is entirely functional, or due to some extraneous reason.

It seems justifiable in every case of irregular heart action to assure the patient that the condition can be improved, which in most instances is the truth. There can be no question of such urgent assurance, if it is decided that the cause is not in the heart itself, or at least is not organic. Irregularities in the heart's action will be discussed later. At this time discussion will be limited to pain which is not true angina pectoris, but which is in the region of the heart or is referred to it.

Intercostal neuralgia is more likely to occur on the left side of the chest than on the right. This is particularly unfortunate, as tending to cause these pains to be referred to the heart. The localization of tender spots along the course of a nerve with demonstration of these to the patient and the diagnosis stated is all the assurance that he requires.

Careful questioning, and if necessary scientific examination of the stomach, may show that the patient has hyperchlorhydria, ulcer of the stomach or duodenum, dilatation of the stomach, or some growth in the stomach as a cause for the pain referred to the region of the heart. Gall-stones in the gall-bladder may also give such referred pains. Other lesions in the abdomen may cause pain referred to the cardiac region. Not only will the demonstration of these causes and their treatment assure the patient that he has not neuralgia of his heart, but also, if curable, the cause of the pain may be removed.

Dry pleurisy of the left chest is not an infrequent cause of these pains, and of course serious disease of the lungs, as tuberculosis, unresolved pneumonia, pleuritic adhesions, emphysema and tumor growths may all be the cause of a referred cardiac pain, the heart being disturbed secondarily.

A stomach cramp is a not infrequent cause of serious pain referred to the heart, and the rare condition of cardiospasm must also be remembered as a cause of pseudo-angina. In other words, the interpretation of these pseudo-anginas means a careful diagnosis of the condition, and as previously stated, not only must the above-named causes be excluded, but also the reverse must be remembered: that many disturbances treated as other conditions really are due to cardiac weakness.

The diagnosis of a real angina pectoris from a false angina may not be difficult. A real angina generally occurs after exertion of some kind, be that exertion ever so slight. False angina may occur at any minute with or without exertion. Pain referred to the heart which awakens a patient at night is not likely to be a true angina; nervous patients are prone to have such night attacks of cardiac disturbance of various kinds. A true angina causes the patient's face to look anxious and pale, with the breathing repressed. A false angina shows no such paleness, allows deep breathing, crying and lamenting, and allows the patient to move about in bed, or about the room. The true angina makes the patient absolutely still and quiet: he hardly dares to speak or tell what he is feeling and fearing. True angina is of course much more frequent in older persons, while false anginas occur in the young, and especially in the neurotic. With all the other manifestations of hysteria, palpitation and cardiac pain are often symptoms.

It should not be decided, however apparently self-evident, that a referred pain is not due to cardiac lesion until a careful examination of the patient has been made. Real cardiac disturbance can of course occur at any time in a neurotic or hysteric patient, and there should be no mistakes of omission from carelessness or neglect on the part of the physician.

Other frequent causes of more or less disturbance of the heart's action, often accompanied by pain, are over-exertion, worry and mental anxiety, and intestinal toxemias due to too much protein or disturbed protein digestion. Frequent causes are tobacco, and the overuse of tea and coffee. Many a patient's pseudo-anginas are corrected by stopping tea and coffee. The effects of caffein and tobacco on the heart will be considered later when toxic disturbances are under discussion.

The above-mentioned causes of pseudo-anginas have only to be named to indicate the treatment that will prevent the pain attacks. At times, the cause being intangible, it may be necessary to change the whole life and metabolism of the patient, as so often necessary in hysteria, neurasthenia, gout, intestinal fermentation and kidney inefficiency. Besides a rearrangement of the diet and measures for causing proper activity of the bowels, massage, exercise and hydrotherapy should be utilized toward the end of improving the nutrition of every part.

TREATMENT OF PSEUDO-ANGINAS

The treatment of these pseudo-anginas depends, of course, on the diagnosis of the cause, and the cause should be eliminated or modified. If the heart shows real disturbance from this reflex cause, the treatment aimed toward it depends on whether the heart action is weak or strong and the circulation poor or good. If the circulation is poor, digitalis in small doses may be needed, either 5 drops of an active tincture twice a day, or 8 or 10 drops once a day. If digitalis is not indicated, strophanthus sometimes is valuable. While strophanthus has been shown not to be a real cardiac tonic like digitalis, still there seems to be a nervous sedative action when it is given by the mouth, and it often does good in these cases. The dose is 5 drops of the tincture, in water, three times a day, after meals. Strychnin in small doses may be needed, but in these patients, who are generally nervous, it is usually better not to give it.

One of the best sedatives to a heart that is irregular in its action and not acting strongly is lime; a good way to administer it is in the form of calcium lactate and the dose is 0.3 gm. (5 grains), in powder or capsule, three times a day, after meals.

If the circulation is good and the heart is strong, and yet these irregular pains and irregular contractions occur, the bromids act favorably and successfully. This is probably on account of their ability to quiet the central nervous system, to quiet and soothe the irritability of the heart, and to relax the peripheral blood-vessels. The dose should be from 0.5 to 1 gm. (7½ to 15 grains), in water, three times a day, after meals. It is not necessary or advisable to continue the bromid very long. Whatever general tonic or eliminative treatment the patient requires should be given. The value of hydrotherapy, massage and graded exercise should not be forgotten.

STOKES-ADAMS DISEASE: HEART-BLOCK

Stokes-Adams disease, or the Stokes-Adams syndrome, is a name applied to a combination of symptoms which was described by Stokes in 1846, and had been observed by Adams in 1827. The disease is characterized by bradycardia and cerebral attacks, either syncope or pseudo-apoplectic or convulsive attacks.

To understand the phenomena of this disease, it will be well to refer to the first article of this series on diseases of the heart. Until 1893, when His described the bundle of muscle fibers which is now known by his name, the transmission of the cardiac stimulus to contraction was not understood. It has been found, by studying the pathology of Stokes-Adams disease, as well as by clinically noting with instruments the contractions of different parts of the heart, that these slow heartbeats are really due to interruptions of the impulse passing from auricle to ventricle through the bundle of His, and degeneration in this region is generally the cause of Stokes-Adams disease. The auricles often beat many times more frequently than the ventricles, even two or three times as frequently, and, of course, these auricular contractions are not transmitted to the arterial system, and the radial pulse notes only the contractions of the ventricles. The phrase that is used to describe this non-transmission of the auricular stimulus to the ventricles is "heart-block."

While this disease almost invariably has a pathology, cases have occurred in which no lesion of the heart could be found, but it generally occurs coincidently with arteriosclerosis, in which the coronary arteries are more or less involved and the arterial system of the brain may be diseased. It occurs more frequently in men than in women, and in them mostly after middle, or in advanced, life. The previous history of the patient has often disclosed syphilis.

The intermittency of the pulse may be regular or irregular, and may not be constant in the early stages of the disease, but when the disease is established the rate of the pulse may be reduced to forty, thirty, or even twenty beats a minute, and it has been known to be even less. When these intermittencies are regular, perhaps two beats to one intermittency or three beats to one intermittency are the most frequent types. When the auricles also beat slowly it is perhaps the pneumogastric or vagi nerves that are for some reason overstimulated and thus inhibit the heart's activity.

The attacks of syncope are doubtless due to anemia of the medulla, because of the infrequent ventricular contractions. This anemia of the medulla and of the brain may also cause an epileptic seizure, or a partial paralytic seizure without any apparent paralysis. It is probable, however, that in these cases there may be coincident arterial disease in the brain. These sudden syncopal attacks are likely to occur when a patient suddenly rises from a reclining posture, especially if he has been asleep. Many persons whose circulation is none too strong may feel faint on suddenly rising, but in a person whose pulse is slow and the circulation weak the danger of causing anemia of the brain by the sudden erect posture is much increased. Slight faint turns are of frequent occurrence with these patients; or the faintness may be so rapid and so intense that the patient may drop in his tracks. Venous pulsation in the neck is generally marked with these patients, showing an impeded contraction of the right auricle.

If the auricles are heard or found by instrumental readings to contract more frequently than the ventricles, the trouble is quite likely to be a heart-block from disease in the heart itself, in the bundle of His. If the heart is slowed as a whole, the trouble might be due to diseased arteries or pressure from a growth, a gumma, perhaps, or other brain tumor in the region of the pons Varolii or medulla oblongata; or a hemorrhage into the fourth ventricle, causing pressure, could be the cause.

TREATMENT

The treatment of Stokes-Adams disease is unsuccessful. If general arteriosclerosis is present, that condition should be treated. Digitalis would seem almost invariably contra-indicated, although it is of value in extrasystoles without heart-block, or in conditions which are not Stokes-Adams disease; but if this disease was considered present, digitalis would probably do harm. Sometimes strychnin is of benefit.

Atropin has sometimes caused stimulation of the heart to more normal rapidity. Its benefit is generally only temporary, as most patients cannot take atropin regularly without having it cause a disagreeable drying of the throat and skin, a stimulation of the brain, and an undesired raising of the blood-pressure, to say nothing of its action on the eyes.

The only value of the nitrites is when the blood-pressure is high and the nitrite action is desired on that account.

Coffee or caffein often causes these hearts to become irritable; it certainly raises the blood-pressure, and therefore is not generally advisable. Both tea and coffee should generally be prohibited.

During the acute faint attack camphor is one of the best stimulants. Ammonia may be used, and alcohol is of benefit.

If syphilis is a cause of the condition, iodids are always valuable. If syphilis is not a cause and arteriosclerosis is present, small doses of iodid given for a long period are beneficial, although it may not much reduce the blood-pressure or decrease the plasticity of the blood. Iodid is a stimulant to the thyroid gland, and therefore it is on this account valuable.

The greatest vasomotor stimulant to the heart that we know is thyroid secretion or thyroid extract. Theoretically thyroid extracts should be the treatment for a slow-acting heart. It sometimes seems of benefit to these patients, but it often causes such nervous excitation and irritability as to preclude its use. The dose of thyroid for this purpose would be small, about 3 grains of the dried substance once a day, or 1 grain three times a day. To be of any value the preparation must be good.

The usual precautions against overeating, overdrinking, severe physical exercise, sudden movements, overuse of tobacco, etc., should all be urged on the patient. The disease is sooner or later fatal, although the patient may live some years. The death is generally sudden.

It is understood that this disease must be separated from the condition of bradycardia inherent in a few persons who have a slow pulse throughout their life, without any untoward symptoms.

ARTERIAL HYPERTENSION

It is essential to study this condition, as it has a distinct relationship to the efficiency of the heart, and has a distinct bearing on the efficiency of the left ventricle, and therefore the systemic circulation. However perfectly the heart may act, it cannot properly circulate the blood without a normal tone of the blood-vessels, and of the arterial system in particular. Abnormal vasodilatation seriously impedes the circulation, and may cause shock and death. An increased tone of the arterial system means immediately an increased effort of the heart to overcome the resistance ahead. This soon means an increased activity and later an increased muscle development of the left ventricle. If a high blood-pressure persists, or becomes still higher, it then is a question as to how many months or how many years the left ventricle can stand the increased strain put on it. If the cause of the increased arterial tension is a fibrosis, sooner or later the heart will partake of the disease, and a chronic myocarditis will result. It is therefore evident that an abnormally high blood-pressure is of serious import, and must be studied, interpreted, prevented if possible, and diminished if advisable.

The various apparatus for determining the blood-pressure have already been referred to. No office is completely equipped without a reliable mechanism for determining the blood-pressure, and a portable apparatus is also almost necessary.

It is not the object of this article to discuss the danger of variations in blood-pressure in seriously ill patients, but to discuss the management of patients in whom hypertension is the only abnormal condition. In seriously ill patients dangerous falls of blood-pressure, as shown by repeated investigations, generally occur with a synchronous rise in the pulse-rate. Therefore, as the pulse-rate rises, the blood-pressure may be known to lower, and a dangerous rise means a dangerously low blood-pressure. It would be wise to take the blood-pressure of the convalescent patient more frequently, and not allow him to become ambulatory or return to his work until his blood-pressure has nearly or quite reached what was normal for him. If this was made a routine practice, fewer patients would suffer from weak heart and poor circulation after they had recovered from acute diseases. The blood-pressure would denote whether the myocardium had returned to its perfect capacity for work.

The blood-pressure varies with the age of the person, and is always lower in females than in males. It is probably rarely above 120 mm. of mercury in childhood and youth, and ranges from 110 to 130 in men and women from the age of 20 to about 40, with the average from 30 to 40 of perhaps about 125 mm. From the age of 45 on the blood-pressure generally increases, and as sclerosis advances, if the heart is competent, the blood-pressure rises to 140 or 150 mm., or even more. The blood-pressure in the female is anywhere from 5 to 10 mm. below what is normal for the corresponding age in the male. Any blood-pressure above 150 as an outside limit is probably always abnormal. This does not mean that it needs to be treated, or that it is necessarily wise to attempt to lower it. Blood-pressures higher than 150 become progressively serious, and anything over 200 is generally indicative of renal insufficiency. If there is marked arteriosclerosis, a blood-pressure reading may be very high, and yet not represent the actual pressure of the blood in the arteries; or it may indicate that the artery of the part cannot be collapsed.

As the blood-pressure depends on two factors, the intensity of the left ventricle contraction and the tone of the arteries, the condition of both must be interrogated and analyzed before it can be determined which is overacting or underacting. A too low blood-pressure is generally caused by insufficiency of the heart; it may be caused by loss of tonicity in the arterioles, or by hemorrhage.

Arterial hypertension may be divided into stages. In the first stage the arteries are healthy, but the tone, owing to muscular contraction, is too great. This stage has been called "chronic arterial hypertension" to designate the period at which there is probably no real sclerosis, and at which the production of sclerosis may be prevented or at least deferred. In the second stage the muscular coats of the arteries have become more or less hypertrophied, and degenerative changes have begun in the intima, with perhaps a beginning fibrosis of the external coat of the arteries; in other words, arteriosclerosis is present. A third stage may be recognized in which the fibrosis has far advanced, an atheromatous condition is developing or is present, and the muscular coat of the arteries is becoming insufficient; the heart can no longer increase the arterial tension, and the blood-pressure begins to be diminished. Also at this stage the heart is generally in serious trouble, and the circulation becomes progressively insufficient, with sooner or later left ventricle dilatation, pendent edemas, and possibly obstruction in some of the peripheral arterioles.

ETIOLOGY

The causes are:

1. Products of protein metabolism, if these are abnormally increased or are abnormal in character. These products are abnormally increased by excessive eating of protein, by insufficiency of excretions (especially of the kidney excretion), and by increased general metabolism (as in pregnancy). These protein products are more or less abnormal in gout, in intestinal fermentation and putrefaction, and in any serious impairment of the digestive system. Such abnormal products are irritant to the blood-vessels and cause considerable increase in pressure, unless they are actual toxins, in which case the blood-pressure may be lowered.

2. The arterial pressure is raised by the blood containing a considerable amount of caffein, nicotin or other blood-pressure-raising substance. It is possible that an abnormal secretion of the adrenal glands may cause a continued high blood-pressure.

3. Cerebral stimulation or excitation will cause contraction of the arterioles and raise the blood-pressure. Such an increase in pressure may be due to nervous intensity, excitement, some forms of hysteria, and fear. It is typically caused by prolonged and active brain work.

4. Hard physical labor and sustained muscular exercise will raise the blood-pressure. Strenuous labor and hard athletic work, especially if competitive, and if it requires sustained effort, will cause high blood-pressure earlier than should normally occur in the individual.

5. Any serious infection or serious illness may be the forerunner of an early arteriosclerosis with a previous high blood-pressure. Sclerosis of the arteries may occur without much previous hypertension. This may occur in cases of hyposecretion of the thyroid, and in syphilis, although probably hypertension almost always precedes arteriosclerosis. If the hypertension persists, and especially if it is due to irritants circulating in the blood, endarteritis will probably sooner or later develop.

While hypertension is not a disease, while it many times should not be combated, and while a large amount of effort is expended to combat hypertension without accomplishing that end, still, hypertension many times causes symptoms. These symptoms are more or less troublesome if not serious, and as prolonged hypertension is a sure forerunner of more serious disease, a careful investigation of the causes of the hypertension should be made in every case, and the withdrawal or abatement of the cause and the modification of the food and life of the patient are absolutely essential. There can be no excuse for considering hypertension a condition unworthy of management. Such a patient can generally be improved; troublesome symptoms are generally removed or lessened; and undoubtedly circulatory health, if not actually life, is prolonged by such management. The exact value of treatment, physical or medicinal, in each case is another question, and can be determined by individualizing. Much good can be done by both means; sometimes both means fail. If physical and medicinal measures and the regulation of the diet and life of the patient fail to reduce the blood-pressure, there is a more serious cause back of the condition, generally a slow-going, latent, interstitial nephritis, if arteriosclerosis has not actually begun.

On the other hand, it is not intended in the above statement to declare that all patients with high tension should be treated either medicinally or physically, as it may be unwise in many instances to reduce the pressure. Such patients, however, are generally those who have some other condition or disease present besides the symptom of hypertension.

It should also be recognized that it is becoming universal with insurance companies to believe that a blood-tension above normal for the individual, even if his physical condition is otherwise perfectly satisfactory, will shorten his prospect of life. Arteriosclerosis, cardiac disease, apoplexy and chronic nephritis are all on the increase, and insurance companies believe that an abnormal increase in blood-tension is a forerunner of these ultimate conditions.

SYMPTOMS

The symptoms of hypertension not noted by the patient are the inability to relax from business as readily as normal; an increased rapidity of doing the simple things of life, notably in eating; an increased stimulation from coffee, tea, alcohol and nicotin; a diminished ability to fall asleep readily, or to sleep without dreams and without turning and tossing in bed; an increased amount of urine (not always the case[8]) or at least an increased frequency of urination; an increased irritability, or an aggravation of any disagreeable trait which the patient may have had. The symptoms noted by the patient are a tendency to be short-breathed on slight exertion; a feeling of fulness in the head at times, especially when leaning over, or on any sudden movement; insomnia; slight waves of dizziness; and perhaps slight cardiac pains after an exertion that has added to the work of the left ventricle, which has been long acting against this increased resistance. Any of the last-named symptoms may send the patient to his physician, and the cause should be discovered. Many other little disturbances may send the patient to his physician, and this hypertension may be noted during the examination. To repeat, in any case of hypertension its cause must be sought and its management carefully decided on. The condition should not be neglected by the physician.

Unfortunately, the other side of this story is a troublesome one — the effect of the knowledge that he has hypertension on the man himself — and it is men with whom we are most frequently dealing in this condition. It is no longer of any use to ridicule the general knowledge that hypertension is a factor in life to be considered. Both business and professional men are now well aware that blood-tension is taken in most insurance examinations. It is new to them, and therefore they are the more inquisitive. In the second place, it is pretty close to their hearts, and a man is always more disturbed if he thinks there is anything the matter with his heart than if any other part of the body is at fault. Therefore he will not so readily forget that the blood-pressure reading is important as he will forget that the examination of the urine is important. Consequently, with the knowledge that his blood-pressure is abnormally high he becomes more or less introspective, even perhaps foolishly so, and will raise his blood-pressure by his own fear and anxiety. To meet this condition, which will little by little wear off as the subject becomes less of a novelty, the physician must minimize the importance of a slightly increased blood-pressure, but must still urge the necessity of a change of the man's intense life and unphysiologic eating, drinking and smoking to the prolongation of that man's years, give him no special treatment, see him only infrequently, and aid him in forgetting that his blood-pressure is somewhat high.

---

[8] *Lawrence*: Am. Jour. Med. Sc., September, 1912, p. 330.

On the other hand, if the patient's blood-pressure is very high and the aortic closure is accentuated, we have no more right to tell him to forget it than we have a right to withhold from him the truth, if he had a damaged heart, or if we had found albumin or sugar in his urine. It means that such a patient should be under more or less frequent observation, and certainly something should be attempted to lower the blood-pressure. If it is not readily lowered, then active treatment need not be persisted in.

MANAGEMENT IN ARTERIAL HYPERTENSION

The above-enumerated causes of hypertension suggest its prevention, its cure or its amelioration. To prevent the condition the physician must constantly preach greater care of patients after serious illnesses, less severe athletic competition, less tobacco, less overeating, and, especially, fewer late dinners, less coffee and tea to those who need such advice, and more careful watching of pregnant women as to their food and their excretions, especially the urine, more prevention of intestinal indigestion and colon putrefaction, and if possible less strenuous life and more periods of rest and relaxation.

The very enumeration of these suggestions of prevention of hypertension names the conditions to be investigated and corrected when hypertension is present: the removal of any cause that can be removed, such as nicotin, caffein, too much meat (if it should not temporarily be interdicted entirely), too much liquid of any kind (even water); more strenuous physical exercise, if the patient's habits are sedentary; warm baths, possibly even Turkish baths; better elimination by the bowels, that is, daily sufficient movements; and more rest, with relaxation—all these before physical or medicinal treatments are even considered.

It is a fact that mental as well as physical strenuosity raises the blood-pressure, and that fear, anxiety and even acute worry or sorrow may raise it. The patient's blood-pressure is lowered during the first part of sleep, at night or whenever he lies down and absolutely relaxes.

Brooks and Carroll[9] have shown that the greatest drop of blood-pressure during sleep occurs during the first part of the night's sleep; the blood-pressure then gradually rises, perhaps to reach its maximum the following afternoon. In other words, a sleepless night, or a night in which one is repeatedly aroused, does not give the normal diurnal lowering of blood-pressure, and it is these periods of relaxation that tend to ease the heart and the arteries. It is not the short-lived increased pressure that is serious; it is the continued increased pressure, or the continually repeated increased high pressure.

These investigators also showed that prolonged bed rest or a longer sleep did not tend to lower the pressure any more than the normal sleep and the normal rest,

---

9. Brooks and Carroll: Arch. Int. Med., August, 1912, p. 97.

but that mental rest or psychic rest, or complete mental relaxation did lower blood-pressure at whatever time of day it was accomplished. We therefore come to the conclusion, as far as rest is concerned as a means to lower blood-pressure, that prolonged rest in bed if a patient is nervous, irritated or objecting is of no value, but vacation relaxation when one entirely forgets his business cares and irritations does reduce blood-pressure. In other words, vacations, even short ones, and repeated daily relaxations, if possible, are essential in the treatment of high pressure.

The physical methods of lowering blood-pressure are hydrotherapeutic, hot air (body baking), electric (autocondensation) and electric-light baths. If there is kidney disturbance, or while there are toxins in the blood still uneliminated, nothing can be of more value than various sweat baths, whether given by body baking, by electric light or by Turkish or Russian baths. Sweating is needed, especially if the patient is overweight and his heart is in good condition. Internal congestions are relieved, surface circulation is improved, capillaries are dilated, and the blood-pressure is temporarily always lowered. If the blood-pressure soon returns to a high mark, such treatments should not be too long continued, or if continued should be used only intermittently. Not infrequently, however, some patients are more or less permanently benefited by these treatments, the benefit lasting for months.

The autocondensation treatment always lowers the blood-pressure, but it must be repeated frequently, and the lasting effect of the treatment, unless the patient cooperates in every other way, is rather unsatisfactory. As an addition to other methods of lowering the blood-pressure it is excellent.

In line with the vacation treatment is the advisability, if possible, of selecting a change of climate to a warm one. Cold always increases the blood-pressure; consequently warm climates tend to lower it, and a patient with high blood-pressure in a cold climate will generally have his pressure lowered in a warm one. A corollary to this is that patients with high tension should be warmly clothed during cold weather, and more or less frequent warm baths are of benefit to them.

REDUCING BLOOD PRESSURE

The most positively successful way of reducing blood-pressure is by venesection, and this method should not be forgotten when the blood-pressure at any time is suddenly enormously excessive. How frequently it should be used in seriously high blood-pressure is a question, as many times the return to high pressure is rather rapid, even after such a radical measure. It is therefore doubtful if it is advisable, except when a heart is seriously in acute trouble, with a very high blood-pressure, and the danger of apoplexy or a congestion of the lungs or a paralysis of the heart is imminent.

Drugs to lower the blood-pressure are mostly nitrites, and of these nitroglycerin, sodium nitrite, erythrol tetranitrate and amyl nitrite are the ones used, and the frequency of their use to lower blood-pressure is in the order named. Other drugs used to lower blood-pressure are iodids, thyroid, alkalies, chloral and bromids.

The nitrites all more or less rapidly lower blood-pressure, but the lowering does not last long. The most intense action is from amyl nitrite, but the action from this drug is so rapid and so intense that it should not be used in ordinary hypertension. Nitroglycerin, especially in tablet form, which is the handiest method for its administration, loses its activity more or less rapidly with age. Well-made nitroglycerin tablets will generally be active for months. If given hypodermatically or dropped on the tongue nitroglycerin will act in from one to two minutes, the greatest intensity of its action occurring within a few minutes. The blood-pressure may be lowered temporarily from 20 to 30 mm. of mercury. The action is stated to last not much longer than half an hour. Such an intense action of nitroglycerin is not advisable and not needed in chronic hypertension. The best method of giving nitroglycerin for this purpose is after meals and on going to bed. Whenever it is given on an empty stomach it should be taken with a large amount of water, in order that the action may be slow, and, if possible, not felt by the patient; throbbing of the head is not desirable. The dose may be from 1/500 to 1/100 grain three or four times a day, or every three hours, as desired. While the blood-pressure may not be perceptibly lowered at first, many of the symptoms of hypertension are removed, even with the small dose named, and when 1/200 grain is given four or five times in twenty-four hours, with the rest of the management of the patient, the blood-pressure is decreased.

The dose of sodium nitrite should not be more than 0.03 to 0.06 gm. (½ to 1 grain). It is always best given on a full stomach, as it is likely to irritate. Its action is slower than nitroglycerin, and it does not ordinarily reduce the pressure quite so much, but such reduction often lasts considerably more than an hour.

Erythrol tetranitrate is thought to act more satisfactorily by some clinicians, its effects often lasting longer than the other substances. On the other hand, nitroglycerin seems to be as satisfactory as the other substances named.

To those clinicians who believe that these short reductions of blood-pressure are of little value in the treatment of hypertension, it should be said that even short periods of relaxation, causing reduction of pressure, to say nothing of the physiologic lowering of blood-pressure during sleep, tend to intermit the continued high pressure, and therefore act for good. It is not asserted that the nitrites will cause a continued low blood-pressure. A tolerance is soon established and the dose must be more and more frequently repeated if they are used to their full effect and with the given frequency. On the other hand, if a nitrite is administered four times a day, the blood-pressure often remains slightly lowered, and even a slight lowering of the blood-pressure is valuable. A blood-pressure of 120 mm. may be perfectly satisfactory to the welfare and ability of a person, and if for any reason his blood-pressure drops to 100, he is generally more or less seriously disabled. In other words, a lowering of the pressure even 20 mm. is of importance to a person with high pressure.

Interesting experiments have been made by Lawrence[10] on the systolic and diastolic fall of blood-pressure under the various nitrites, as well as by other measures. He concludes that the diastolic fall of blood-pressure from almost any method of treatment is approximately one-half the systolic fall, and that sodium nitrite reduces the diastolic pressure more rapidly than the more complex compounds.

It is sometimes noticeable that when there is no kidney lesion a very high blood-pressure shows more of a fall under nitrites than does a medium high pressure.

It has long been thought that small doses of iodids, doses that are not sufficient to cause indigestion, will act as vasodilators and render the blood more aplastic or alter the viscosity. This has been recently shown by Capps[11] not to be a fact. Capps concludes that iodids in therapeutic doses are not active vasodilators, and when long continued do not materially affect blood-pressure; also that iodids do not alter the viscosity of the blood to any marked degree. Of course these findings do not mitigate against their value in syphilitic sclerosis, or sclerosis from any cause. As previously stated in these articles, small doses of iodid, daily, stimulate the thyroid gland to greater activity, and one of its activities is to lower the blood-pressure.

---

[10] Lawrence: Arch. Int. Med., April, 1912, p. 409.

[11] Capps, Joseph A.: The Journal A. M. A., Oct. 12, 1912, p. 1350.

If a patient is obese, or if there are other signs of subsecretion of the thyroid, and hypertension is present, thyroid is the proper medicament to give. The dose need not be large, and from 0.1 to 0.2 gm. (1½ to 3 grains) of an active thyroid preparation (the dried gland), given daily for a considerable length of time, will be of value in such instances.

In many instances the blood-pressure is lowered under the action of an alkali such as potassium citrate. It is quite probable that when an alkali is given the diet has also been regulated, and that the alkali may combat some acid irritants. In other words, alkalies may aid in combating gouty tendencies or uric acid tendencies, and these causes of hypertension may be ameliorated. At any rate, they are often of benefit.

Unless a patient had very high tension and for some reason other drugs were unsatisfactory or inadvisable, hydrated chloral would probably rarely be used; but hydrated chloral in small doses, 0.20 to 0.25 gm. (from 3 to 4 grains) given three times a day, after meals, sometimes diminishes the blood-pressure.

The blood-pressure is always lowered by bromids, but of course such treatment is rarely advisable. If insomnia calls for medicinal treatment, bromids and chloral and many other hypnotics will cause a lowering of the blood-pressure, and, if frequently used, may lower the blood-pressure more or less continuously. Such a lowering of pressure, however, is quite likely to be due to some weakening of the heart. In other words, active physical measures should not be taken to lower the blood-pressure while these active drugs are being administered.

CARDIOVASCULAR-RENAL DISEASE

The name "cardiovascular-renal disease" is so suggestive of the exact condition in many instances that it has come to be used as a distinct name for the disease of these three parts of the organism, the name representing the pathology and the prognosis. It is impossible to tell which of these three parts of the organism will become pathologically the most diseased before the patient succumbs. "Bright's disease" is the old name for this condition, but it was used by Bright and is now used by clinicians as a more distinctive term for chronic interstitial nephritis. No patient dies of Bright's disease, however, without having more or less chronic endarteritis and more or less hypertrophy and subsequent chronic myocarditis and dilatation of the heart.

In cardiovascular-renal disease it is often doubtful for a long time whether the patient will finally succumb to cardiac disability, to apoplexy from diseased arteries, to a general serious arteriosclerosis, possibly coronary disease, or to uremia from chronic interstitial nephritis. It is also many times impossible to determine where the general disease first started. The beginning may be chronic endarteritis with preceding hypertension, or a slowly developing chronic interstitial nephritis or cirrhosis of the kidneys; in some instances an insufficient heart may cause such irritations of the kidneys from chronic passive congestion that they become cirrhotic. A more serious passive congestion of the kidneys may occur from an insufficient heart, with profuse albuminuria, all kinds of casts, and finally kidney insufficiency, without developing real chronic interstitial nephritis or the small contracted kidney before death occurs.

Cardiovascular-renal disease generally begins insidiously, with the first indication prolonged hypertension with the symptoms described for that condition. Interstitial nephritis also is always insidious in its development, and for months there may be nothing abnormal in the urine. Whether the hypertension and later chronic endarteritis is the primary lesion, or whether the slowly developing interstitial nephritis is the primary lesion, in which case there is also hypertension, the left ventricle will hypertrophy to overcome the increased resistance in the arteries. This hypertension, this beginning chronic endarteritis, this beginning chronic interstitial nephritis, and this hypertrophy of the left ventricle may all occur without any symptoms at all, and the condition is often discovered only on insurance examination, the man considering himself perfectly well. The insidious beginning of this serious disease should therefore cause careful examination of patients who present themselves for some apparently simple functional disturbance, if the patient is over 40, when this disease is most likely to occur.

The diagnostic symptoms are hypertension and slight enlargement of the heart. The urine may or may not aid in the diagnosis. The absence of albumin and even the absence of casts is no indication that chronic interstitial nephritis has not begun. If the kidneys are suspected, frequent examinations should be made of the urine passed early in the morning, after meals and after exercise; a quantitative test of the twenty-four hours' urine for the excretion of the different elements will often disclose an insufficiency in excretive power. The amount of urine passed by these patients is generally large, and the specific gravity low.

DIAGNOSIS

An early diagnosis of this disease, and the proper management of the patient, may mean several years of life to the patient.

The symptoms of the beginning of this disease are those already mentioned under the subject of hypertension, and in its incipiency it is hypertension. The various rules and regulations laid down for the management of a patient with hypertension are the very methods of treatment for a patient who has actually begun to have cardiovascular-renal disease. If albumin and casts, or either, are found in the urine, the diet must be more carefully regulated than in hypertension without such signs of renal disease, and the treatment becomes that laid down for Bright's disease or chronic interstitial nephritis. If the blood-pressure is nearly 200 or over 200 mm., it is almost indicative of interstitial nephritis, even if the urine, on repeated examinations, is absolutely negative.

As previously stated, the heart, sooner or later, from hypertension, endarteritis, interstitial nephritis, or from any combination of these conditions, will hypertrophy. The heart will be able to overcome the increased blood-pressure for a shorter or longer time, depending on the age of the patient, the previous health of the myocardium, whether or not the patient's general condition (i. e., his nutrition) is good, and whether or not the kidneys are seriously damaged. Sooner or later, however, the left ventricle generally fails, if the patient does not die intercurrently of an apoplexy, an angina pectoris or uremia. The gradually dilating left ventricle begins to give more and more symptoms of insufficiency, and finally, if the dilatation is considerable, the symptoms are the same as they are in valvular disease with loss of compensation. Pendent edemas, more or less serious, develop; there is more or less dyspnea, and albumin, at this period, is pretty sure to be present in the urine, often in large amount. Besides all the disturbances due to the insufficiency of the heart, the two main disturbances, chronic nephritis and myocardial degeneration, progress together.

Not infrequently it is difficult to determine which condition is the more serious at a given time with a given patient, whether it is the failing heart that has made the kidneys worse, or whether the kidneys are on the point of secreting totally insufficient urine both in amount of liquid and in amount of salts, and uremia is imminent. The clue is often given by the edema being pendent, mostly in the feet and legs, gradually coming into the genitals, with possibly ascitic fluid in the peritoneal cavity and with but little puffing or swelling of the hands and face, all of which is indicative of heart failure. Signs of impending uremia are more general edemas, including the hands and face, nausea, vomiting, headache and drowsiness; or, on the other hand, muscular twitchings and sharp neuralgic pains, as indicative of possible convulsions.

It is beyond the scope of this article to treat chronic interstitial nephritis or Bright's disease. The endeavor to prevent this finale of cardiovascular-renal disease has been sufficiently described under hypertension. If cardiac failure occurs, it should be noted that sufficient doses of digitalis may reestablish the power of the left ventricle and remove temporarily, at least, the beginning edemas. On the other hand, if the kidneys are seriously damaged, suppression of urine may occur under full doses of digitalis, and uremia may be precipitated. It is therefore necessary to make a careful decision on the totality of the above-mentioned symptoms to determine if possible whether digitalis to the physiologic effect should be given, or should not be used at all. Very small doses are unsatisfactory in this condition; very large doses are inexcusable; but good full-sized doses to help the heart, if possible, are often justifiable and valuable. The salt-free diet is often indicated, and may do good in preventing and reducing edema.

The Nauheim baths, with sodium chlorid and with carbonic acid gas, are not always efficient in reducing blood-pressure, and may even do serious harm in cardiovascular-renal disease. Sometimes they raise blood-pressure. If it is decided to try them, the patient should be carefully watched after the bath to see whether he is benefited or injured.

Unless the kidneys are in serious trouble and the patient is almost uremic, when the heart fails the blood-pressure is generally much lowered from what it has been, and this is another indication for digitalis.

Patients who have become ill enough to remain in their room or perhaps in bed with this disease are likely to be sleepless and require hypnotics, and it should be remembered that nothing repairs the heart more than good sleep. The hypnotic selected should not be one that would seriously weaken the heart, but almost any of them may be used intermittently, perhaps, first one and then another. If the patient's heart is very weak, even if the kidneys are seriously damaged, there is sometimes nothing more safe than a small sleeping dose of morphin. It is contrary to the general belief and general teaching, but from the cardiac point of view it is the safest of the hypnotics, and from the kidney point of view it generally does as little harm as any other.

If the patient has not reached the point in this disease just described, or if he has had an exacerbation of the trouble and has become improved, he should never think of returning to his active life. If possible, he should go to a warm climate during the cold weather. He should make up his mind that his activity is over, and that his whole time should be spent in prolonging his life by the easiest possible regimen laid down for him, depending on the condition of his kidneys on the one hand and his heart on the other. The diet depends on which part of his disease is the most in evidence; if his heart, the diet should be that previously described under myocarditis and high blood-pressure; if his kidneys, that for chronic nephritis.

DISTURBANCES OF THE HEART RATE

ARRHYTHMIA

While this term really signifies irregularity and intermittency of the heart, it may also be broadly used to indicate a pulse that is abnormally slow or one that is abnormally fast, a rhythm that is not correct for the age, condition and activity of the patient. Irregularity in the pulse-beat as to volume, force and pressure, except such variation in the pulse wave as caused by respiration, is always abnormal. While an intermittent pulse is of course abnormal, it may be caused in certain persons by a condition that does not in the least interfere with their health and well-being.

As to whether a slow or a more or less (but not excessively) rapid pulse in any one is abnormal depends entirely on whether that speed is normal or abnormal to him. As a general rule the heart is more rapid in women than in men. It is always more rapid in children than in adults, and generally diminishes in frequency after the age of 60, unless there is cardiac weakness or some cardiac muscle degeneration. The average frequency of the pulse in an adult who is at rest is 72 beats per minute, but a frequency of 80 is not abnormal, and a frequency of 65 in men is common; 60 is infrequent in men but normal, while up to 90 is not abnormal, especially in women, at the time the pulse is being counted. It should always be considered that in the majority of patients the pulse is slightly increased while the physician is noting its rapidity. Anything over 90 should always be considered rapid, unless the patient is very nervous and this rapidity is considered accidental. Anything below 60 is abnormally slow. In children under 10 or 12 years of age anything below 80 is unusual, and up to 100 is perfectly normal, at least at such time as the pulse is counted and the patient is awake.

Referring to the first article in this series on disturbances of the heart, it will be noted that many physiologic factors must enter into the production of the normal regularity of the pulse. The stimulus must regularly begin in the auricle, must be perfectly transmitted through the bundle of His to the ventricles, the ventricles must normally contract with the normal and regular force, the valves must close normally and at the proper time, the blood-pressure in the aorta must be normally constant to insure the perfect transmission of the blood to the peripheral arteries and to insure the normal circulation through the coronary arteries, and the arterioles must be normally elastic. The nervous inhibitory control through the vagi nerves must also be normal, and there must be no abnormal reflexes of any part of the body to interfere with the normal vagus control of the heart.

While the heart beats from an inherent musculonervous mechanism, nervous interference easily upsets its normal regularity. It may be seriously slowed by nervous shock, fear or sudden peripheral contractions, spasm of muscles, or convulsive contractions, or it may be stimulated to greater rapidity by nervous excitement. It may be slowed or made rapid by reflex irritations, and it may be seriously interfered with by cerebral lesions; it may be made very rapid, or any pressure on the vagi centers in the medulla oblongata will make it very slow. Various kinds of poisons circulating in the blood, both depressants and excitants, may affect the rapidity or the regularity of the heart. Therefore, if it is decided that a given heart is abnormally slow or abnormally rapid, or is decidedly irregular or intermittent, the various causes for such interference with its normal activity must be investigated and admitted or excluded as causative factors.

Many investigations of the rhythm of children's pulses have been made, and some of the later investigations seem to show that not more than 40 per cent. are regular, the remaining 60 per cent. varying from mild irregularity to extreme irregularity.

Scientifically to determine the exact character of a pulse which is discovered by the finger on the radial artery and the stethoscope on the heart to be irregular, tracings of one or more arteries, veins and the heart should be taken. As previously stated in other sections of this series, two synchronous tracings are more accurate than one, and three of more value than two in interpreting the exact activity and regularity of the heart.

ETIOLOGY

The cause of an irregularly acting heart in an adult may be organic, as in the various forms of myocarditis, in broken compensation of valvular disease, Stokes-Adams disease, coronary disease, auricular fibrillation, cerebral disease, and toxemias from various kinds of serious organic disease. The cause may be more or less functional and removable, such as tea, coffee, alcohol, tobacco, gastric indigestion and intestinal fermentation or putrefaction; or it may be due to functional disturbances of the heart such as that due to what has been termed extrasystole, or to irregular ventricular contractions. A frequent cause of irregular heart action in women, more especially of increased rapidity, is hyperthyroidism.

There may be an arhythmia due to some nervous stimulation, probably through the pneumogastric, so that the pulse varies abnormally during respiration, being accelerated during inspiration and retarded during expiration more than is normally found in adults. This condition is frequent in children, and is noticed in neurotic adults and sometimes during convalescence from a serious illness. Nervous and physical rest, with plenty of sleep and fresh, clean air so that the respiratory center is normally stimulated, will generally improve this condition in an adult.

Extrasystoles causing arhythmia give a more or less regularly intermittent pulse, while the examination of the heart discloses an imperfect beat or the extrasystole which is not transmitted or acted on by the ventricles, and hence the intermittency in the peripheral arteries. This condition may be due to some toxemia, nervous irritability, or some irritation in the heart muscle. Good general elimination by catharsis, warm baths to increase the peripheral circulation, a low diet for a few days, abstinence from any toxin that could cause this cardiac irritation, extra physical and mental rest, sometimes nervous sedatives such as bromids, and perhaps a lowering of the blood-pressure bv nitroglycerin if such is indicated, or an increase of the cardiac tone by digitalis if that is indicated, will generally remove the cardiac irritation and prevent the extrasystoles, when the heart will again become regular. It should be carefully decided whether there is beginning heart-block or beginning Stokes-Adams disease, in which case digitalis should not be used. This disease is not frequent, while extrasystoles of a functional character are very frequent. Sometimes this functional disease persists without any apparent injury to the individual as long as the ventricle does not take note of these extra auricular systoles and does not also become extra rapid. If the ventricle does contract with this increased rapidity it soon wears itself out, and the condition becomes serious.

In this kind of arhythmia, if there are no contraindications to digitalis, it is the logical drug to use from its physiologic activities, slowing the heart by its action on the vagi and causing a steadier contraction of the heart; clinically this treatment is generally successful. If digitalis should cause the individual heart to become more irritable, it is acting for harm, and should be stopped.

TREATMENT

One has but to refer to the enumerated causes of irregular heart action to determine the treatment. In that caused by extrasystole the treatment has just been suggested. In irregular heart caused by serious cardiac or other lesions the treatment has already been described, or is that of the disease that has a badly acting heart as a complication. If the irregularity is caused by toxins, the treatment is to stop the ingestion of the toxin and to promote the elimination of what is already in the system; how much of the irregularity was due to the toxin and how much is inherent disturbance in the heart can then be determined. If the cause of a toxemia developed in the system, perhaps most frequently from intestinal putrefaction, increased elimination and a regulation of the diet will cure the condition.

The valvular lesions most likely to cause irregular action of the heart are mitral insufficiency or mitral stenosis. The lesion that is most likely to cause auricular fibrillation and more or less permanently irregular heart is perhaps mitral stenosis. Another frequent cause of more or less permanent irregularity is the excessive use of alcohol.

While an irregular pulse and an irregular heart are always of more or less serious import, still, as the extrasystoles of the auricle are better understood and more frequently recognized, and the habits and life of the patients (most frequently men) are regulated and revised, frequently a pulse and heart that would be rejected by any medical examiner for an insurance company becomes, in a few weeks or a few months, a perfectly acting heart, and remains so sometimes for years. It also is not quite determinable whether a heart that is so misbehaving has a recurrence of such misbehavior more readily than a heart that has never been so affected. However this may be, the cause having been determined or presumed by the physician, it should be so impressed on the patient that he does not again repeat the insult to his heart.

AURICULAR FIBRILLATION

While auricular fibrillation is a clinical entity, it is often difficult of diagnosis, and sometimes can be excluded only by treatment and the results of treatment, or by watching the patient for some time. When completely present, it consists really of a paralysis of the auricles; normal systolic contractions of the auricles do not occur, although there are little rapid twitchings of different muscle fibers of the auricles, which give the name to the condition.

In 1902 Dr. James Mackenzie of England discovered that in these irregular hearts the auricular wave in the jugular tracings did not occur, and assumed that the auricle was more or less paralyzed. In 1908 he concluded that from a stimulus, perhaps occurring in the auriculoventricular node, both auricles and ventricles contracted synchronously, and to these he gave the term of "nodal rhythm." But in 1909 Lewis[12] determined by experimentation that when fibrillations were caused fibrillation, and that the ventricle fortunately generally in the dog the tracings were perfectly similar to those observed by Mackenzie, although Cushny and Edmonds, in 1906, had already suggested this similarity. It is now known that there is no nodal rhythm in auricular responds to only a certain number of the auricular stimuli. The auricle loses coordinate activity but is probably not paralyzed.

The irregular pulse in auricular fibrillation is more or less distinctive, being generally rapid, from 110 upward. Occasionally the pulse-rate may be much slower, if the heart is under the influence of digitalis. The irregularity of the pulse in this condition is excessive; the rate, strength and apparent intermittency during a half minute may not at all represent the condition in the next half minute, or in the next several minutes. It has been thought that auricular fibrillation, while prevented many times by digitalis, is perhaps incurable. This is probaly not true in the early stages of the condition. If digitalis does not cure the irregularity, the condition has been termed the "absolutely irregular heart." Other terms applied to the condition have been "ventricular rhythm," "nodal rhythm" and "rhythm of auricular paralysis." The condition of the pulse has been Latinized as *pulsus irregularis perpetuus.*

---

[12] Lewis, Thomas A : Heart, i, 311; also Brit. Med. Jour., Jan. 13, 1912.

While the condition is best diagnosed by tracings taken simultaneously of the heart, jugular and radial, still the jugular tracing is almost conclusive in the absence of the auricular systolic wave. The radial tracing is exceedingly suggestive, and if it is taken with a careful stethoscoping of the heart, an almost certain presumptive diagnosis may be made.

OCCURRENCE

This condition of auricular fibrillation occurs occasionally in valvular disease, and perhaps most frequently in mitral stenosis; but it can occur without valvular lesions, and with any valvular lesion. If it occurs in younger patients, valvular disease is likely to be a cause; if in older patients, sclerosis or myocardial degeneration is generally present.

It may also follow depressing infections such as diphtheria, or some infection that has caused a myocarditis. Rarely this fibrillation may be caused by some of the drugs used to stimulate the heart.

It is astonishing how few symptoms may be present with auricular fibrillation and an absolutely irregular heart action. The patient may be able to perform all of his duties, however strenuous, until coincident, concomitant or causative ventricular weakening and dilatation or broken compensation occurs, and then the symptoms are those due to the cardiac failure. Often in the first stage of this weakening and later fibrillation of the auricles the patient may recognize the cardiac irregularity and disturbances. Generally, however, he soon becomes accustomed to the sensations, and, unless he has cardiac pains or dyspnea, he becomes oblivious to the irregularity. At other times he may be conscious of irregular, strong throbs or pulsations of the heart, as such hearts often give an occasional extra sturdy ventricular contraction. These he notes. Real attacks of tachycardia may be superimposed on the condition. Sooner or later, however, if the condition is not stopped, cardiac weakness and loss of compensation, with all the usual symptoms, occur. It seems to be probable that more than half of all cases of heart failure are due to auricular fibrillation, or at least are aggravated by it.

TREATMENT

The condition may be stopped by relieving the heart and circulation of all possible toxins and irritants, and by the administration of digitalis. One attack is frequently followed by others, perhaps of longer duration. Occasionally, however, the patient may be observed for many years without the condition again being present. If the pulse, in spite of treatment, is permanently irregular, and auricular insufficiency is permanent, the patient is of course in danger of cardiac failure; but still he may live for years and die of some other cause than heart failure. The prognosis seems to be better when the pulse is not rapid—below a hundred. This shows that the ventricles are not much excited and do not tend to wear themselves out.

Any treatment that lowers the heart-rate is of advantage, such as the stopping of tea and coffee, the administration of digitalis, and an increased amount of rest and quiet. Digitalis should be increased gradually until a fair dose is given, and it is better to administer one dose a day than several. If it causes undesired symptoms, such as cardiac pain, a tight feeling in the chest, nausea or vomiting, or a diminished amount of urine, it is not acting well and should be stopped. If the pulse is gradually slowed to about what is normal, its action should be considered successful.

If the pulse is intermittent and there is apparently a heart-block, Stokes-Adams disease should be considered as possibly present, and digitalis would be contra-indicated and would do harm.

A scientific indication as to whether a heart is disturbed through the action of the vagi nerves or whether the disturbance is due to muscle degeneration may be obtained by the administration of atropin. Talley[13] of Philadelphia shows the diagnostic value of this drug. It is a familiar physiologic fact that stimulation of the vagi slows the heart or even stops it. Stimulation of these nerves by the electric current, however, does not destroy the irritability of the heart; indeed, the heart may act by local stimulation after it has been stopped by pneumogastric stimulation. It is also a well-known fact that anything that inhibits or removes vagus control of the heart allows the heart to become more rapid, since these nerves act as a governor to the heart's contractions. Under the influence of atropin the heart-rate is increased by paralysis of the vagi. Talley states that a hypodermic injection of from 1/50 to 1/25 grain of atropin produces the same paralytic and rapid heart effect in man. He advises the use of 1/25 grain of atropin in robust males, and 1/50 grain in females and in less robust males, and he has seen no serious trouble occur from such injections. The throat is of course dry, and the eyesight interfered with for a day or more, but Talley has not seen even insomnia occur, to say nothing of nervous excitation or delirium. Theoretically, however, before such atropin dosage, an idiosyncrasy against belladonna should be determined.

---

[13] Talley, James: Am. Jour. Med. Sc., October, 1912.

The value of such an injection rests on the fact that atropin thus injected will increase the normal heart from thirty to forty beats a minute, and Talley believes that if the heart-beat is increased only twenty or less, if the patient has not been suffering from an exhausting disease, it shows "a degenerative process in the cardiac tissue which makes the outlook for improvement under treatment unpromising." He also believes that when the heart in auricular fibrillation is increased the normal amount or more than normal, the prognosis is good. He still further advises in auricular fibrillation an injection of atropin before digitalis has been administered, and another after digitalis is thoroughly acting. Comparison of the findings after these two injections will determine which factor, vagal or cardiac tissue, is the greater in the condition present. The patients with a large cardiac factor are the ones who may be more improved by the digitalis treatment than those in whom the fibrillation is caused by vagus disturbance.

BRADYCARDIA

The first decision to be made is what constitutes a slow pulse or slow heart. A pulse below 58 or 60 beats per minute should be considered slow, and anything below 50 should be considered abnormally slow and a condition of more or less suspicion. A pulse from 45 to 50 per minute occasionally occurs when no pathologic excuse can be found, but such a slow rate is unusual. Before determining that the heart is slow, it must of course be carefully examined to determine if there are beats which are not transmitted to the wrist; also whether a slow radial rate is not due to intermittency or a heart-block. Auricular fibrillation, while generally causing a rapid pulse (though by no means all beats are transmitted to the peripheral arteries), may cause a slow pulse because some of the contractions of the heart are not transmitted.

While any pulse-rate below 50 should be considered abnormal and more or less pathologic, still a pulse-rate no lower than 60 may be very abnormal for the individual. For athletes and those who work hard physically a slow pulse is normal. Such hearts are often not even normally stimulated by high fever, so that the pulse is unusually slow considering the patient's temperature, unless inflammation of the heart has occurred.

Some chronic diseases cause a slow pulse; this is especially true of chronic interstitial nephritis. In fact, it may be stated that any disease or condition that increases the blood-pressure generally slows the pulse, unless the heart itself is affected. This is true of hypertension, of arteriosclerosis, of nicotin unless the heart has become injured, and often of caffein, unless it acts in the individual as a nervous stimulant. Chronic lead-poisoning causes a slow pulse on account of the increased blood-pressure.

A slow pulse may occur during convalescence from acute infections such as typhoid fever and pneumonia, and sometimes after septic processes. While it may not be serious in these conditions, it should always be carefully watched, as it may show a serious myocarditis.

While weakness generally and myocarditis, at least on exertion or nervous excitation or after eating, cause a heart to be rapid, still such a heart may act sluggishly when the patient is at rest, so that he feels faint and weak and disinclined to attempt even the slightest exertion. In such a condition calcium, iron and strychnin, not too frequently or in too large doses, and perhaps caffein, are indicated. Camphor is always a valuable stimulant, more or less frequently administered, during such a period of slow heart. This slow heart sometimes occurs after rheumatic fever; it is quite frequent after diphtheria, and may show a disturbance of the vagi.

Although the prognosis of such slow hearts after serious illness is generally good, a heart that is too rapid after illness is often more readily brought to normal by proper management than a heart that is too slow. Either condition needs proper treatment and proper management.

It is well recognized that serious, almost major hysteria may be present and the heart not only not be increased, but it may even be slowed. The heart in this condition of course requires no treatment. In cerebral disturbances, especially when there is cerebral pressure, and more particularly if there is pressure in the fourth ventricle, the pulse may be much slowed. It is often slowed in connection with Cheyne-Stokes respiration. It may be very slow after apoplexy, and when there are brain tumors. It is often much slowed in narcotic poisoning, especially in opium, chloral and bromid poisoning. Serious toxemia from alcohol may cause a heart to be very slow. It is more likely, however, to cause a heart to be rapid, unless there is actual coma.

A frequent condition causing a slowing of the heart is the presence of bile in the blood, typically true of catarrhal jaundice. Uremic poisoning and acidemia and coma of diabetes may cause a pulse to be very slow.

Not infrequently after parturition the heart quiets down from its exertion to a rate below normal. If the urine is known to be free from albumin and casts and there are no signs of impending eclampsia, the slow pulse is indicative of no serious trouble; but the urine should be carefully examined and a possible uremia or other cause of eclampsia carefully considered. Sometimes with serious edema and after serious hemorrhage the heart becomes very slow, unless some exertion is made, when it will beat more rapidly than normal. This probably represents a diminished cardiac nutrition.

The cardiac lesions that cause a pulse to be slow are sclerosis or thrombosis of the coronary arteries, fatty degeneration of the myocardium and Stokes-Adams disease.

It is therefore seen that when a pulse is slower than normal, even below 65 beats per minute, the cause should be sought. If no functional or pathologic excuse is discovered, it must be considered normal for the individual, and as above stated, even 58 or 60 beats per minute are in many instances normal for men. This is especially true with beginning hypertension, and may be true in young men who are athletic or who are oversmoking but are not being poisoned by the nicotin, as shown by the fact that their hearts are not rapid, that they are not having cardiac pains, that they do not perspire profusely, and that they do not have muscle cramps. A pulse of from 50 to 55 is liable to be seriously considered by an insurance company in deciding the advisability of the risk, and below 50 must be considered as abnormal.

SYMPTOMS

If a person has been long accustomed to a slow-acting heart, there are no symptoms. If the heart has become slowed from disease or from any acute condition, the patient is likely to feel more or less faint, perhaps have some dizziness, and often headache, which is generally relieved by lying down. Sometimes convulsions may occur, epileptiform in character, due possibly to anemia or irritation of the brain. If the slow heart does not cause these more serious symptoms, the patient may feel weak and unable to attend to his ordinary duties. As previously urged, an abnormally slow heart after serious illness should be as carefully cared for as a too rapid heart under the same conditions. Probably often a myocarditis and perhaps some fatty degeneration are at the base of such a slowed heart after serious infections.

A heart that has not always been slow but has gradually become slow with the progress of hypertension and arteriosclerosis will often disclose on post-mortem examination serious lesions of the coronary arteries.

Deficiency in the thyroid secretion will always cause a heart to be slower than normal. The more marked and serious the hypothyroidism, the slower the heart is likely to be. When such a condition is diagnosed the treatment is thyroid extract; or if the insufficiency is not great, small doses of an iodid should be given. In either case it is sometimes astonishing how rapidly a slow, sluggishly acting heart improves and how much improvement there is in the mental condition of the patient.

In acute slowing of the heart, as in syncope, the patient must immediately lie down with the head low, possibly with the feet and legs elevated, and all constricting clothing of the abdomen and chest should be removed. Whiffs of smelling-salts may be given; whisky, brandy or other quickly-acting stimulant, not much diluted, may also be given. Aromatic spirit of ammonia, camphor, a hypodermatic dose of strychnin or atropin if deemed necessary, a hot-water bag over the heart, and massaging of the arms and legs to aid the return circulation, are all means which are generally successful in restoring the patient's circulation to normal. Caffein is another valuable stimulant, perhaps best administered as a cup of coffee. Digitalis is not indicated; neither is nitroglycerin, unless the slow heart is due to cardiac pain or to an angina.

Some patients have syncopal attacks with the least injury or with any mental shock. Such patients as soon as restored are as well as ever. Other patients who faint or have attacks of syncope should remain at rest on a couch or bed for some hours.

A tangible cause being discovered for an unusually slow heart is sufficiently indicative of the treatment not to require further comment. While generally toxins from intestinal indigestion make a heart irritable and more rapid, sometimes they slow a heart, and in such cases the heart will be improved when catharsis has been caused and a modification of the diet is ordered.

PAROXYSMAL TACHYCARDIA

This condition is generally termed by the patient a "palpitation," and palpitation of the heart is recognized by most physicians as meaning a too rapidly-acting heart, the term "tachycardia" being reserved for an excessive rapidity of the heart. Some persons normally have a pulse and heart too rapid; children more or less constantly have a heart-beat of from 90 to 100. Women have more rapid heart action than men, and it becomes more rapid with their varying functions, specifically increasing in rapidity before, and perhaps during, menstruation. Many patients have a rapid heart action with the slightest increase in temperature and in any fever process. Some have a rapid heart action after the least exertion without any cardiac lesion or assignable excuse for such rapidity. Others have a rapid heart with mental activity and excessive excitement. Therefore in deciding that a heart is abnormally rapid, one must individualize the patient.

During or after illness many patients are said to have palpitation when the real cause is an unhealed myocarditis. Tuberculosis almost invariably causes increased heart action, even when there is no fever. All high fever increases the heart's action, but not so markedly in typhoid fever as in other fevers; in fact, the heart in typhoid fever, during the early stages, is likely to be slower than the temperature would seem to call for. In anemia when the patient is active the heart is generally rapid. The rapid heart from cardiac disease has already been considered. For the palpitation or rapid heart just described there is little necessity for other treatment than what the acute or chronic condition would call for. With proper management the condition will improve unless the patient has an idiosyncrasy for intermittent attacks of slightly rapid heart, as from 100 to 120 beats per minute.

A permanently rapid heart, when the patient is at rest, is generally due to hypersecretion of the thyroid, which will be discussed later. Paroxysmal tachycardia, the condition now under discussion, is a name applied to very rapid heart attacks in persons who are more or less subject to their recurrence. They may occur without any tangible excuse, and are liable to occur during serious illness, after a large meal, after a cup of tea or coffee, or after taking alcohol. The heart may beat as rapidly as from 150 to 200 times a minute, or even more, with no other symptoms than a feeling of constriction or tightness in the chest, an inability to respire properly and a feeling of "air hunger." The patient almost invariably must sit up, or at least have his head raised. Attacks of cardiac delirium may occur with serious lesions of the heart, as valvular disease or sclerosis, but paroxysmal tachycardia occurs in certain persons without any tangible cardiac excuse. The auricles of the heart may act more energetically than normal and precede as usual the ventricular contraction; or the auricles and ventricles may contract almost together — a so-called "nodal" type of contraction. Rarely does a patient die of paroxysmal tachycardia. The length of time the attack may last varies from a few minutes to an hour, or even for a day or more.

MANAGEMENT

There is no specific treatment for paroxysmal tachycardia. What is of value in one patient may be of no value in another; in fact, drugs are rarely successful in ameliorating or preventing the condition. Patients who are accustomed to these attacks often learn what particular position or management stops the attack. Sometimes a patient rises and walks about. Sometimes an ice-bag over the heart will stop the attack. If there is no serious illness present, and no serious cardiac disease causing the condition, and a patient is known to have an overloaded stomach or bowels, an emetic or a briskly-acting cathartic is the best possible treatment. The attack often terminates as suddenly as it begins, without leaving any knowledge as to which particular treatment has been beneficial. A patient who is well and has an attack of tachycardia should be allowed to assume the position that he finds to give him the most comfort, and to use the means of stopping his attack which he has found the most successful. In the absence of his success or of his knowledge of any successful treatment, a hypodermatic injection of 1/6 or even ¼ grain of morphin sulphate is often curative. Atropin should not be given, as it may increase the cardiac delirium.

If an attack lasts more than an hour or so, one of the best treatments is the bromids, which should be given either as potassium or sodium bromid in a dose of 2 or 3 gm. (30 or 45 grains) at once. Sometimes one good-sized dose of digitalis may be of benefit, but it is often disappointing, and unless there is a valvular lesion with signs of broken compensation, it is rarely indicated. It should also be remembered that, if the patient is receiving digitalis in good dosage for broken compensation, cardiac delirium may be caused by an overaction of the digitalis. Such overaction would be indicated by previous symptoms of nausea, vomiting, intestinal irritation, a diminished amount of urine, headache and a tight, band-like feeling in the head, cold hands and feet, and a day or two of very slow pulse. If none of these symptoms is present, though a patient has received digitalis for broken compensation, a tachycardia occurring might not contra-indicate digitalis, as much of the digitalis on the market is useless, and a patient may not actually have been obtaining digitalis action.

If the tachycardia occurs in a patient with arteriosclerosis, especially if there is much cardiac pain, nitroglycerin is of advantage; also warm foot-baths. If there is prostration and a flaccid, flabby abdomen, a tight abdominal bandage may be of benefit.

Gastric flatulence, while perhaps not a cause of the tachycardia, is liable to develop and be a troublesome symptom. Anything that causes eructations of gases is of benefit, as spirit of peppermint, aromatic spirit of ammonia or plain hot water. If there is hyperacidity of the stomach, bicarbonate of sodium or milk of magnesia will be of benefit.

The ability of some patients to stand a rapid heart action without noting it or being incapacitated by it is astonishing. It may generally be stated that a rapid heart is noted, and a pulse above 120 generally prostrates, at least temporarily, a patient who is otherwise well, provided the cause is anything but hyperthyroidism. A patient who has hypersecretion of the thyroid will be perfectly calm, collected, often perhaps not seriously nervous, and, with a heart beating at the rate of 140, 150, 160 and even 170 per minute, will state that she has no palpitation now, although she sometimes has it. A heart thus fast, with a patient not noting it and not prostrated by it, is almost diagnostic of a thyroid cause.

Some patients, both men and women, cannot take even a small cup of tea or coffee without an attack of paroxysmal tachycardia. Such patients, of course, quickly learn their limitations.

HYPERTHYROIDISM

The presence of a well-marked case of Graves' disease or exophthalmic goiter is not necessary for the secretion of the thyroid to be increased sufficiently to cause tachycardia; in fact, an increased heart rapidity in women often has hyperthyroidism as its cause. The thyroid gland hypersecretes in women before every menstrual period and during each pregnancy, and with an active, emotional, nervous life, social excitement, theaters, too much coffee, and, unfortunately to-day among women, too much alcohol, it readily gives the condition of increased secretion; and the organ that notes this increased secretion the quickest is the heart.

The tachycardia of a developed exophthalmic goiter is difficult to inhibit. Digitalis is of no avail, and no other single medicinal treatment is of any great value. The tachycardia will improve as the disease improves. On the other hand, nothing is more serious for this patient than her rapid heart, and if it cannot be soon slowed, operative interference is absolutely necessary. If the rapid heart continues until a myocarditis has developed and a weakening of the muscle fibers occurs or dilatation is imminent or has actually occurred, operative interference is serious, and most patients under these conditions die after a complete operation, i. e., the removal of from one-half to two-thirds of the thyroid. In such cases the only excusable operative interference is the graded one, namely, the tying of first one artery and then another of the thyroid to inhibit the blood-supply to the gland so that it may not furnish so much secretion. If the heart then improves, a more radical operation may be done without much serious danger. Therefore the working rule should be that, if a heart does not quickly improve under medical management, operative interference should not be delayed until the heart has become degenerated.

If tachycardia is the only serious symptom present in a patient who is considered to have hyperthyroidism, it may generally be successfully treated by insistence on quiet, cessation of all physical and exciting mental activities, more or less complete rest, the absolute interdiction of all tea, coffee, or other caffein-bearing preparations, total abstinence from alcohol, the restriction to a milk, cereal and fruit diet (the withdrawal of all meat from the diet), the administration of calcium, as the calcium glycerophosphate in dose of 0.3 gram (5 grains) in powder three times a day, and for a time, perhaps, the administration of bromids. If the depressing action of bromids on the heart is counteracted by the coincident administration of digitalis, they will act only for good by quieting the nervous system and more or less inhibiting the secretion of the thyroid gland.

If a patient has exophthalmic goiter fully developed, absolute rest in bed, with the treatment above outlined, should soon cause improvement. If it does not, the operative treatment as above advised should be considered. If myocarditis has been diagnosed, the minor operations should be done if the patient does not soon improve. The prolongation of the treatment depends on the condition and the amount of improvement.

If the physician is in doubt as to whether this particular tachycardia is caused by hyperthyroidism or not, the administration of sodium iodid in doses of 0.25 gm. (4 grains) three times a day will make the diagnosis positive within a few days. If the trouble is due to hyperthyroidism, all of the symptoms will be aggravated; there will be more palpitation, more nervousness, more restlessness, more sweating and more sleeplessness. In such cases the iodid should be stopped immediately, of course, and the proper treatment begun.

TOXIC DISTURBANCES AND HEART RATE

Under this head it is not proposed to consider disturbances of the heart due to infections, to cardiac disease, or to localized or general acute or chronic disease, but to discuss disturbances due to the absorption of irritants from the intestines, and to alcohol, tobacco and caffein.

It is hardly necessary to repeat that various toxins which may seriously irritate the heart may be absorbed from the intestines during fermentation or putrefactive processes in either the small or the large intestines. The heart may be slowed by some, made rapid by others, and it is often made irregular. The relation of the absorption of intestinal toxins to increased blood-pressure has already been described, and the necessity of removing from the diet anything that perpetuates or increases intestinal indigestion in all cases of high blood-pressure has already been referred to several times. The indications that such a condition of the intestines is present are irregular action of the bowels, a large amount of intestinal gas, sometimes watery stools, often a coated tongue, and the presence of indican in the urine.

INTESTINAL PUTREFACTION

The most successful procedure in the management of intestinal putrefaction is to remove meat from the diet absolutely. Laxatives in some form are generally indicated, and one of the best is agar-agar. Of course aloin and cascara are always good laxatives, with an occasional dose of calomel or saline, if such seem indicated. Some of the solid peroxid-of-hydrogen-carrying preparations (magnesium peroxid, calcium peroxid, see N. N. R., 1912, p. 174) have been advised as bowel antiseptics, but they are not more successful than many of the salicylic acid preparations, and perhaps none is more efficient than salol (phenyl salicylate) in a dose of 0.3 gm. (5 grains), three or four times a day. Washing out the colon with high injections is often of great value, but should not be continued too long lest the rectum become habituated to distention and bowel movements not take place without an enema.

Lactic acid bacilli, best the Bulgarian, are often of value in intestinal fermentation. A tablet may be eaten with a little milk-sugar or a small lump of cane-sugar after each meal. Or yeast may be taken in the form of brewer's yeast, a tablespoonful in a glass of water, two or three times a day, or one-sixth of an ordinary compressed yeast cake dissolved in a glass of water and taken once or twice a day. Or various forms of lactic acid fermented milk may be successful.

Any particular food that causes fermentation in the intestine of the subject should be eliminated from his diet; in other words, the patient must be individualized as to fruits, cereals and vegetables but, as above stated, meat should ordinarily be withheld for a time at least.

ALCOHOL

Enough has already been said of the value and limitations of alcohol as a therapeutic agent. As a beverage, when constantly used, it is likely to cause obesity, gastric indigestion, arteriosclerosis, myocardial degeneration, chronic nephritis and cirrhosis of the liver. Its first action is undoubtedly as a food, if not too large amounts are taken, and therefore it is a protector of other food, especially of fat and starch. A habitué, then, especially if he has reached the age at which he normally adds weight, increases his tendency to obesity, and the first mistake in his nutrition is made. If he takes too much alcohol when he eats or afterward, his digestion will be interfered with. Sooner or later, then, gastritis and stomach indigestion develop, with consequent intestinal indigestion. If he takes strong alcohol, like whisky, on an empty stomach, he may sooner or later cause serious disease of the mucous membrane of the stomach, first chronic gastritis, and later atrophy of the glands of the stomach. Alcohol with meals which contain meat tends to the production of an increased amount of uric acid. Alcohol taken before meals on an empty stomach causes sudden vasodilatation after absorption. It goes quickly to the liver, irritates it, and little by little causes congestions of the liver, so that sooner or later sclerosis of this organ develops.

Alcohol probably causes arteriosclerosis not by its action *per se,* but indirectly by causing gastro-intestinal indigestion and insufficiency of the liver, as a result of which more toxins circulate in the blood, tending to produce arteriosclerosis. Sooner or later these irritants cause kidney irritation, and chronic interstitial nephritis may develop. Just which process becomes the farthest advanced and finally kills the patient is an individual proposition and cannot be foretold. The finale may be cirrhosis of the liver, uremia, arteriosclerosis, apoplexy or myocarditis with dilatation or coronary disease.

While small, more or less undiluted doses of alcohol, as whisky or brandy, may cause quick stimulation of the heart by reflex irritation of the esophagus and stomach, vasodilatation occurs as soon as the alcohol is absorbed, and if large doses are absorbed, vasomotor paresis may occur, temporarily at least.

During acute fever processes with an increased pulse-rate, provided shock or collapse is not present, small or medium-sized doses of alcohol, by dilating the peripheral blood-vessels and increasing the peripheral circulation, may relieve the tension of the heart and slow the pulse by the equalization of the circulation. Some of this action may be due to the narcotic effect of alcohol on the cerebrum. Alcohol may thus in many instances act for good. Overdoses, as shown by cerebral excitation, flushing of the face and increased pulse-rate, will do harm; in fact, many a patient with a serious illness, as typhoid fever or pneumonia, is made delirious by alcohol, and the delirium is not a complication of the disease. Large doses of alcohol in shock or collapse are contraindicated.

Chronic overuse of alcohol may cause chronic myocarditis and fatty degeneration of the heart, with later weakening of the heart muscle and dilatation.

In acute alcohol poisoning the pulse may become very rapid and weak, and the patient may die of heart failure. This is often seen in delirium tremens. The administration in this condition of enormous doses of digitalis by the stomach is inexcusable, and the reason that such patients survive such digitalis poisoning is that the stomach does not absorb during this cardiac prostration.

A treatment as successful as any in this heart weakness in delirium tremens is morphin sulphate, ¼ grain, and atropin, 1/150 grain, given hypodermatically, with the administration of digitalinum hypodermatically for its later action on the heart. If the heart is contracting very rapidly, an ice-bag over the precordia will often quiet it. If the pulse is very weak, the cerebral sedatives more frequently used in delirium tremens, such as chloral, bromids, paraldehyd, etc., are generally contra-indicated. A hot foot-bath and an ice-cap on the head sometimes aid in establishing a more general equalization of the circulation. It may often be necessary to administer strychnin, although if the patient is greatly excited it should be withheld as long as possible. For the same reason camphor, coffee and other cardiac stimulants which cause cerebral excitation should be withheld.

If the patient is in alcoholic coma, the pulse is generally slow, although it may be of low pressure unless the patient is otherwise diseased. Caffein or coffee is here indicated, and the patient should be kept warm lest he lose necessary heat. The stomach should be emptied by an emetic, often best by apomorphin hypodermatically, unless the pulse is excessively weak. Strychnin may also be given, and digitalin, hypodermatically, if it seems indicated. Camphor is another cardiac and cerebral stimulant that is valuable in these cases.

The treatment of an actual degeneration of the heart from overuse of alcohol is similar to the same condition from other causes.

CAFFEIN

Caffein can irritate the heart and cause irregularity and tachycardia, especially in certain persons. In fact, some can never take a single cup of coffee without having an attack of palpitation, and many times when coffee and tea have been unsuspected by the patient as the cause of cardiac irritability, their removal from the diet has stopped the symptoms, and the heart has at once acted normally.

Caffein is a stimulant and tonic to the heart, increasing its rapidity and the strength of the contractions. It is also a cerebral stimulant, one of the most active that we possess among the drugs. It increases the blood-pressure, principally by stimulating the vasomotor center and by increasing the heart strength. It acts as a diuretic, not only by increasing the circulatory force and blood-pressure, but also by acting directly on the kidney. This action on the kidney contra-indicates the use of caffein in any form, except in rare instances when there is acute or chronic nephritis. The increased blood-pressure caused by caffein also contra-indicates its use when there is hypertension. Caffein first accelerates the heart and later may slow it and strengthen it; but if the dose is large or too frequently repeated the apex of the heart ceases to relax properly and there is an interference with the contraction of the ventricles, the heart muscle becomes irritable, and a tachycardia may develop.

Therefore when a heart has serious lesions, whether of the myocardium or of the valves, with compensation only sufficient, the action of caffein in any form is contra-indicated. The fact that it raises the blood-pressure, thus increasing the force against which the heart must act, and that it irritates the heart muscle to more sturdy or irregular contraction, indicates that a patient with a heart lesion or with a nervously irritable heart should never drink tea and coffee or take caffein in any beverage.

Many patients cannot sleep for many hours after they have taken coffee or tea, as the cerebral stimulation of caffein is projected for hours after its ingestion. Caffein does not absorb so quickly and therefore does not act so quickly when taken in the form of tea and coffee as it does when taken as the drug or as a beverage which contains the alkaloid. Persons who are nervously irritable, excited and overstimulated cerebrally, with or without high blood-pressure, should not take this cerebral and nervous excitant. This is true in early childhood and in youth, and continues true as age advances, in most persons. It is a crime to present caffein as a soda fountain beverage to children and young persons when the excitement of the age is such as already to overstimulate all nervous systems and all hearts.

A considerable majority of persons over 40 learn that they cannot drink tea or coffee with their evening meal without finding it difficult to sleep. Such patients, of course, should omit this stimulant. Some patients have already recognized this fact and its cause; others must be told. The majority of adults are probably no worse and may be distinctly benefited by the morning cup of coffee and the noon coffee or tea, provided that the amount taken is not large. It seems to be a fact that the drinking of coffee is on the increase, especially as to frequency. Certainly the five o'clock tea, with women, is on the increase, and we must deal with one more cerebral and nervous excitant in our consideration of what we shall do to slow this rapid age.

TOBACCO

In spite of the fact that a large number of men to-day do not smoke, more and more frequently every clinician has a patient who smokes too much. The accuracy with which he investigates these cases depends somewhat on his personal use of tobacco, and therefore his leniency toward a fellow-user. Perhaps the percentage of young boys who smoke excessively is larger than the percentage of men. Whether the term "excessive" should be applied to any particular amount of tobacco consumed depends entirely on the person. What may be only a large amount for one person may be an excessive amount for another, and even one cigar a day may be too much for a person—as much for him as five or more cigars for another. If one is to judge by the internal revenue report it will appear that, in spite of the public school instruction as to the physiologic action of tobacco and its harm, and in spite of the antitobacco leagues, the consumption of tobacco is enormously on the increase.

The most important action of nicotin is on the circulation. Except during the stage when the person is becoming used to the tobacco habit, in which stage the heart is weakened and the vasomotor pressure lowered by his nausea and prostration, the blood-pressure is always raised during the period of smoking. The length of time the increased pressure continues depends on the person, and it is this diminishing pressure that causes many to take another smoke. The heart is slowed by the action of nicotin on the vagi, as these nerves are stimulated both centrally and peripherally. An overdose of nicotin will paralyze the vagi. The heart action then becomes rapid and perhaps irregular. The heart muscle is first stimulated, and if too large a dose is taken, or too much in twenty-four hours, the muscle becomes depressed and perhaps debilitated. The consequence of such action on the heart muscle, sooner or later, is a dilatation of the left ventricle if the overuse of the tobacco is continued.

There is, then, no possible opportunity for any discussion as to the action of tobacco on the circulation. Its action is positive, constantly occurs, and is always to be considered. The only point at issue is as to whether such an activity is of consequence to the individual. The active principle of tobacco is nicotin, besides which it contains an aromatic camphor-like substance, cellulose, resins, sugars, etc. Other products developed during combustion are carbon monoxid gas, a minute amount of prussic acid and in some varieties a considerable amount of furfural, a poison. From any one cigar or cigaret but little nicotin is absorbed, else the user would be poisoned. It is generally considered that the best tobacco comes from Cuba, and in the United States from Virginia. While it has not been definitely shown that any stronger narcotic drug occurs in cigarets sold in this country, it still is of great interest to note that a user who becomes habituated to one particular brand will generally have no other, and the excessive cigaret-smoker will generally select the strongest brand of cigarets. The same is almost equally true of cigar smokers.

Besides the effect on the circulation, no one who uses tobacco can deny that it has a soothing, narcotic effect. If it did not have this quieting effect on the nervous system, the increased blood-pressure would stimulate the cerebrum. Following a large meal, especially if alcohol has been taken, the blood-vessels of the abdomen are more or less dilated by the digestion which is in process. During this period of lassitude it is possible that tobacco, through its contracting power, by raising the blood-pressure in the cerebrum to the height at which the patient is accustomed, will stimulate him and cause him to be more able to do active mental work. On the other hand, if a person is nervously tired, irritable, or even muscularly weary, a cigar or several cigarets will increase his blood-pressure, take away his circulatory tire, soothe his irritability, and stop temporarily his muscular pains or aches and muscle weariness. If the user of the tobacco has thorough control of his habit, is not working excessively, physically or mentally, has his normal sleep at night and therefore does not become weary from insomnia, he may use tobacco with sense and in the amount and frequency that is more or less harmless as far as he is concerned. If such a man, however, is sleepless, overworked or worried, if he has irregular meals or goes without his food, and has a series of "dinners," or uses a lot of alcohol, which gives him vasomotor relaxation, he finds a constantly growing need for a frequent smoke, and soon begins to use tobacco excessively. Or the young boy, stimulated by his associates, smokes cigarets more and more frequently until he uses them to excess.

Just what creates the intense desire for tobacco to the habitué has not been quite decided, but probably it is a combination of the irritation in the throat, especially in inhalers, of the desire for the rhythmic puffing which is a general cerebral and circulatory stimulant, for the increased vasomotor tension which many a patient feels the need of, for the narcotic, sedative, *quieting* effect on his brain or nerves, for the alluring comfort of watching the smoke curl into the air, or for the quiet, contented sociability of smoking with associates. Probably all of these factors enter into the desire to continue the tobacco habit in those who smoke, so to speak, normally.

The abnormal smokers, or those who use tobacco excessively, have a more and more intense nervous desire or physical need of the narcotic or the circulatory stimulant effect of the tobacco, and they smoke more and more constantly. They are largely inhalers, and frequently cigaret fiends.

It is probable that tobacco smoked slowly and deliberately, when the patient is at rest, and when he is leading a lazy, inactive, non-hustling life such as occurs in the warmer climates, is much less harmful than in our colder climates where life is more active. Something at least seems to demonstrate that cigaret-smoking is more harmful in our climate than in the tropics.

It has been shown by athletic records and by physicians' examinations of boys and young men in gymnasiums that perfect circulation, perfect respiration and perfect normal growth of the chest is not compatible with the use of tobacco during the growing period. It is also known that tobacco, except possibly in minute quantities, prevents the full athletic power, circulatorily and muscularly, of men who compete in any branch of athletics that requires prolonged effort.

The chronic inflammation of the pharynx and subacute or chronic irritation of the lingual tonsil, causing the tickling, irritating, dry cough of inhalers of tobacco is too well known to need description.

Many patients who oversmoke lose their appetites, have disturbances from inhibition of the gastric digestion, and may have an irregular action of the bowels from overstimulation of the intestines, since nicotin increases peristalsis. Such patients look sallow, grow thin and lose weight. These are the kind of patients who smoke while they are dressing in the morning, on the way to their meals, to and from their business, and not only before going to bed, but also after they are in bed. It might be a question as to whether such patients do not need conservators. The use of tobacco in that way is absolutely inexcusable, if the patient is not mentally warped. Cancer of the mouth caused by smoking, blindness from the overuse of tobacco, muscular trembling, tremors, muscle cramps and profuse perspiration of the hands and feet are all recognized as being caused by tobacco poisoning, but such symptoms need not be further described here.

The reason for which physicians most frequently must stop their patients from using tobacco, however, is that the heart itself has become affected by the nicotin action. The heart muscle is never strengthened by nicotin, but is always weakened by excessive indulgence in nicotin, the nerves of the heart being probably disturbed, if not actually injured. The positive symptoms of the overuse of tobacco on the heart are attacks of palpitation on exertion lasting perhaps but a short time, sharp stinging pains in the region of the heart, less firmness of the apex-beat, perhaps irregularity of the heart, and cold hands and feet. Clammy perspiration frequently occurs, more especially on the hands. Before the heart muscle actually weakens, the blood-pressure has been increased more or less constantly, perhaps permanently, until such time as the left ventricle fails. The left ventricle from tobacco alone, without any other assignable cause, may become dilated and the mitral valve become insufficient. Before the heart has been injured to this extent the patient learns that he cannot lie on his left side at night without discomfort, that exertion causes palpitation, and that he frequently has an irregularly acting heart and an irregular pulse. He may have cramps in his legs, leg-aches and cold hands and feet from an imperfect systemic circulation. In this condition if tobacco is entirely stopped, and the patient put on digitalis and given the usual careful advice as to eating, drinking, exertion, exercise and rest, such a heart will generally improve, acquire its normal tone, and the mitral valve become again sufficient, and to all intents and purposes the patient becomes well.

On the other hand, a heart under the overuse of tobacco may show no signs of disability, but its reserve energy is impaired and when a serious illness occurs, when an operation with the necessary anesthesia must be endured or when any other sudden strain is put on this heart, it goes to pieces and fails more readily than a heart that has not been so damaged.

If a patient does not show such cardiac weakness but has high tension, the danger of hypertension is increased by his use of tobacco, and certainly in hypertension tobacco should be prohibited. The nicotin is doing two things for him that are serious; first, it is raising his blood-pressure, and second, it will sooner or later weaken his heart, which may be weakened by the high blood-pressure alone. Nevertheless a patient who is a habitual user of tobacco and has circulatory failure noted more especially about or during convalescence from a serious illness, particularly pneumonia, may best be improved by being allowed to smoke at regular intervals and in the amount that seems sufficient. Such patients sometimes rapidly improve when their previous circulatory weakness has been a subject of serious worry. Even such patients who were actually collapsed have been saved by the use of tobacco.

Whether the tobacco in a given patient shall be withdrawn absolutely, or only modified in amount, depends entirely on the individual case. As above stated, no rule can be laid down as to what is enough and what is too much. Theoretically, two or three cigars a day is moderate, and anything more than five cigars a day is excessive; even one cigar a day may be too much.

MISCELLANEOUS DISTURBANCES

SIMPLE HYPERTROPHY

Like any other muscular tissue the heart hypertrophies when it has more work to do, provided this work is gradually increased and the heart is not strained by sudden exertion. To hypertrophy properly the heart must go into training. This training is necessary in valvular lesions after acute endocarditis, and is the reason that the return to work must be so carefully graduated. When the heart is hypertrophied sufficiently and compensation is perfect, a reserve power must be developed by such graded exercise as represented by the Nauheim, Oertel or Schott methods. Anything that increases the peripheral resistance causes the left ventricle to hypertrophy. Anything that increases the resistance in the lungs causes the right ventricle to hypertrophy. The right ventricle hypertrophy caused by mitral lesions has already been sufficiently discussed. The right ventricle also hypertrophies in emphysema, after repeated or prolonged asthma attacks, perhaps generally in neglected pleurisies with effusion, in certain kinds of tuberculosis, and whenever there is increased resistance in the lung tissue or in the chest cavity.

The term "simple hypertrophy" is generally restricted to hypertrophy of the left ventricle without any cardiac excuse—the hypertrophy caused by hypertension and hard physical labor. It is well recognized that it hypertrophies with hypertension and with chronic interstitial nephritis. It also becomes hypertrophied when the subject drinks largely of liquid—water or beer—and overloads his blood-vessels and increases the work the heart must do. This kind of hypertrophy develops slowly because the resistance in the circulation is gradual or intermittent. In athletes and in soldiers who are required to march long distances hypertrophy generally occurs. This hypertrophy, if slowly developed by gradual, careful training, is normal and compensatory. In effort too long sustained, especially in those untrained in that kind of effort, and even in the trained if the effort is too long continued, the left ventricle will become dilated and the usual symptoms of that condition occur. Such dilatation is always more or less serious. It may be completely recovered from, and it may not be. Therefore a proper understanding of the physics of the circulation by the medical trainer of young men to decide whether or not one should compete in a prolonged effort, as rowing in a race, for instance, is essential. It is wrong for a single athlete to have an incurable condition occur from competition.

Sometimes simple hypertrophy of the left ventricle occurs from various kinds of conditions that increase the peripheral circulation. It may occur from oversmoking, from the overuse of coffee and tea, from certain kinds of physical labor, or from high-tension mental work. It is a part of the story of hypertension. Many times such patients as well as occasionally trained athletes, and sometimes patients with arteriosclerosis or chronic interstitial nephritis, complain of unpleasant throbbing sensations of the heart. Added to these sensations are a feeling of fulness in the head, flushing of the face, and possibly dizziness—all symptoms not only of hypertension but of too great cardiac activity. Various drugs used to stimulate the heart may cause this condition; when digitalis is given and is not indicated or is given in overdosage these symptoms occur.

The treatment is simply to lower the diet, cause catharsis, give hot baths, stop the tobacco, tea and coffee, stop the drinking of large amounts of liquid at any one time, and administer bromids and perhaps nitroglycerin, when all the symptoms of simple hypertrophy will temporarily, at least, disappear.

If the heart is enlarged from hypertrophy, if it is the right ventricle that is the most hypertrophied, the apex is not only pushed to the left, but the beat may be rather diffuse, as the enlarged right ventricle will prevent the apex from acting close to the surface of the chest. If the left ventricle is the most hypertrophied, the apex is also to the left, but the impact is very decided and the aortic closure is accentuated.

SIMPLE DILATATION

The term "simple dilatation" may be applied to the dilatation of one or both ventricles when there is no valvular lesion and when the condition may not be called broken compensation. Broken compensation has been sufficiently discussed. Dilatation of the heart occurs when there is increased resistance to the outflow of the blood from a ventricle, or when the ventricle is overfilled with blood and the muscular wall is unable to compete with the increased work thrown on it. In other words, it may be weakened by myocarditis or fatty degeneration; or it may be a normal heart that has sustained a strain; or it may be a hypertrophied heart that has become weakened. Heart strain is of frequent occurrence. It occurs in young men from severe athletic effort; it occurs in older persons from some severe muscle strain, and may even occur from so simple an effort as rapid walking by one who is otherwise diseased and whose heart is unable to sustain even this extra work. All of the conditions which have been enumerated as causing simple hypertrophy may have dilatation as a sequence.

The symptoms of heart strain, if the condition is acute, are those of complete prostration, lowered blood-pressure, and a sluggishly, insufficiently acting heart. The heart is found enlarged, the apex-beat diffuse and there may be a systolic blow at the mitral or tricuspid valve. Sometimes, although the patient recognizes that he has hurt himself and strained his heart, he is not prostrated, and the full symptoms do not occur for several hours or perhaps several days, although the patient realizes that he is progressively growing weaker and more breathless.

The treatment of this acute or gradual dilatation is absolute rest, with small doses of digitalis gradually but slowly increased, and when the proper dosage is decided on, administered at that dosage but once a day. Cardiac stimulants should not be given, except when faintness or syncope has occurred, and if strychnin is used, it should be in small doses. The whole care, diet, massage and return to full work should be on the plan advised for a myocarditis following an acute disease. The heart may completely recover its usual powers, but subsequently it is more readily strained again by any thoughtless laborious effort. The patient must be warned as carefully as though he had a valvular lesion and had recovered from a broken compensation, and his life should be regulated accordingly, at least for some months. If he is young, and the heart completely and absolutely recovers, the force of the circulation may remain as strong as ever.

Sometimes the heart strain is not so severe, and after a few hours of rest and quiet the patient regains complete cardiac power and is apparently as well as ever; but for some time subsequently his heart more easily suffers strain.

ACUTE DILATATION OF THE HEART IN ACUTE DISEASE

It has for a long time been recognized that in all acute prolonged illness the heart fails, sooner or later, often without its having been attacked by the disease. The prolonged high temperature causes the heart to beat more rapidly, while the toxins produced by the fever process cause muscle degeneration of the heart or a myocarditis, and at the same time the nutrition of the heart becomes impaired either by improper feeding or by the imperfect metabolism of the food given; hence the heart muscle becomes weakened, and cardiac failure or cardiac relaxation or dilatation occurs.

The specific germ of the disease or the toxin elaborated by the specific germ of the disease may be especially depressant to the heart, as in diphtheria, or the germ may be particularly prone to locate in the heart, as in rheumatism and pneumonia. But all feverish processes, sooner or later, if sufficiently prolonged, cause serious cardiac weakness and more or less dilatation.

Just exactly what changes take place in the muscle fibers of the heart in some of these fevers has not been decided. Whether an albuminous or parenchymatous degeneration of the muscle fibers or a fatty degeneration occurs; whether there is a real myocarditis that always precedes the dilatation, or whether the weakening and loss of muscle fibers or a diminished power of the muscle fibers occurs without inflammation, dilatation of the heart is always a factor to be considered, and frequently occurs in acute disease.

While it is denied that acute dilatation can occur in a sound heart, at the latter end of a serious illness the heart is never sound, and acute dilatation can most readily occur, though fortunately it is generally preventable. When the dilatation occurs suddenly, as indicated by a fluttering heart, a low tension, rapid pulse, dyspnea and perhaps cyanosis with venous stasis in the capillaries, death is imminent, although such patients may be saved by proper aid. Even when the dilatation is slower, as evidenced by a gradually increasing rapidity of the heart and a gradually lowering blood-pressure, and with more evidences of exhaustion, death may occur from such heart failure in spite of all treatment.

Unless a patient dies from accident, as from a hemorrhage, from cerebral pressure or from some organic lesion in acute disease, the physician frequently feels that if he can hold the power and force of the circulation for several hours or days, the patient will recover *from* the disease, for in most acute diseases the patient has a good chance of recovery if his circulation will only hold out until the crisis has occurred or until the disease is ready to end by lysis. Therefore anything during the disease that tends to sustain, nourish, quiet and guard the heart means so much more chance of recovery, whatever else may or may not be done for the disease itself.

The best treatment of dilatation of the heart in acute disease is its prevention, and to prevent it means to recognize the conditions that can cause it. These are:

1. Prolonged high temperature. A short-lived temperature, even if high, is not serious. Prolonged temperature of even 103 F. or more is serious, and even that of 101 is serious if too long continued.

2. Exertion and excitement. Every possible means should be inaugurated to prevent muscular exertion and strain of the patient while in bed. Proper help in lifting and turning the patient should be employed, the bed-pan should be used, proper feeding methods should be adopted, and friends should be excluded so that the patient may not be excited by conversation.

3. Bad feeding. The diet should of course be sufficient for the patient and proper for the disease, but any diet that causes a large amount of gas in the stomach, or tympanites, is harmful to the patient's circulation, to say nothing of any other harm such as indigestion may do. All of the nutriments needed to keep the body in perfect condition should be given to a patient who is ill; in some manner he should receive the proper amounts of iron, salt, calcium, starch, protein, sugar and water.

4. Intestinal sluggishness. This not only means that tympanites should not be allowed, but necessary laxatives should be given. It would be wrong to prostrate a patient with frequent saline purgatives, but the bowels must move at least once every other day, generally better daily; and if the case is one of typhoid fever, they should be moved by some carefully selected laxative, and after the bowels have sufficiently moved, the diar-

rhea should be stopped by 1/10 grain of morphin, and the next day the bowels properly moved again.

5. Depressant drugs. In this age of cardiac failure heart depressants of all types, and especially the synthetic products, should be given only with careful judgment, and never frequently repeated or long continued.

6. Pain. This is one of the most serious depressants a heart has to combat; acute pain must not be allowed, and prolonged subacute pain must be stopped. Even peripheral troublesome irritations must be removed, as tending to wear out a heart that has all of the trouble that it can endure.

7. Insomnia. Nothing rests a heart or recuperates a heart more than sleep. Insomnia and acute disease make a combination that will wear a heart out quicker than any other combination. Sleep, then, must be produced in the best, easiest and safest manner possible.

8. A too speedy return to activity. The convalescence must be prolonged until the heart is able to sustain the work required of it.

The treatment of gradual dilatation in acute disease has been sufficiently discussed under the subject of acute myocarditis. The treatment of acute dilatation is practically the same as the treatment of shock plus whatever treatment must coincidently be given to a patient for the disease with which he is suffering. The treatment of shock will be discussed under a separate heading.

THE HEART IN PNEUMONIA

As pneumonia heads the list of the causes of death in this country, and as the heart fails so quickly, sometimes almost in the beginning, in pneumonia, a special discussion of the management of the heart in this disease is justifiable.

Acute lobar pneumonia may kill a patient in twenty-four or forty-eight hours; he may live for a week and die of heart failure or toxemia, or he may live for

---

5. Brauer: Beitr. z. Klin. d. Tuberk., 1912, xxiii, No. 2.

several weeks and die of cardiac weakness. If he has double pneumonia he may die almost of suffocation. It is to-day just as frequent to see a slowly developing and slowly resolving pneumonia as to see one of the sthenic type that attacks one lobe with a rush, has a crisis in seven, eight or nine days, and then a rapid resolution. In fact the asthenic type, in which different parts of the lung are involved but not necessarily confined to or even equivalent to one lobe, is perhaps the most frequent form of pneumonia.

The serious acute congestion of the lung in sthenic pneumonia in a full-blooded, sturdy person with high tension pulse may be relieved by cardiac sedatives, vasodilators, brisk purging, or by the relaxing effect of antipyretics. Venesection is often the best treatment.

When the sputum almost from the first is tinged with venous blood, or even when the sputum is very bloody, of the prune-juice variety, the heart is in serious trouble, and the right ventricle has generally become weak and possibly dilated. The heart may have been diseased and therefore is unable to overcome the pressure in the lungs during the congestion and consolidation.

There is a great difference in the belief of clinicians as to the best treatment for this condition. It would seem to be a positive indication for digitalis, and good-sized doses of digitalis given correctly, provided always that the preparation of the drug used is active, is good and, many times, efficient treatment. Small doses of strychnin may be of advantage, and camphor may be of value. In the condition described, however, reliance should be placed on digitalis. Later in the disease when the heart begins to fail, perhaps the cause is a myocarditis. In this condition digitalis would not work so well and might do harm. It is quite possible that the difference between digitalis success and digitalis nonsuccess or harm may be as to whether or not a myocarditis is present.

If the expectoration is not of the prune-juice variety and is not more than normally bloody, or in other words, typically pneumonic, and the heart begins to fail, especially if there is no great amount of consolidation, the left ventricle is in trouble as much as the right, if not more. In this case all of the means above described for the prevention of any dilatation of the heart will be means of preventing dilatation from the pneumonia, if possible. The treatment advisable for this gradually failing heart is camphor; strychnin in not too large doses, at the most 1/30 grain hypodermatically once in six hours; many times ergot intramuscularly once in six hours for two or three doses and then once in twelve hours; plenty of fresh air, or perhaps the inhalation of oxygen. Oxygen does not cure pneumonia, but may relieve a dyspnea and aid a heart until other drugs have time to act.

If there is insomnia, morphin in small doses will not only cause sleep, but will not hurt the heart. In the morning hours of the day the value of caffein as a cardiac stimulant and vasocontractor, either in the form of caffein or as black coffee, should be remembered. Strophanthin may be given hypodermatically. One of the greatest cares in the treatment of heart failure in pneumonia should be not to give too many drugs or to do too much.

SHOCK

The treatment of shock will probably always be unsatisfactory as the cause is so varied, and, although circulatory prostration and vasomotor paresis always constitute the acute condition, the physiologic health of the heart and blood-vessels is so varied. The patient in shock has low temperature, low blood-pressure, and a pulse either rapid or slow, but excessively feeble; the face is pale, the surface of the body cold, and there is more or less clammy perspiration; there may be dyspnea and cardiac anxiety, or the patient may hardly breathe.

An acute cause, as terrible pain or hemorrhage, must of course be stopped immediately. There is more or less *anemia* of the brain, and therefore the legs and perhaps the lower part of the body should be elevated. It may even be wise to drive the blood from the legs by Esmarch bandages into the rest of the circulation. As there is always more or less paresis and dilatation of the large veins of the splanchnic system, a tight bandage about the abdomen is of great advantage in raising the blood-pressure to the safety mark.

Strophanthin, given hypodermatically, is valuable as a quick restorative of the heart. 1 Digitalis is so slow that it is of little value in an emergency. Camphor hypodermatically, and hot liquids (nothing is better than black coffee) given by the mouth, are valuable remedies. 2 The camphor may be repeated frequently. Strychnin, the long-used stimulant, should generally be given, but in not too large doses and not too frequently repeated; 1/30 grain hypodermatically is generally a large enough dose; this dose may be repeated in three or four hours, but should ordinarily not be given oftener than once in six hours. An aseptic preparation of ergot given Fair intramuscularly is most efficient in raising the blood-pressure and aiding the heart. Aromatic spirit of Fai ammonia, if at hand, may be the first remedy tried, and one dose of brandy or whisky may do no harm. Alcohol, and however, should not be pushed. dilitat

A most important procedure in all kinds of shock is 3 to surround the patient with dry heat, hot-water bags, and hot flannels; gentle friction of the arms and legs, unless the patient is too exhausted, may be of benefit. A hot-water bag to the heart is always a stimulant. Sometimes friction over the base of the heart in the region of the auricles is of benefit.

If the collapse is not acute and there is gradual profound prostration, or if the patient is improved but still in a serious condition of shock, too energetic measures must not be used; neither should too many drugs be administered, or drugs in too large doses. Absolute quiet and the administration of liquid nourishment in but small amounts at a time are essential.

+ ransient

The hypodermatic administration of epinephrin solutions, 1:10,000, or solutions of pituitary extract, 1:10,000, should be considered; they are often valuable.

V. G.

If the shock occurs in ether or chloroform anesthesia, the vasopressor stimulating effect of inhalations of carbon dioxid gas may be considered, as advised by Henderson.[1]

If the shock is due to hemorrhage and the hemorrhage has ceased, a transfusion of physiologic saline solution is generally indicated. Sometimes transfusion of blood under the same conditions would be still better. Rarely is transfusion indicated in shock from other causes; it often adds to the difficulty rather than improves it. Occasionally if shock is decided to be due to a toxemia, the toxins may be diluted by the withdrawal of a small amount of blood and the transfusion of an equal amount of saline solution.

ACUTE DILATATION OF THE STOMACH

This condition is not well understood, nor is its frequency known, but not a few instances of shock are due to dilatation of this organ. The shock to the heart may be a reflex one through the pneumogastric nerves. It perhaps not infrequently occurs after abdominal operations and is more or less serious, the symptoms being 3 persistent vomiting, upper abdominal distention and collapse. The vomiting is of bloody or coffee-ground material.

Sometimes the ordinary treatment of the collapse and washing out the stomach save the patient; at other times the patient with this series of symptoms dies in spite of all treatment.

? It has been shown that acute dilatation of the stomach may occur in pneumonia, and may be one of the causes of cardiac collapse in pneumonia.

When the condition is diagnosed the treatment would be that of shock plus an abdominal bandage and washing out the stomach with warm solutions, if the patient is not too collapsed, or at any rate the frequent administration of hot water in small quantities.

---

[1] Henderson: Am. Jour. Physiol., February and April, 1909.

Sometimes when the stomach is dilated the pylorus becomes insufficient, and bile regurgitates into the stomach, and is a cause of the profound nausea and vomiting and the subsequent collapse. In these cases not infrequently small doses of dilute hydrochloric acid seem to aid the pylorus to maintain its normal contraction, the regurgitation of bile does not take place, and the stomach may soon acquire a more normal muscle tone. Not infrequently when a stomach is in this kind of trouble and all foods are rejected, and yet the patient seriously needs nourishment, a warm, thin cereal, as oatmeal gruel or something similar, may be retained. Such patients, as has been repeatedly stated, need starch as soon as possible, lest an acidosis develop.

[illegible]

In these vomiting and collapse cases the hypodermatic administration of morphin and atropin will not only stop the vomiting, at least temporarily, but will give necessary rest. The dose of morphin need not be large, and the atropin may prevent nausea from the drug.

ANESTHESIA IN HEART DISEASE

While no physician likes to give an anesthetic to a patient who has valvular disease of the heart, and no surgeon cares to operate on such a patient unless operation is absolutely necessary, still in valvular disease with good compensation the prognosis of either ether or chloroform narcosis is good.

When there are evidences of chronic myocarditis or a history of broken compensation, and the borderline of compensation and dilatation is very narrow, or when there is arteriosclerosis, the danger from an anesthetic and an operation is much greater; it may be serious in fact, and the decision must be made whether the operation is absolutely necessary or not. Under any circumstances it is understood that the anesthetist must be an expert, as there can be no carelessness and nothing but the best of judgment in giving anesthesia in cardiac defect.

The anesthetic to select is a subject for careful decision, as one cannot assert which anesthetic is the best. While chloroform seems occasionally to cause a fatty degeneration of the heart, or if given too rapidly at first may cause sudden death, especially in cardiac weakness, ether has its disadvantages, owing to the increased tension (especially if there is likely to be much muscular or cerebral excitement), and the greater amount of ether that must be given, with the attendant danger to the kidneys, which may have been disturbed from the cardiac conditions. Generally, however, the better method is perhaps to administer first chloroform to the point of producing sleep and then to change to ether, the first mild chloroform narcosis preventing the ether from causing acute stimulation, and ether being better for the operation as it is more of a stimulant.

Some believe that it is better to administer morphin, with perhaps atropin hypodermatically before the anesthesia, and then to use ether. Nitrous oxid gas would be contra-indicated as tending to increase arterial pressure, and therefore endanger a damaged heart; it is a serious danger to damaged blood-vessels.

hyp.

INDEX TO SUBJECTS

PAGE
Alcohol and heart rate.................................. 190
    In broken compensation.............................. 103
    in endocarditis..................................... 52
Alkalies in endocarditis................................ 46
Ammonia in broken compensation.......................... 104
Anesthesia in heart disease............................. 211
Angina pectoris... 137
    management 141
    symptoms 139
Aortic insufficiency 80
    lesions, physics of................................. 70
    regurgitation 80
    stenosis 71, 77
Arhythmia 169
    etiology 171
    treatment 173
Arterial hypertension................................... 151
    etiology 154
    management 157
    symptoms 156
    tracings, interpretation of......................... 11
Atropin in broken compensation.......................... 110
Auricular fibrillation 174
    treatment 176

Baths in endocarditis................................... 51
    in heart diseases 114
Blood-pressure 10
    reducing 160
Bradycardia 178
    symptoms 181
Broken compensation 90
    alcohol in.. 103
    ammonia in.. 104
    atropin in.. 110
    caffein in.. 100
    calcium in.. 108
    camphor in.. 102
    convalescence in.................................... 111
    diet in 91
    digitalis in 95
    elimination in 92
    emergency drugs in.................................. 108
    epinephrin in 109

Broken compensation (continued): PAGE
- ergot in 108
- hygiene in 90
- iodids in 106
- iron in 107
- morphin in 111
- nitroglycerin in 105
- nutritives, cardiac, in 107
- physical measures in 93
- physics of 65
- pituitary extract in 109
- stimulants in 102
- strophanthus in 99
- strychnin in 101
- thyroid extract in 107
- tonics in 95
- treatment of 90
- vasodilators in 105

Bromids and chloral in endocarditis 48

Caffein and heart rate 192
- in broken compensation 100

Calcium in broken compensation 108
Camphor in broken compensation 102
Cardiac: See Heart.
Cardiovascular-renal disease 163
- diagnosis 165

Children, heart disease in 123
Chloral and bromids in endocarditis 48
Coronary sclerosis 130
- pathology 131
- symptoms 132
- treatment 135

Degenerations 130
Diet in broken compensation 91
- in endocarditis 50
- in heart diseases 114

Digitalis in broken compensation 95
Dilatation, acute, of the heart, in acute diseases 203
- simple 202

Elimination in broken compensation 92
Endocarditis 38
- acute 38
  - alcohol in 52
  - alkalies in 46
  - baths in 51
  - bromids and chloral in 48
  - cardiac drugs in 50
  - diet in 50
  - iron in 47
  - laxatives in 50
  - opium in 47

Endocarditis (continued): PAGE
  pathology of ... 40
  prevention of ... 49
  prognosis and convalescence in ... 52
  strychnin in ... 52
  symptoms and signs ... 43
  treatment ... 44
 chronic ... 57
  etiology ... 58
  pathology ... 60
  treatment ... 60
 malignant ... 54
  diagnosis ... 55
  treatment ... 56
 ulcerative ... 54
Epinephrin in broken compensation ... 109
Ergot in broken compensation ... 108
Exercise in broken compensation ... 94

Fatty degeneration ... 85
Fatty heart ... 86

Heart, anesthesia in diseases of the ... 211
 acute attack ... 84
  drugs in ... 87
  management of ... 85
 block ... 148
  treatment ... 150
 diseases, classification of ... 16
 diseases in children ... 123
 diseases, prophylaxis of ... 15
 diseases, symptoms and signs ... 12
 during pregnancy ... 127
 in pneumonia ... 206
 in shock ... 208
 rate, disturbances of ... 169
Hydrotherapy in broken compensation ... 93
Hygiene in broken compensation ... 90
Hyperthyroidism ... 186
Hypertrophy, simple ... 200

Intestinal putrefaction ... 189
Iodids in broken compensation ... 106
Iron in broken compensation ... 107
  in endocarditis ... 47

Laxatives in endocarditis ... 50

Mitral insufficiency ... 76
  regurgitation ... 76
  stenosis ... 73
  stenosis, physics of ... 68
Morphin in broken compensation ... 111
Myocardial disturbances ... 29

PAGE
Myocarditis, acute 28
  chronic 31
    management 33
    prognosis 35
    symptoms and signs 32

Nauheim baths 119
Nitroglycerin in broken compensation 105
Nutritives, cardiac, in broken compensation 107

Oertel treatment of chronic diseases 118
Opium in endocarditis 47

Pericarditis, acute 17
    adherent 25
    convalescence in 25
    exudate in 21
    pain in, stopping the 20
    symptoms and signs 17
    treatment 18
Pituitary extract in broken compensation 109
Pneumonia, heart in 206
Pregnancy, heart during 127
Pseudo-angina 144
    treatment 147
Pulmonary insufficiency 83
    regurgitation 83
    stenosis 83
Pulse tracings, clinical interpretation of 8

Resort treatment of chronic heart diseases 117

Schott treatment 119
Shock, heart in 208
Stimulants, cardiac 102
Stokes-Adams disease 148
Strophanthus in broken compensation 99
Strychnin in broken compensation 101
    in endocarditis 52

Tachycardia, paroxysmal 183
    management 184
Thyroid extract in broken compensation 107
Tobacco and heart rate 194
Tonics in broken compensation 95
Toxic disturbances and heart rate 188
Tricuspid insufficiency 71, 81
    stenosis 82

Valves, chronic diseases of 62
    pathologic physiology of 62
Valvular lesions, chronic, symptomatology and treatment 73
Vasodilators 105

LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on
or before the date last stamped below.

JUN 11 '28

JUL 11 '30

AUG -9 1930

JAN 30 1931

MAR 25 1931

L681 Osborne, O.T.
081 Disturbances of the
1913 heart. 54037

NAME	DATE DUE
O. F. Johnson	JUN 11 28
Room 309.	NOV 16 1928
Henry Specker	JUL 11
M. C. Baldwin	JUN 2 1930
	JUL 11 30
	AUG 9 1930
J. H. Padg[illegible]	JAN 26 1931
Lewis G. Jacobs Jr	MAR 24 1931